

ANALYSIS & PDE

Volume 10

No. 1

2017

ANNE-LAURE DALIBARD AND DAVID GÉRARD-VARET

NONLINEAR BOUNDARY LAYERS FOR ROTATING FLUIDS

NONLINEAR BOUNDARY LAYERS FOR ROTATING FLUIDS

ANNE-LAURE DALIBARD AND DAVID GÉRARD-VARET

We investigate the behaviour of rotating incompressible flows near a nonflat horizontal bottom. In the flat case, the velocity profile is given explicitly by a simple linear ODE. When bottom variations are taken into account, it is governed by a nonlinear PDE system, with far less obvious mathematical properties. We establish the well-posedness of this system and the asymptotic behaviour of the solution away from the boundary. In the course of the proof, we investigate in particular the action of pseudodifferential operators in nonlocalized Sobolev spaces. Our results extend an older paper of Gérard-Varet (*J. Math. Pures Appl.* (9) **82**:11 (2003), 1453–1498), restricted to periodic variations of the bottom, using the recent linear analysis of Dalibard and Prange (*Anal. & PDE* **7**:6 (2014), 1253–1315).

1. Introduction

The general concern of this paper is the effect of rough walls on fluid flows, in a context where the rough wall has very little structure. This effect is important in several problems, like transition to turbulence or drag computation. For instance, understanding the connection between roughness and drag is crucial for microfluidics, because friction at solid boundaries is a major factor of energy loss in microchannels. This issue has been much studied over recent years, through both theory and experiments [Lauga et al. 2007; Bocquet and Barrat 2007]. Conclusions are ambivalent. On the one hand, rough surfaces may increase the friction area, and thus enstrophy dissipation. On the other hand, recent experiments have shown that rough hydrophobic surfaces may lead to drag decrease: air bubbles can be trapped in the humps of the roughness, generating some slip [Vinogradova and Yakubov 2006; Ybert et al. 2007].

Mathematically, these problems are often tackled by a homogenization approach. Typically, one considers Stokes equations over a rough plate, modelled by an oscillating boundary of small wavelength and amplitude:

$$\Gamma^\varepsilon : \quad x_3 = \varepsilon\gamma(x_1/\varepsilon, x_2/\varepsilon), \quad \varepsilon \ll 1, \quad (1-1)$$

where the function $\gamma = \gamma(y_1, y_2)$ describes the roughness pattern. Within this formalism, the understanding of roughness-induced effects comes down to an asymptotic problem, as $\varepsilon \rightarrow 0$. The point is to derive effective boundary conditions at the flat plate Γ^0 , retaining in this boundary condition an averaged effect of the roughness. We refer to the works [Achdou et al. 1998a; 1998b; 1998c; Amirat et al. 2001; Jäger and Mikelić 2001; 2003; Neuss et al. 2006; Bresch and Milisic 2010; Mikelić et al. 2013] on this topic. In all of these works, a restrictive hypothesis is made, namely periodicity of the roughness pattern γ . This hypothesis simplifies greatly the construction of the so-called boundary layer corrector, describing the

MSC2010: primary 35Q30; secondary 35Q86.

Keywords: fluid mechanics, geophysical fluids, Ekman layers, boundary layers.

small-scale variations of the flow near the boundary. This corrector is an analogue of the cell corrector in classical homogenization of heterogeneous media.

The main point and difficulty is the mathematical study of the boundary layer equations, which are satisfied formally by the boundary layer corrector. When γ is periodic in y_1, y_2 , the solution of the boundary layer system is itself sought periodic, so that well-posedness and qualitative properties of the system are easy to determine. When the periodicity structure is relaxed, and replaced by general ergodicity properties, the analysis is still possible, but much more involved, as shown in [Basson and Gérard-Varet 2008; Gérard-Varet 2009; Gérard-Varet and Masmoudi 2010]. A key feature of these articles is the linearity of the boundary layer system: after the rescaling $y = x/\varepsilon$, it is governed by Stokes equations in the boundary layer domain

$$\Omega_{\text{bl}} = \{y : y_3 > \gamma(y_1, y_2)\}. \quad (1-2)$$

It thus reads

$$\begin{cases} -\Delta v + \nabla p = 0 & \text{in } \Omega_{\text{bl}}, \\ \operatorname{div} v = 0 & \text{in } \Omega_{\text{bl}}, \\ v|_{\partial\Omega_{\text{bl}}} = \phi \end{cases} \quad (1-3)$$

for some Dirichlet boundary data ϕ that has no decay as y_1, y_2 go to infinity, but no periodic structure. As a consequence, spaces of infinite energy, such as H_{uloc}^s , form a natural functional setting for such equations.

A natural challenge is to extend this type of analysis to nonlinear systems. This is the goal of the present paper. Namely, we will study a nonlinear boundary layer system that describes a rotating fluid near a rough boundary. The dynamics of rotating fluid layers are relevant in the context of geophysical flows, for which the Earth's rotation plays a dominant role. The system under consideration reads

$$\begin{cases} v \cdot \nabla v + \nabla p + e \times v - \Delta v = 0 & \text{in } \Omega_{\text{bl}}, \\ \operatorname{div} v = 0 & \text{in } \Omega_{\text{bl}}, \\ v|_{\partial\Omega_{\text{bl}}} = \phi. \end{cases} \quad (1-4)$$

These are the incompressible Navier–Stokes equations written in a rotating frame, which is the reason for the extra Coriolis force $e \times u$, where $e = e_3 = (0, 0, 1)^t$. The equations in (1-4) can be obtained through an asymptotics of the full rotating fluid system

$$\operatorname{Ro}(\partial_t u + u \cdot \nabla u) + e \times u - E\Delta u = 0, \quad \operatorname{div} u = 0, \quad (1-5)$$

where Ro and E are the so-called Rossby and Ekman numbers. These parameters are small in many applications. In the vicinity of the rough boundary (1-1), and in the special case where

$$E \sim \varepsilon^2, \quad \operatorname{Ro} \sim \varepsilon, \quad (1-6)$$

it is natural to look for an asymptotic behaviour of the type

$$u^\varepsilon(t, x) \sim v(t, x_1, x_2, x/\varepsilon),$$

where $v = v(t, x_1, x_2, y)$, $y \in \Omega_{\text{bl}}$. Injecting this ansatz in (1-5) yields the first two equations in the system (1-4), where the “slow variables” (t, x_1, x_2) are only parameters and are eluded.

The main goal of this paper is to construct a solution v of system (1-4), under no structural assumption on γ . We shall moreover provide information on the behaviour of v away from the boundary. We will in this way generalize [Gérard-Varet 2003] by the second author in which periodic roughness was considered. See also [Gérard-Varet and Dormy 2006]. Before stating the main difficulties and results of our study, several remarks are in order:

(1) The choice of the scaling (1-6), which leads to the derivation of the boundary layer system, may seem peculiar. It is, however, the richest possible, as it retains all terms in the equation for the boundary layer. All other scaling would provide a degeneracy of system (1-4).

(2) In the flat case, that is, for the roughness profile $\gamma = 0$, and for $\phi = (\phi_1, \phi_2, 0)$, with ϕ_1, ϕ_2 independent of y , the solution of (1-5) is explicitly given in complex form by

$$(v_1 + iv_2)(y) = (\phi_1 + i\phi_2) \exp(-(1+i)y_3/\sqrt{2}), \quad v_3 = 0. \quad (1-7)$$

This profile, sometimes called the Ekman spiral, solves the linear ODE

$$e \times v - \partial_3^2 v = 0.$$

Considering roughness turns this linear ODE into a nonlinear PDE, and as we will see, changes drastically the properties of the solution.

(3) Rather than the Dirichlet condition $v|_{\partial\Omega_{\text{bl}}} = \phi$, some slightly different settings could be considered:

- One could for instance prescribe a homogeneous Dirichlet condition $v|_{\partial\Omega_{\text{bl}}} = 0$, and add a source term with enough decay in y_3 . This would correspond to a localized forcing of the boundary layer.
- One could replace the Dirichlet condition by a Navier condition, that is, a condition of the type

$$D(u)n \times n|_{\partial\Omega_{\text{bl}}} = f, \quad u \cdot n|_{\partial\Omega_{\text{bl}}} = 0,$$

with $D(u)$ the symmetric part of ∇u , and n the normal unit vector at the boundary. For instance, one could think of (1-1) as modelling an oscillating free surface, under the rigid lid approximation. In this context, the Navier condition would model a wind forcing, and the boundary layer domain would model the water below the free surface (changing the direction of the vertical axis). We refer to [Pedlosky 1987] for some similar modelling, and to [Casado-Díaz et al. 2003; Bucur et al. 2008; Bonnivard and Bucur 2012; Dalibard and Gérard-Varet 2011] for the treatment of such Navier condition. As shown in those papers, some hypothesis on the nondegeneracy of the roughness is necessary to the mathematical analysis.

However, our analysis does not extend to the important case of an inhomogeneous Dirichlet condition at infinity, which models a boundary layer driven by an external flow. For linear systems, one can in general lift this Dirichlet data at infinity, and recover the case of a Dirichlet data at the bottom boundary, like in (1-3). But for our nonlinear system (1-4), this lift would lead to the introduction of an additional drift term in the momentum equation, which would break down its rotational invariance.

2. Statement of the results

Our main result is a well-posedness theorem for the boundary layer system (1-4), where ϕ is a given boundary data, with no decay tangentially to the boundary, and satisfying $\phi \cdot n|_{\partial\Omega_{\text{bl}}} = 0$. As usual in the theory of steady Navier–Stokes equations, the well-posedness will be obtained under a smallness hypothesis. We first introduce, for any unbounded $\Omega \subset \mathbb{R}^d$, the spaces

$$L_{\text{uloc}}^2(\Omega) = \left\{ f : \sup_{k \in \mathbb{Z}^d} \int_{B(k,1) \cap \Omega} |f|^2 < +\infty \right\},$$

$$\text{and for all } m \geq 0, \quad H_{\text{uloc}}^m(\Omega) = \{ f : \partial^\alpha f \in L_{\text{uloc}}^2(\Omega) \forall \alpha \leq m \}.$$

These spaces are of course Banach spaces when endowed with their natural norms.

Theorem 1. *Let γ be bounded and Lipschitz and Ω_{bl} be defined as in (1-2). There exists $\delta_0, C > 0$, such that for all $\phi \in H_{\text{uloc}}^2(\partial\Omega_{\text{bl}})$ satisfying $\phi \cdot n|_{\partial\Omega_{\text{bl}}} = 0$ and $\|\phi\|_{H_{\text{uloc}}^2} \leq \delta_0$ system (1-4) has a unique solution (v, p) with*

$$(1 + y_3)^{1/3} v \in H_{\text{uloc}}^1(\Omega_{\text{bl}}), \quad (1 + y_3)^{1/3} p \in L_{\text{uloc}}^2(\Omega_{\text{bl}}),$$

and

$$\|(1 + y_3)^{1/3} v\|_{H_{\text{uloc}}^1} + \|(1 + y_3)^{1/3} p\|_{L_{\text{uloc}}^2} \leq C \|\phi\|_{H_{\text{uloc}}^2}.$$

This theorem generalizes the result of [Gérard-Varet 2003], dedicated to the case of periodic roughness pattern γ . In this case, the analysis is much easier, as the solution v of (1-4) is itself periodic in y_1, y_2 . Through standard arguments, one can then build a solution v satisfying

$$\int_{\mathbb{T}^2} \int_{y_3 > \gamma(y_1, y_2)} |\nabla v|^2 < +\infty.$$

Moreover, one can establish exponential decay estimates for v as y_3 goes to infinity. This exponential decay is related to the periodicity in the horizontal variables, which provides a Poincaré inequality for functions with zero mean in x_1 . When the periodicity assumption is removed, one expects the exponential convergence to be no longer true: this has been notably discussed in [Gérard-Varet and Masmoudi 2010; Prange 2013] in the context of the Laplace or the Stokes equation near a rough wall. It is worth noting that in such context, the convergence can be arbitrarily slow. In fact, there is in general no convergence when no ergodicity assumption on γ is made. A remarkable feature of our theorem for rotating flows is that decay to zero persists, despite the nonlinearity, and without any ergodicity assumption on γ . We emphasize that this decay comes from the rotation term. However, exponential decay is replaced by polynomial decay, with rate $O(y_3^{-1/3})$ for v .

Let us comment on the difficulties associated with Theorem 1. Of course, the first issue is that the data ϕ does not decay as (y_1, y_2) goes to infinity, so that the solution v is not expected to decay in the horizontal directions. If Ω_{bl} were replaced by

$$\Omega_{\text{bl}}^M := \{y : M > y_3 > \gamma(y_1, y_2)\}, \quad M > 0,$$

together with a Dirichlet condition at the upper boundary, one could build a solution v in $H_{\text{uloc}}^1(\Omega_{\text{bl}}^M)$, adapting ideas of Ladyženskaya and Solonnikov [1980] on Navier–Stokes flows in tubes. Among those ideas, an important one is to obtain an a priori differential inequality on the local energy

$$E(t) := \int_{\{|(y_1, y_2)| \leq t\}} \int_{\{M > y_3 > \gamma(y_1, y_2)\}} |\nabla v|^2.$$

Such a differential inequality, known in the literature as a *Saint-Venant estimate*, appeared previously in other contexts; see for instance [Wheeler and Horgan 1976; Wheeler et al. 1975]. Namely, one shows an inequality of the type

$$E(t) \leq C_M(E'(t) + E'(t)^{3/2} + t^2).$$

However, the derivation of this differential inequality relies on the Poincaré inequality between two planes, or in other words on the fact that Ω_{bl}^M has a bounded direction. For the boundary layer domain Ω_{bl} , this is no longer true, and no a priori bound can be obtained in this way. Moreover, contrary to what happens for the Laplace equation, one cannot rely on maximum principles to get an L^∞ bound.

Under a periodicity assumption on γ , one can restrict the domain to the periodic slab

$$\{y : (y_1, y_2) \in \mathbb{T}^2, y_3 > \gamma(y_1, y_2)\}.$$

In this manner, one has again a domain with a bounded direction (horizontal rather than vertical). One can establish again Saint-Venant estimates leading to the exponential decay mentioned above. It allows one to prove well-posedness of the boundary layer system. However, this approach does not work in our framework, where no structure is assumed on the roughness profile γ .

For the Stokes boundary layer flow

$$-\Delta v + \nabla p = 0, \quad \operatorname{div} v = 0 \quad \text{in } \Omega_{\text{bl}}, \quad v|_{\partial\Omega_{\text{bl}}} = v_0, \quad (2-1)$$

this problem is overcome in [Gérard-Varet and Masmoudi 2010] by N. Masmoudi and the second author. The main idea there is to get back to the domain Ω_{bl}^M by imposing a so-called transparent boundary condition at $y_3 = M$. This transparency condition involves the Stokes analogue of the Dirichlet-to-Neumann operator, and, despite its nonlocal nature (contrary to the Dirichlet condition), allows then to apply the method of Solonnikov. We refer to [Gérard-Varet and Masmoudi 2010] for more details.¹ Of course, the use of an explicit transparent boundary condition at $y_3 = M$ is possible because v satisfies a homogeneous Stokes equation in the half-space $\{y_3 > M\}$, which gives access to explicit formulas.

Such simplification does not occur in the context of our rotating flow system: in particular, the main issue is the quasilinear term $u \cdot \nabla u$ in system (1-4), in contrast with previous linear studies. In fact, even without this convective term, the analysis is not easy. In other words, the Coriolis–Stokes problem

$$\begin{cases} e \times v + \nabla p - \Delta v = 0 & \text{in } \Omega_{\text{bl}}, \\ \operatorname{div} v = 0 & \text{in } \Omega_{\text{bl}}, \\ v|_{\partial\Omega_{\text{bl}}} = \phi \end{cases} \quad (2-2)$$

¹Actually, [Gérard-Varet and Masmoudi 2010] is concerned with the 2D case. For adaptation to 3D, we refer to [Dalibard and Prange 2014].

already raises difficulties. For instance, to use a strategy based on a transparent boundary condition, one needs to construct the solution of the Dirichlet problem in a half-space for the Stokes–Coriolis operator, when the Dirichlet data has uniform local bounds. But contrary to the Stokes case, there is no easy integral representation. Still, such a linear problem was tackled in the recent paper [Dalibard and Prange 2014] by the first author and C. Prange. To solve the Dirichlet problem, they use a Fourier transform in variables y_1, y_2 , leading to accurate formulas. The point is then to be able to translate information on the Fourier side to uniform local bounds on v . This requires careful estimates, as spaces like L^2_{uloc} are defined through truncations in space, which are not so suitable for a Fourier treatment. Similar difficulties arise in [Alazard et al. 2016], devoted to water waves equations in locally uniform spaces.

The linear study [Dalibard and Prange 2014] is a starting point for our study of the nonlinear system (1-4), but we will need many refined estimates, combined with a fixed point argument. More precisely, the outline of the paper is the following.

- Section 3, the main section of the paper, will be devoted to the system

$$\begin{cases} e \times v + \nabla p - \Delta v = \operatorname{div} F & \text{in } \{y_3 > M\}, \\ \operatorname{div} v = 0 & \text{in } \{y_3 > M\}, \\ v|_{y_3=M} = v_0. \end{cases} \quad (2-3)$$

The data v_0 and F will have no decay in horizontal variables (y_1, y_2) . The source term F , which is reminiscent of $u \otimes u$, will decay typically like $|y_3|^{-2/3}$ as y_3 goes to infinity. This exponent is coherent with the decay of u given in Theorem 1. The point will be to establish a priori estimates on a solution v of (2-3), with no decay in (y_1, y_2) , decaying like $|y_3|^{-1/3}$ at infinity. Functional spaces will be specified in due course.

- On the basis of previous a priori estimates, we will show well-posedness of the system

$$\begin{cases} v \cdot \nabla v + e \times v + \nabla p - \Delta v = 0 & \text{in } \{y_3 > M\}, \\ \operatorname{div} v = 0 & \text{in } \{y_3 > M\}, \\ v|_{y_3=M} = v_0 \end{cases} \quad (2-4)$$

for small enough boundary data v_0 (again, in a functional space to be specified). This will be done in the first subsection of Section 4.

- Finally, through the next subsections of Section 4, we will establish Theorem 1. The solution v of (1-4) will be constructed with the help of a mapping $\mathcal{F} = \mathcal{F}(\psi, \phi)$, defined in the following way:

- (1) First, we will introduce the solution (v^-, p^-) of

$$\begin{cases} v^- \cdot \nabla v^- + e \times v^- + \nabla p^- - \Delta v^- = 0 & \text{in } \Omega_{\text{bl}}^M, \\ \operatorname{div} v^- = 0 & \text{in } \Omega_{\text{bl}}^M, \\ v^-|_{\partial\Omega_{\text{bl}}} = \phi, \\ \Sigma(v^-, p^-)e_3|_{y_3=M} = \psi, \end{cases} \quad (2-5)$$

where $\Sigma(v, p) = \nabla v - (p + \frac{1}{2}|v|^2)\operatorname{Id}$. Note that a quadratic term $\frac{1}{2}|v|^2$ is added to the usual Newtonian tensor in order to handle the nonlinearity.

(2) Then, we will introduce the solution (v^+, p^+) of (2-4), with $v_0 := v^-|_{y_3=M}$.

(3) Eventually, we will define $\mathcal{F}(\psi, \phi) := \Sigma(v^+, p^+)e_3|_{y_3=M} - \psi$.

The point will be to show that for small enough ϕ , the equation $\mathcal{F}(\psi, \phi) = 0$ has a solution ψ , knowing that $\mathcal{F}(0, 0) = 0$. This will be obtained via the inverse function theorem (using the linear analysis of [Dalibard and Prange 2014]). For such ψ , the field v defined by v^\pm over $\{\pm y_3 > M\}$ will be a solution of (1-4). Indeed, v is always continuous at $y_3 = M$ by the definition of v^+ , while the condition $\mathcal{F}(\psi, \phi) = 0$ means that the normal component of the stress tensor $\Sigma(v, p)$ is also continuous at $y_3 = M$.

3. Stokes–Coriolis equations with source

A central part of the work is the analysis of system (2-3). For simplicity, we take $M = 0$. The case without source term ($F = 0$) was partially analyzed in [Dalibard and Prange 2014], but we will establish new estimates, notably related to low frequencies. Let us emphasize that the difficulty induced by low frequencies already appeared in Proposition 2.1 on page 6 of the above work, even in the case of classical Sobolev data: in such case, some cancellation of the Fourier transform $\hat{v}_{0,3}$ at frequency $\xi = 0$ was assumed. We make a similar hypothesis here. The main theorem of the section is:

Theorem 2. *Let $m \in \mathbb{N}$, $m \gg 1$. Let $v_0 \in H_{\text{uloc}}^{m+1}(\mathbb{R}^2)$ with third component satisfying $v_{0,3} = \partial_1 v_1^* + \partial_2 v_2^*$, with v_1^*, v_2^* in $L_{\text{uloc}}^2(\mathbb{R}^2)$. Let $F \in H_{\text{loc}}^m(\mathbb{R}_+^3)$ such that $(1 + y_3)^{2/3} F \in H_{\text{uloc}}^m(\mathbb{R}_+^3)$. There exists a unique solution v of system (2-3) such that*

$$\|(1 + y_3)^{1/3} v\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}_+^3)} \leq C \left(\|v_0\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|(1 + y_3)^{2/3} F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)} \right) \quad (3-1)$$

for a universal constant C .

Prior to the proof of the theorem, several simplifying remarks are in order:

- Obviously, uniqueness comes down to showing that if $F = 0$ and $v_0 = 0$, the only solution v of (2-3) such that $(1 + y_3)^{1/3} v \in H_{\text{uloc}}^m(\mathbb{R}_+^3)$ is $v = 0$. This result follows from [Dalibard and Prange 2014, Proposition 2.1], in which even a larger functional space was considered. Hence, the key statement our theorem is the existence of a solution satisfying the estimate (3-1).
- In order to show existence of such a solution, we can assume $v_{0,1}, v_{0,2}, v^* := (v_1^*, v_2^*)$ and F to be smooth and compactly supported (resp. in \mathbb{R}^2 and \mathbb{R}_+^3). Indeed, let us introduce

$$\begin{aligned} (v_{0,1}^n, v_{0,2}^n, v^{*,n})(y_1, y_2) &:= \chi((y_1, y_2)/n) \rho^n \star (v_{0,1}, v_{0,2}, v^*)(y_1, y_2), \\ F^n(y) &:= \tilde{\chi}(y/n) \tilde{\rho}^n(y) \star F(y), \end{aligned}$$

where $\chi \in C_c^\infty(\mathbb{R}^2)$, $\tilde{\chi} \in C_c^\infty(\mathbb{R}^3)$ are 1 near the origin, and $\rho^n, \tilde{\rho}^n$ are approximations of unity. These functions are smooth, compactly supported, and satisfy

$$\begin{aligned} \|(v_{0,1}^n, v_{0,2}^n)\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2)} &\leq C \|(v_{0,1}, v_{0,2})\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2)}, \\ \|v^{*,n}\|_{H_{\text{uloc}}^{m+2}(\mathbb{R}^2)} &\leq C \|v^*\|_{H_{\text{uloc}}^{m+2}(\mathbb{R}^2)}, \\ \|(1 + y_3)^{2/3} F^n\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)} &\leq C \|(1 + y_3)^{2/3} F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)} \end{aligned}$$

for a universal constant C . Moreover, $(v_{0,1}^n, v_{0,2}^n)$, $v^{*,n}$ and F^n converge strongly to $(v_{0,1}, v_{0,2})$, v^* and F in $H^{m+1}(K)$, $H^{m+2}(K)$ and $H^m(K')$ respectively for any compact set K of \mathbb{R}^2 and any compact set K' of \mathbb{R}_+^3 . Now, assume that for all $n \in \mathbb{N}$, there exists a solution v^n corresponding to the data $v_{0,1}^n, v_{0,2}^n, v^{*,n}$, and F^n , for which we can get the estimate

$$\|(1 + y_3)^{1/3} v^n\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}_+^3)} \leq C (\|(v_{0,1}^n, v_{0,2}^n)\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2)} + \|v^{*,n}\|_{H_{\text{uloc}}^{m+2}(\mathbb{R}^2)} + \|(1 + y_3)^{2/3} F^n\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)})$$

for a universal constant C . Then,

$$\|(1 + y_3)^{1/3} v^n\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}_+^3)} \leq C' (\|(v_{0,1}, v_{0,2})\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2)} + \|v^*\|_{H_{\text{uloc}}^{m+2}(\mathbb{R}^2)} + \|(1 + y_3)^{2/3} F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)})$$

for a universal constant C' . We can then extract a subsequence weakly converging to some v , which is easily seen to satisfy (2-3) and (3-1).

• Finally, if $v_{0,1}, v_{0,2}, v^*$ and F are smooth and compactly supported, the existence of a solution v of (2-3) can be obtained by standard variational arguments. More precisely, one can build a function v such that

$$\begin{aligned} \int_{\mathbb{R}_+^3} |\nabla v|^2 &\leq C (\|F\|_{L^2(\mathbb{R}^2)} + \|v_0\|_{H^{1/2}(\mathbb{R}^2)}), \\ \int_{\mathbb{R}^2 \times \{y_3 < a\}} |v|^2 &\leq C_a (\|F\|_{L^2(\mathbb{R}^2)} + \|v_0\|_{H^{1/2}(\mathbb{R}^2)}) \quad \forall a > 0. \end{aligned}$$

Higher-order derivatives are then controlled by elliptic regularity. Hence, the whole problem is to establish the estimate (3-1) for such a solution.

We are now ready to tackle the proof of [Theorem 2](#). We forget temporarily about the boundary condition and focus on the equations

$$e \times v + \nabla p - \Delta v = \text{div } F, \quad \text{div } v = 0 \quad \text{in } \mathbb{R}_+^3, \quad (3-2)$$

Our goal is to construct some particular solution of these equations, satisfying for some large enough m ,

$$\|(1 + z)^{1/3} v\|_{L^\infty} \leq C \|(1 + z)^{2/3} F\|_{L^\infty(H_{\text{uloc}}^m)}. \quad (3-3)$$

We will turn to the solution of the whole system (2-3) in a second step.

3.1. Orr–Sommerfeld formulation. To handle (3-2), we rely on a formulation similar to Orr and Sommerfeld's rewriting of Navier–Stokes. Namely, we wish to express this system in terms of v_3 and $\omega := \partial_1 v_2 - \partial_2 v_1$. First, we apply ∂_2 to the first line, $-\partial_1$ to the second line, and combine to obtain

$$\partial_3 v_3 + \Delta \omega = s_3 := \partial_2 f_1 - \partial_1 f_2, \quad \text{with } f := \text{div } F = \left(\sum_j \partial_j F_{ij} \right)_i. \quad (3-4)$$

Similarly, we apply $\partial_1 \partial_3$ to the first line of (3-2), $\partial_2 \partial_3$ to the second line, and $-(\partial_1^2 + \partial_2^2)$ to the third line. Combining the three, we are left with

$$-\partial_3 \omega + \Delta^2 v_3 = s_\omega := \partial_1 \partial_3 f_1 + \partial_2 \partial_3 f_2 - (\partial_1^2 + \partial_2^2) f_3. \quad (3-5)$$

From ω and v_3 , one recovers the horizontal velocity components v_1, v_2 using the system

$$\partial_1 v_1 + \partial_2 v_2 = -\partial_3 v_3, \quad \partial_1 v_2 - \partial_2 v_1 = \omega.$$

We are led to the (so far formal) expressions

$$\begin{aligned} v_1 &= (\partial_1^2 + \partial_2^2)^{-1}(-\partial_3 \partial_1 v_3 - \partial_2 \omega), \\ v_2 &= (\partial_1^2 + \partial_2^2)^{-1}(-\partial_3 \partial_2 v_3 + \partial_1 \omega). \end{aligned} \tag{3-6}$$

Our goal is to construct a solution (v_3, ω) of (3-4)–(3-5), by means of an integral representation. Since the vertical variable will play a special role in this construction, we will denote it by z instead of y_3 : $y = (y_1, y_2, z)$. We write (3-4)–(3-5) in the compact form

$$L(D, \partial_z)V = S, \quad V := \begin{pmatrix} v_3 \\ \omega \end{pmatrix}, \quad S := \begin{pmatrix} s_3 \\ s_\omega \end{pmatrix}, \quad D := \frac{1}{i}(\partial_1, \partial_2),$$

where $L(D, \partial_z)$ is a Fourier multiplier in variables x_1, x_2 associated with

$$L(\xi, \partial_z) := \begin{pmatrix} \partial_z & (\partial_z^2 - |\xi|^2) \\ (\partial_z^2 - |\xi|^2)^2 & -\partial_z \end{pmatrix}.$$

We will look for a solution of the form

$$V(\cdot, z) = \int_0^{+\infty} G(D, z - z')S(\cdot, z') dz' + V_h, \tag{3-7}$$

where:

- $G(D, z)$ is a matrix Fourier multiplier, whose symbol $G(\xi, z)$ is the fundamental solution over \mathbb{R} of $L(\xi, \partial_z)$ for any $\xi \in \mathbb{R}^2$:

$$L(\xi, \partial_z)G(\xi, z) = \delta_{z=0} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

- V_h is a solution of the homogeneous equation. The purpose of the addition of V_h is to ensure the decay of the solution V . More details will be given in due course.

3.1.1. Construction of the Green function. We start with the construction of the fundamental solution $G(\xi, z)$. Away from $z = 0$, it should satisfy the homogeneous system, which requires one to understand the kernel of the operator $L(\xi, \partial_z)$. This kernel is a combination of elements of the form $e^{\lambda z}V$, where λ is a root of the characteristic equation

$$\det L(\xi, \lambda) = 0, \quad \text{i.e., } -\lambda^2 - (\lambda^2 - |\xi|^2)^3 = 0, \tag{3-8}$$

and V is an associated “eigenelement”, meaning a nonzero vector in $\ker L(\xi, \lambda)$. A careful study of the characteristic equation was carried out recently in [Dalibard and Prange 2014]. Notice that (3-8) can be seen as an equation of degree three on $Y = \lambda^2 - |\xi|^2$ (with negative discriminant). Using Cardano’s formula gives access to explicit expressions. The roots can be written as $\pm\lambda_1(\xi)$, $\pm\lambda_2(\xi)$ and $\pm\lambda_3(\xi)$, where $\lambda_1 \in \mathbb{R}_+$, λ_2, λ_3 have positive real parts, $\lambda_1 \in \mathbb{R}$, $\overline{\lambda_2} = \lambda_3$, $\text{Im}\lambda_2 > 0$. The λ_i are continuous

functions of ξ (see [Remark 4](#) below for more). The above work also provides their asymptotic behaviour at low and high frequencies. This behaviour will be very important to establish our estimates.

Lemma 3 [[Dalibard and Prange 2014](#), Lemma 2.4]. *As $\xi \rightarrow 0$, we have*

$$\lambda_1(\xi) = |\xi|^3 + O(|\xi|^5), \quad \lambda_2(\xi) = e^{i\pi/4} + O(|\xi|^2), \quad \lambda_3(\xi) = e^{-i\pi/4} + O(|\xi|^2).$$

As $\xi \rightarrow \infty$, we have

$$\lambda_1(\xi) = |\xi| - \frac{1}{2}|\xi|^{-1/3} + O(|\xi|^{-5/3}),$$

$$\lambda_2(\xi) = |\xi| - \frac{1}{2}j^2|\xi|^{-1/3} + O(|\xi|^{-5/3}), \quad \lambda_3(\xi) = |\xi| - \frac{1}{2}j|\xi|^{-1/3} + O(|\xi|^{-5/3}), \quad \text{where } j = \exp(2i\pi/3).$$

Remark 4. We insist that λ_2 and λ_3 are distinct and have a positive real part for all values of ξ , whereas $\lambda_1 \neq 0$ for $\xi \neq 0$. Moreover, it can be easily checked that λ_i^2 is a C^∞ function of $|\xi|^2$ for $i = 1, \dots, 3$. Using the fact that λ_2 and λ_3 never vanish or merge, while λ_1 vanishes for $\xi = 0$ only, we deduce that λ_2, λ_3 are C^∞ functions of $|\xi|^2$, and that $\lambda_1(\xi) = |\xi|^3 \Lambda_1(\xi)$, where $\Lambda_1 \in C^\infty(\mathbb{R}^2)$, $\Lambda_1(0) = 1$ and Λ_1 does not vanish on \mathbb{R}^2 .

Regarding the eigenelements, an explicit computation shows that for all $i = 1, \dots, 3$,

$$V_i^\pm := \begin{pmatrix} 1 \\ \pm\Omega_i \end{pmatrix} \quad \text{and} \quad \Omega_i := \frac{-\lambda_i}{\lambda_i^2 - |\xi|^2} \quad \text{satisfy } L(\xi, \pm\lambda_i)V_i^\pm = 0. \quad (3-9)$$

We can now determine G ; our results are summarized in [Lemma 5](#) below. We begin with its first column $G_1 = \begin{pmatrix} G_{11} \\ G_{21} \end{pmatrix}$, a solution of $L(\xi, \partial_z)G_1 = \delta \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. As explained above, for $z \neq 0$, we know $G_1(\xi, z)$ is a linear combination of $e^{\pm\lambda_i z} V_i^\pm$. Furthermore, we want to avoid any exponential growth of G as $z \rightarrow \pm\infty$. Thus G_1 should be of the form

$$G_1 = \begin{cases} \sum_{i=1}^3 A_i^+ e^{-\lambda_i z} V_i^-, & z > 0, \\ \sum_{i=1}^3 A_i^- e^{\lambda_i z} V_i^+, & z < 0. \end{cases}$$

We now look at the jump conditions at $z = 0$. For $f = f(z)$, recall that $[f]|_{z=z'} := f(z'^+) - f(z'^-)$ denotes the jump of f at z' . Since

$$\begin{cases} (\partial_z^2 - |\xi|^2)^2 G_{11} - \partial_z G_{21} = 0, \\ \partial_z G_{11} + (\partial_z^2 - |\xi|^2) G_{21} = \delta_{z=0}, \end{cases}$$

we infer that

$$[G_{21}]|_{z=0} = 0, \quad [\partial_z G_{21}]|_{z=0} = 1, \quad [\partial_z^k G_{11}]|_{z=0} = 0, \quad k = 0, \dots, 3.$$

This yields a linear system of six equations on the coefficients A_i^\pm . One finds $A_i := A_i^+ = -A_i^-$, and the system

$$\sum_i \lambda_i \Omega_i A_i = \frac{1}{2}, \quad \sum_i A_i = 0, \quad \sum_i \lambda_i^2 A_i = 0.$$

Note that

$$\sum_i \lambda_i \Omega_i A_i = - \sum_i \frac{\lambda_i^2}{\lambda_i^2 - |\xi|^2} A_i = - \sum_i \frac{|\xi|^2}{\lambda_i^2 - |\xi|^2} A_i$$

taking into account the second equality. Hence, we find

$$\begin{pmatrix} |\xi|^2/(\lambda_1^2 - |\xi|^2) & |\xi|^2/(\lambda_2^2 - |\xi|^2) & |\xi|^2/(\lambda_3^2 - |\xi|^2) \\ 1 & 1 & 1 \\ \lambda_1^2 & \lambda_2^2 & \lambda_3^2 \end{pmatrix} \begin{pmatrix} A_1 \\ A_2 \\ A_3 \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} \\ 0 \\ 0 \end{pmatrix}.$$

The determinant of the matrix is

$$D_1 := |\xi|^2 D,$$

where

$$D := \begin{vmatrix} 1/(\lambda_1^2 - |\xi|^2) & 1/(\lambda_2^2 - |\xi|^2) & 1/(\lambda_3^2 - |\xi|^2) \\ 1 & 1 & 1 \\ \lambda_1^2 & \lambda_2^2 & \lambda_3^2 \end{vmatrix}.$$

After a few computations, we find that

$$D_1 = |\xi|^2 (\lambda_2^2 - \lambda_1^2) (\lambda_3^2 - \lambda_1^2) \left(\frac{1}{(\lambda_1^2 - |\xi|^2)(\lambda_2^2 - |\xi|^2)} - \frac{1}{(\lambda_1^2 - |\xi|^2)(\lambda_3^2 - |\xi|^2)} \right), \quad (3-10)$$

and

$$A_1 = -\frac{1}{2D_1} (\lambda_3^2 - \lambda_2^2), \quad A_2 = -\frac{1}{2D_1} (\lambda_1^2 - \lambda_3^2), \quad A_3 = -\frac{1}{2D_1} (\lambda_2^2 - \lambda_1^2). \quad (3-11)$$

Computations for the second column G_2 of G are similar. It is of the form

$$G_2 = \begin{cases} \sum_{i=1}^3 B_i^+ e^{-\lambda_i z} V_i^-, & z > 0, \\ \sum_{i=1}^3 B_i^- e^{\lambda_i z} V_i^+, & z < 0, \end{cases}$$

with jump conditions

$$[\partial_z^k G_{22}]|_{z=0} = 0, \quad k = 0, 1, \quad [\partial_z^k G_{12}]|_{z=0} = 0, \quad k = 0, \dots, 2, \quad [\partial_z^3 G_{12}]|_{z=0} = 1.$$

We find $B_i := B_i^+ = B_i^-$ and the system

$$\begin{pmatrix} \Omega_1 & \Omega_2 & \Omega_3 \\ \lambda_1 & \lambda_2 & \lambda_3 \\ \lambda_1^3 & \lambda_2^3 & \lambda_3^3 \end{pmatrix} \begin{pmatrix} B_1 \\ B_2 \\ B_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ -\frac{1}{2} \end{pmatrix}.$$

The determinant of the matrix is now $D_2 := -\lambda_1 \lambda_2 \lambda_3 D$, and

$$\begin{aligned} B_1 &= \frac{\lambda_2 \lambda_3}{2D_2} \left(\frac{1}{\lambda_2^2 - |\xi|^2} - \frac{1}{\lambda_3^2 - |\xi|^2} \right), & B_2 &= \frac{\lambda_1 \lambda_3}{2D_2} \left(\frac{1}{\lambda_3^2 - |\xi|^2} - \frac{1}{\lambda_1^2 - |\xi|^2} \right), \\ B_3 &= \frac{\lambda_1 \lambda_2}{2D_2} \left(\frac{1}{\lambda_1^2 - |\xi|^2} - \frac{1}{\lambda_2^2 - |\xi|^2} \right). \end{aligned} \quad (3-12)$$

This concludes the construction of the matrix G . We sum up our results in the following lemma, in which we also give the asymptotic behaviours of the coefficients A_i , B_i , V_i^\pm and of G as $\xi \rightarrow 0$ and $|\xi| \rightarrow \infty$. The latter follow from [Lemma 3](#) and [Remark 4](#) and are left to the reader.

Lemma 5. *We have*

$$G_1 = \begin{cases} \sum_{i=1}^3 A_i e^{-\lambda_i z} V_i^-, & z > 0, \\ -\sum_{i=1}^3 A_i e^{\lambda_i z} V_i^+, & z < 0, \end{cases} \quad G_2 = \begin{cases} \sum_{i=1}^3 B_i e^{-\lambda_i z} V_i^-, & z > 0, \\ \sum_{i=1}^3 B_i e^{\lambda_i z} V_i^+, & z < 0, \end{cases}$$

where

$$V_i^\pm = \begin{pmatrix} 1 \\ \mp \lambda_i / (\lambda_i^2 - |\xi|^2) \end{pmatrix}$$

and where A_i and B_i are defined by (3-11) and (3-12) respectively.

Asymptotic behaviour:

• For $|\xi| \gg 1$, there exists $N > 0$ such that $A_i, B_i, \Omega_i = O(|\xi|^N)$ for $i = 1, \dots, 3$, and $|\Omega_i| \gtrsim |\xi|^{-N}$. As a consequence, $G(\xi, z) = O(|\xi|^N)$ for all z .

• As $\xi \rightarrow 0$, we have

$$\begin{aligned} A_i(\xi) &\rightarrow \bar{A}_i \in \mathbb{C}^*, \quad i = 1, \dots, 3, \\ B_1(\xi) &\sim \frac{\bar{B}_1}{|\xi|}, \quad \bar{B}_1 \in \mathbb{C}^*, \quad B_i(\xi) \rightarrow \bar{B}_i \in \mathbb{C}^*, \quad i = 2, 3, \\ \Omega_1 &\sim \bar{\Omega}_1 |\xi|, \quad \bar{\Omega}_1 \in \mathbb{C}^*, \quad \Omega_i(\xi) \rightarrow \bar{\Omega}_i \in \mathbb{C}^*, \quad i = 2, 3. \end{aligned} \quad (3-13)$$

More precisely, we can write, for instance,

$$B_1(\xi) = \frac{\bar{B}_1}{|\xi|} \beta_1(\xi) \quad \forall \xi \in \mathbb{R}^2$$

for some function $\beta_1 \in C^\infty(\mathbb{R}^2)$ such that $\beta_1(0) = 1$. Similar statements hold for the other coefficients.

It follows that

$$G(\xi, z) = \begin{pmatrix} O(1) & O(|\xi|^{-1}) \\ O(1) & O(1) \end{pmatrix}$$

as $|\xi| \rightarrow 0$ for all $z \in \mathbb{R}$.

3.1.2. Construction of the homogeneous correction. We will see rigorously below that the field

$$V_G(\cdot, z) := \int_0^{+\infty} G(D, z-z') S(\cdot, z') dz' = \int_0^{+\infty} \mathcal{F}_{\xi \rightarrow (y_1, y_2)}^{-1} (G(\cdot, z-z') \mathcal{F}_{(y_1, y_2) \rightarrow \xi} S(\cdot, z')) dz' \quad (3-14)$$

is well-defined and satisfies (3-4)–(3-5). However, the corresponding velocity field does not have a good decay with respect to z . This is the reason for the additional field V_h in formula (3-7). To be more specific, let us split the source term S into $S(z') = S^0(z') + \partial_{z'} S^1(z') + \partial_{z'}^2 S^2(z')$, with

$$S^0(z') := \begin{pmatrix} \partial_2(\partial_1 F_{11} + \partial_2 F_{12}) - \partial_1(\partial_1 F_{21} + \partial_2 F_{22}) \\ -(\partial_1^2 + \partial_2^2)(\partial_1 F_{31} + \partial_2 F_{32}) \end{pmatrix} \quad (3-15)$$

and

$$\begin{aligned} S^1(z') &:= \begin{pmatrix} \partial_2 F_{13} - \partial_1 F_{23} \\ \partial_1(\partial_1 F_{11} + \partial_2 F_{12}) + \partial_2(\partial_1 F_{21} + \partial_2 F_{22}) - (\partial_1^2 + \partial_2^2)F_{33} \end{pmatrix}, \\ S^2(z') &:= \begin{pmatrix} 0 \\ \partial_1 F_{13} + \partial_2 F_{23} \end{pmatrix}. \end{aligned} \quad (3-16)$$

Roughly, the idea is that

$$V(\cdot, z) := \int_0^{+\infty} (G(D, z - z')S^0(z') + \partial_z G(D, z - z')S^1(z') + \partial_z^2 G(D, z - z')S^2(z')) dz'$$

has a better decay. Using the fact that

$$\partial_z G(D, z - z') = -\partial_{z'} G(D, z - z'),$$

we see that going from V_G to V is possible through integrations by parts in the variable z' , which generates boundary terms. We recall that the jump of $G(D, z - z')$ at $z = z'$ is zero, and that

$$[\partial_z G(D, z - z')] \Big|_{z=z'} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

On the other hand, the first component of S^2 is zero, so that the jump of $\partial_z G_{21}$ at $z = z'$ is not involved in the two integrations by parts of $\partial_z^2 G(D, z - z')S^2(z')$. Formal computations eventually lead to

$$\begin{aligned} V_h(\cdot, z) &:= V(\cdot, z) - V_G(\cdot, z) \\ &= -[G(D, z - z')(S^1(\cdot, z') + \partial_z S^2(\cdot, z'))]_0^{+\infty} + [\partial_z G(D, z - z')S^2(\cdot, z')]_0^{+\infty} \\ &= G(D, z)(S^1(\cdot, 0) + \partial_{z'} S^2(\cdot, 0)) - \partial_z G(D, z)S^2(\cdot, 0). \end{aligned}$$

Back to the expression of the Green function, we get

$$\begin{aligned} V_h(\cdot, z) &= -(\sum_i A_i e^{-\lambda_i z} V_i^- - \sum_i B_i e^{-\lambda_i z} V_i^-)(S^1(\cdot, 0) + \partial_{z'} S^2(\cdot, 0)) \\ &\quad + (\sum_i A_i \lambda_i e^{-\lambda_i z} V_i^- - \sum_i B_i \lambda_i e^{-\lambda_i z} V_i^-)S^2(\cdot, 0). \end{aligned} \quad (3-17)$$

It is a linear combination of terms of the form $e^{-\lambda_i z} V_i^-$, and therefore satisfies the homogeneous Orr–Sommerfeld equations. Hence, V is (still formally) a solution of (3-4)–(3-5).

We now need to put these formal arguments on rigorous grounds. As mentioned after [Theorem 2](#), there is no loss of generality assuming that F is smooth and compactly supported.

Lemma 6. *Let F be smooth and compactly supported. The formula (3-7), with V_h given by (3-17), defines a solution $V = (v_3, \omega)^t$ of (3-4)–(3-5) satisfying*

$$V \in L_{\text{loc}}^\infty(\mathbb{R}_+, H^m(\mathbb{R}^2)), \quad |D|^{-1}\omega \in L_{\text{loc}}^\infty(\mathbb{R}_+, H^m(\mathbb{R}^2)) \quad \text{for any } m.$$

Proof. Let us first show that the integral term V_G (see (3-14)) satisfies the properties of the lemma. The main point is to show that for any $z, z' \geq 0$, the function

$$J_{z,z'} : \xi \rightarrow G(\xi, z - z')\hat{S}(\xi, z') \quad \text{belongs to} \quad L^2((1 + |\xi|^2)^{m/2} d\xi) \times L^2(|\xi|^{-1}(1 + |\xi|^2)^{m/2} d\xi)$$

for all m . Therefore, we recall that $\hat{F} = \hat{F}(\xi, z')$ is in the Schwartz class with respect to ξ , smooth and compactly supported in z' . Also, $G(\xi, z - z')$ is smooth in $\xi \neq 0$ (see [Remark 4](#)), and continuous in z, z' . It implies that $J_{z,z'}$ is smooth in $\xi \neq 0$, continuous in z, z' . It remains to check its behaviour at high and low frequencies.

- At high frequencies ($|\xi| \gg 1$), from [Lemma 5](#), it is easily seen that $J_{z,z'}$ is bounded by

$$|J_{z,z'}(\xi)| \leq C|\xi|^N \sum_{k=0}^2 |\partial_{z'}^k \hat{F}(\xi, z')|$$

for some N . As \hat{F} and its z' -derivatives are rapidly decreasing in ξ , it will belong to any L^2 with polynomial weight.

- At low frequencies ($\xi \sim 0$), one can check that $|\hat{S}(\xi, z')| \leq C|\xi|$. Hence, using again the bounds derived in [Lemma 5](#),

$$G(\xi, z - z') \hat{S}(\xi, z') = \begin{pmatrix} O(1) \\ O(|\xi|) \end{pmatrix}.$$

The result follows.

From there, by standard arguments, V_G defines a continuous function of z with values in $H^m(\mathbb{R}^2) \times |D|^{-1}H^m(\mathbb{R}^2)$ for all m . Moreover, a change of variable gives

$$V_G(\cdot, z) = \int_0^{+\infty} G(D, z') S(\cdot, z - z') dz'.$$

By the smoothness of S , we deduce that V_G is smooth in z with values in the same space. The fact that it satisfies (3-4)–(3-5) comes of course from the properties of the Green function G , and is classical. We leave it to the reader.

To conclude the proof of the lemma, we still have to consider the homogeneous correction V_h . Again, V_h is smooth in $\xi \neq 0$ and z . Thanks to the properties of F , it is decaying fast as $|\xi|$ goes to infinity. Moreover, from the asymptotics above, one can check that $V_h = \begin{pmatrix} O(1) \\ O(|\xi|) \end{pmatrix}$ for $|\xi| \ll 1$. Finally, as its Fourier transform is a linear combination of $e^{-\lambda_i(\xi)z} V_i^-(\xi)$, it satisfies (3-4)–(3-5) without source. \square

Let us stress that, with the same kind of arguments, one can justify the integration by parts mentioned above, and write

$$V(\cdot, z) := \int_0^{+\infty} \sum_{k=0}^2 \partial_z^k G(D, z - z') S^k(z') dz'. \quad (3-18)$$

We will now try to derive the estimate (3-3), starting from this formulation.

3.1.3. Main estimate. By [Lemma 6](#), we know that formula (3-7) (or equivalently (3-18)) defines a solution V of (3-4)–(3-5). Our main goal in this section is to establish that V obeys inequality (3-3). Our main ingredient will be:

Lemma 7. *Let $\chi = \chi(\xi) \in C_c^\infty(\mathbb{R}^2)$, and $P = P(\xi) \in C^\infty(\mathbb{R}^2 \setminus \{0\})$ defined by*

$$P(\xi) = p_k(\xi) |\xi|^{\alpha-k} Q(\xi)$$

near $\xi = 0$, with p_k a homogeneous polynomial in ξ_1, ξ_2 of degree k , $\alpha > 0$, and $Q \in C^\infty(\mathbb{R}^2)$. Assume furthermore that $\alpha - k \geq -2$. For $\underline{v}_0 \in L^1_{\text{uloc}}(\mathbb{R}^2)$, we define $u^i = u^i(y_1, y_2, z)$ by

$$u^i(\cdot, z) := \chi(D)P(D)e^{-\lambda_i(D)z}\underline{v}_0. \quad (3-19)$$

Then, there exists C and $\delta > 0$ independent of \underline{v}_0 such that

$$\|e^{\delta z}u^2\|_{L^\infty(\mathbb{R}_+^3)} + \|e^{\delta z}u^3\|_{L^\infty(\mathbb{R}_+^3)} \leq C\|\underline{v}_0\|_{L^1_{\text{uloc}}}.$$

Moreover, there exists C and $\delta > 0$ independent of \underline{v}_0 such that

$$\|(1+z)^{\frac{\alpha}{3}}u^1\|_{L^\infty(\mathbb{R}_+^3)} \leq C\|\underline{v}_0\|_{L^1_{\text{uloc}}}.$$

Remark 8. Showing that the definition (3-19) makes sense is part of the proof of the lemma. Namely, it is shown that for any $z > 0$, the kernel

$$K(x_1, x_2, z) := \mathcal{F}_{\xi \rightarrow (x_1, x_2)}^{-1}(\chi(\xi)P(\xi)e^{-\lambda_i(\xi)z})$$

defines an element of $L^1(\mathbb{R}^2)$. In particular, (3-19) is appropriate: $u^i = K(\cdot, z) \star \underline{v}_0$ defines (at least) an L^1_{uloc} function as the convolution of functions of L^1 and L^1_{uloc} .

We refer to [Appendix A](#) for a proof. [Lemma 7](#) is the source of the asymptotic behaviour of the solution v of (1-4). As always in this type of boundary layer problem, the asymptotic behaviour is given by low frequencies, corresponding to the cut-off χ . In particular, the decay is given by the characteristic root $\lambda_1(\xi)$, which vanishes at $\xi = 0$.

Proof of estimate (3-3). We distinguish between low and high frequencies.

Low frequencies. We introduce some $\chi = \chi(\xi) \in C_c^\infty(\mathbb{R}^2)$ equal to 1 near $\xi = 0$. We consider

$$V^b = \int_{\mathbb{R}^+} \sum_{k=0}^2 I^k(\cdot, z, z') dz', \quad (3-20)$$

$$I^k(\cdot, z, z') := \chi(D)\partial_z^k G(D, z - z')S^k(\cdot, z').$$

In what follows, we write

$$S^k = (s_3^k, s_\omega^k)^t \quad \text{and} \quad I^k = (I_3^k, I_\omega^k)^t.$$

We will use the following fact, which is a straightforward consequence of (3-15)–(3-16): \widehat{s}_3^0 and \widehat{s}_ω^1 are homogeneous of degree 2 and \widehat{s}_ω^0 is homogeneous of degree 3, while \widehat{s}_3^1 and \widehat{s}_ω^2 are homogeneous of degree 1.

Study of I^0 . We find

$$I_3^0(\cdot, z, z') = \text{sgn}(z - z')\chi(D) \sum A_i(D)e^{-\lambda_i(D)|z - z'|}s_3^0(\cdot, z') + \chi(D) \sum B_i(D)e^{-\lambda_i(D)|z - z'|}s_\omega^0(\cdot, z'),$$

$$I_\omega^0(\cdot, z, z') = -\chi(D) \sum A_i(D)\Omega_i(D)e^{-\lambda_i(D)|z - z'|}s_3^0(\cdot, z') \\ - \text{sgn}(z - z')\chi(D) \sum B_i(D)\Omega_i(D)e^{-\lambda_i(D)|z - z'|}s_\omega^0(\cdot, z').$$

We also have

$$\begin{aligned} \partial_z I_3^0(\cdot, z, z') &= -\chi(D) \sum A_i(D) \lambda_i(D) e^{-\lambda_i(D)|z-z'|} s_3^0(\cdot, z') \\ &\quad - \operatorname{sgn}(z-z') \chi(D) \sum B_i(D) \lambda_i(D) e^{-\lambda_i(D)|z-z'|} s_\omega^0(\cdot, z'). \end{aligned}$$

We note that $\widehat{s}_3^0(\xi, z')$ and $\widehat{s}_\omega^0(\xi, z')$ are products of components of $\widehat{F}(\xi, z')$ by homogeneous polynomials of degrees 2 and 3 respectively in ξ . Using the asymptotic behaviours derived in [Lemma 5](#) together with [Lemma 7](#), we deduce

$$\begin{aligned} \|I_3^0(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{2/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\ \|I_\omega^0(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{1+|z-z'|} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\ \left\| \frac{D}{|D|^2} I_\omega^0(\cdot, z, z') \right\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{2/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\ \left\| \frac{D}{|D|^2} \partial_z I_3^0(\cdot, z, z') \right\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{4/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}. \end{aligned} \tag{3-21}$$

The last two bounds will be useful when estimating the horizontal velocity components through [\(3-6\)](#). We insist that $\partial_z I_3^0$ has a better behaviour than I_3^0 , because there is an extra factor $\lambda_1(D)$ in front of A_1 and B_1 , which gives a higher degree of homogeneity at low frequencies for the term in $\exp(-\lambda_1(D)z)$. This is why we can apply $D/|D|^2$ to that term. As for the terms in $\exp(-\lambda_i(D)z)$ for $i=2, 3$, there is no singularity near $\xi=0$ when we apply $D/|D|^2$ because of the homogeneity of degrees 2 and 3 in $\widehat{s}_3^0(\xi, z')$ and $\widehat{s}_\omega^0(\xi, z')$ respectively.

Study of I^1 . We find

$$\begin{aligned} I_3^1(\cdot, z, z') &= -\chi(D) \sum A_i(D) \lambda_i(D) e^{-\lambda_i(D)|z-z'|} s_3^1(\cdot, z') \\ &\quad - \operatorname{sgn}(z-z') \chi(D) \sum B_i(D) \lambda_i(D) e^{-\lambda_i(D)|z-z'|} s_\omega^1(\cdot, z'), \end{aligned}$$

$$\begin{aligned} I_\omega^1(\cdot, z, z') &= \operatorname{sgn}(z-z') \chi(D) \sum A_i(D) \lambda_i(D) \Omega_i(D) e^{-\lambda_i(D)|z-z'|} s_3^1(\cdot, z') \\ &\quad + \chi(D) \sum B_i(D) \lambda_i(D) \Omega_i(D) e^{-\lambda_i(D)|z-z'|} s_\omega^1(\cdot, z'), \end{aligned}$$

and also

$$\begin{aligned} \partial_z I_3^1(\cdot, z, z') &= \operatorname{sgn}(z-z') \chi(D) \sum A_i(D) (\lambda_i(D))^2 e^{-\lambda_i(D)|z-z'|} s_3^1(\cdot, z') \\ &\quad + \chi(D) \sum B_i(D) (\lambda_i(D))^2 e^{-\lambda_i(D)|z-z'|} s_\omega^1(\cdot, z'). \end{aligned}$$

Thanks to the derivation of the Green function with respect to z , an extra factor $\lambda_1(D)$ appears together with $A_1(D)$ or $B_1(D)$. This provides a higher degree of homogeneity in $|\xi|$ at low frequencies. It compensates for the loss of homogeneity of S^1 compared to S^0 . More precisely, we note that $\widehat{s}_3^1(\xi, z')$ and $\widehat{s}_\omega^1(\xi, z')$ are products of components of $\widehat{F}(\xi, z')$ by homogeneous polynomials of degrees 1 and 2

respectively in ξ . We also get

$$\begin{aligned}
\|I_3^1(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{4/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\
\|I_\omega^1(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{5/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\
\left\| \frac{D}{|D|^2} I_\omega^1(\cdot, z, z') \right\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{4/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\
\left\| \frac{D}{|D|^2} \partial_z I_3^1(\cdot, z, z') \right\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^2} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}.
\end{aligned} \tag{3-22}$$

Study of I^2 . We find

$$\begin{aligned}
I_3^2(\cdot, z, z') &= \text{sgn}(z-z')\chi(D) \sum A_i(D)(\lambda_i(D))^2 e^{-\lambda_i(D)|z-z'|} s_3^2(\cdot, z') \\
&\quad + \chi(D) \sum B_i(D)(\lambda_i(D))^2 e^{-\lambda_i(D)|z-z'|} s_\omega^2(\cdot, z'),
\end{aligned}$$

as well as

$$\begin{aligned}
I_\omega^2(\cdot, z, z') &= -\chi(D) \sum A_i(D)(\lambda_i(D))^2 \Omega_i(D) e^{-\lambda_i(D)|z-z'|} s_3^2(\cdot, z') \\
&\quad - \text{sgn}(z-z')\chi(D) \sum B_i(D)(\lambda_i(D))^2 \Omega_i(D) e^{-\lambda_i(D)|z-z'|} s_\omega^2(\cdot, z'),
\end{aligned}$$

and

$$\begin{aligned}
\partial_z I_3^2(\cdot, z, z') &= -\chi(D) \sum A_i(D)(\lambda_i(D))^3 e^{-\lambda_i(D)|z-z'|} s_3^2(\cdot, z') \\
&\quad - \text{sgn}(z-z')\chi(D) \sum B_i(D)(\lambda_i(D))^3 e^{-\lambda_i(D)|z-z'|} s_\omega^2(\cdot, z').
\end{aligned}$$

This time, $s_3^2 = 0$ and $\widehat{s_\omega^2}$ is homogeneous of degree 1. We get as before that

$$\begin{aligned}
\|I_3^2(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^2} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\
\|I_\omega^2(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{7/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\
\left\| \frac{D}{|D|^2} I_\omega^2(\cdot, z, z') \right\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^2} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\
\left\| \frac{D}{|D|^2} \partial_z I_3^2(\cdot, z, z') \right\|_{L^\infty(\mathbb{R}^2)} &\leq \frac{C}{(1+|z-z'|)^{8/3}} \|F(\cdot, z')\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}.
\end{aligned} \tag{3-23}$$

Combining (3-21)–(3-23), we find

$$\begin{aligned}
\|v_3^b(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C \int_0^{+\infty} \frac{1}{(1+|z-z'|)^{2/3}} \frac{1}{(1+z')^{2/3}} dz' \|(1+z'^{2/3})F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}, \\
\|\omega^b(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C \int_0^{+\infty} \frac{1}{1+|z-z'|} \frac{1}{(1+z')^{2/3}} dz' \|(1+z'^{2/3})F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))},
\end{aligned}$$

and

$$\begin{aligned} \left\| \frac{D}{|D|^2} \omega^b(\cdot, z) \right\|_{L^\infty(\mathbb{R}^2)} &\leq C \int_0^{+\infty} \frac{1}{(1+|z-z'|)^{2/3}} \frac{1}{(1+z')^{2/3}} dz' \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}, \\ \left\| \frac{D}{|D|^2} \partial_z v_3^b(\cdot, z) \right\|_{L^\infty(\mathbb{R}^2)} &\leq C \int_0^{+\infty} \frac{1}{(1+|z-z'|)^{4/3}} \frac{1}{(1+z')^{2/3}} dz' \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}. \end{aligned}$$

We deduce that (see [Lemma 16](#) in [Appendix B](#))

$$\begin{aligned} \|v_3^b(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C(1+z)^{-1/3} \|(1+z^{2/3})F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}, \\ \|\omega^b(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C(1+z)^{-2/3} \ln(2+z) \|(1+z^{2/3})F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}, \end{aligned} \quad (3-24)$$

and

$$\begin{aligned} \left\| \frac{D}{|D|^2} \omega^b(\cdot, z) \right\|_{L^\infty(\mathbb{R}^2)} &\leq C(1+z)^{-1/3} \|(1+z^{2/3})F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}, \\ \left\| \frac{D}{|D|^2} \partial_z v_3^b(\cdot, z) \right\|_{L^\infty(\mathbb{R}^2)} &\leq C(1+z)^{-2/3} \ln(2+z) \|(1+z^{2/3})F\|_{L^\infty(L^1_{\text{uloc}}(\mathbb{R}^2))}. \end{aligned} \quad (3-25)$$

High frequencies. To obtain the estimate (3-3), we still have to control the high frequencies

$$V^\# = \int_{\mathbb{R}^+} \sum_{k=0}^2 J^k(\cdot, z, z') dz', \quad J^k(\cdot, z, z') := (1 - \chi(D)) \partial_z^k G(D, z - z') S^k(\cdot, z'). \quad (3-26)$$

Instead of [Lemma 7](#), we shall use this (see [Appendix A](#) for a proof):

Lemma 9. *Let $\chi \in C_c^\infty(\mathbb{R}^2)$, with $\chi = 1$ in a ball $B_r := B(0, r)$ for some $r > 0$. Let $P = P(\xi) \in C_b^3(\mathbb{R}^2 \setminus B_r)$. For $\underline{v}_0 = \underline{v}_0(y_1, y_2) \in H_{\text{uloc}}^N(\mathbb{R}^2)$, $N \in \mathbb{N}$, we define $u^i = u^i(y_1, y_2, z)$ by*

$$u^i(\cdot, z) := (1 - \chi(D)) P(D) e^{-\lambda_i(D)z} \underline{v}_0. \quad (3-27)$$

Then, for N large enough and $\delta > 0$ small enough,

$$\|e^{\delta z} u^1\|_{L^\infty(\mathbb{R}_+^3)} + \|e^{\delta z} u^2\|_{L^\infty(\mathbb{R}_+^3)} + \|e^{\delta z} u^3\|_{L^\infty(\mathbb{R}_+^3)} \leq C \|\underline{v}_0\|_{H_{\text{uloc}}^N(\mathbb{R}^2)}.$$

Remark 10. As in the proof of [Lemma 7](#), part of the proof of [Lemma 9](#) gives a meaning to (3-27). In particular, it is shown that for n large enough, and any $z > 0$, the kernel

$$K_n(x_1, x_2, z) := \mathcal{F}^{-1} \left((1 + |\xi|^2)^{-n} (1 - \chi(\xi)) P(\xi) e^{-\lambda_i(\xi)z} \right)$$

belongs to $L^1(\mathbb{R}^2)$ so that $u^i = K_n \star ((1 - \Delta)^n \underline{v}_0)$ defines at least an element of L^2_{uloc} as the convolution of functions in L^1 and L^2_{uloc} (assuming $N \geq 2n$).

The analysis is simpler than for low frequencies. From (3-26), (3-15)–(3-16) and [Lemma 5](#), we decompose the components of J^k for $k = 0, 1, 2$ into terms of the form

$$(1 - \chi(D)) R(D) e^{-\lambda_i(D)|z-z'|} \partial_1^{a_1} \partial_2^{a_2} F_{jl},$$

where F_{jl} are components of our source term F , $a_1, a_2 = 0, 1, 2$ with $1 \leq a_1 + a_2 \leq 3$, and $R(D)$ is of the form

$$R(D) = \mathcal{R}(\lambda_1(D), \lambda_2(D), \lambda_3(D), D)$$

for some rational expression $\mathcal{R} = \mathcal{R}(\lambda_1, \lambda_2, \lambda_3, \xi)$. Considering the behaviour of $\lambda_i(\xi)$ at infinity (see [Lemma 7](#) and [Remark 4](#)), it can be easily seen that $|\xi|^{-2n} R(\xi) \in C_b^3(\mathbb{R}^2 \setminus B_r)$ for some n large enough. Thus, we can apply [Lemma 9](#) with

$$P(\xi) = |\xi|^{-2n} R(\xi), \quad \underline{v}_0 = (\partial_1^2 + \partial_2^2)^n \partial_1^{a_1} \partial_2^{a_2} F_{jl}(\cdot, z').$$

This shows that for m large enough ($m = N + 2n + 3$),

$$\|J^k(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} \leq C e^{-\delta|z-z'|} \|F(\cdot, z')\|_{H_{\text{uloc}}^m(\mathbb{R}^2)}. \quad (3-28)$$

Also, up to taking a larger m , one can check that

$$\|\partial_z J^k(\cdot, z, z')\|_{L^\infty(\mathbb{R}^2)} \leq C e^{-\delta|z-z'|} \|F(\cdot, z')\|_{H_{\text{uloc}}^m(\mathbb{R}^2)}. \quad (3-29)$$

We deduce from (3-28)–(3-29) that for m large enough

$$\begin{aligned} \|V^\#(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} + \|\partial_z V^\#(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C \int_0^{+\infty} e^{-\delta|z-z'|} (1+z')^{-2/3} dz' \|(1+z)^{2/3} F\|_{L^\infty(H_{\text{uloc}}^m)} \\ &\leq C (1+z)^{-2/3} \|(1+z)^{2/3} F\|_{L^\infty(H_{\text{uloc}}^m)}. \end{aligned} \quad (3-30)$$

Together with (3-24), this inequality implies the estimate (3-3). \square

Together with (3-25), inequality (3-30) further yields

$$\left\| (1+z)^{1/3} \frac{D}{|D|^2} \partial_z v_3 \right\|_{L^\infty(\mathbb{R}_+^3)} + \left\| (1+z)^{1/3} \frac{D}{|D|^2} \omega \right\|_{L^\infty(\mathbb{R}_+^3)} \leq C \|(1+z)^{2/3} F\|_{L^\infty(H_{\text{uloc}}^m)}. \quad (3-31)$$

3.2. Proof of [Theorem 2](#). In the last section, we have constructed a particular solution of (3-4)–(3-5) satisfying (3-3) and (3-31); in the rest of this section, we denote this particular solution as $V^P = (v_3^P, \omega^P)^t$. The bound (3-31) implies in particular that

$$\left\| (1+z)^{1/3} (v_1^P, v_2^P) \right\|_{L^\infty(\mathbb{R}_+^3)} \leq C \|(1+z)^{2/3} F\|_{L^\infty(H_{\text{uloc}}^m)}, \quad (3-32)$$

where v_1^P, v_2^P are recovered from v_3^P, ω^P through formula (3-6).

We still need to make the connection with the solution of (2-3). Following the discussion after [Theorem 2](#), for smooth and compactly supported data, such a solution exists, and the point is to establish (3-1). We introduce

$$\underline{v} := v - v^P, \quad \underline{\omega} = \omega - \omega^P.$$

Functions \underline{v}_3 and $\underline{\omega}$ satisfy the homogeneous version of the Orr–Sommerfeld equations:

$$\partial_3 \underline{v}_3 + \Delta \underline{\omega} = 0, \quad -\partial_3 \underline{\omega} + \Delta^2 \underline{v}_3 = 0. \quad (3-33)$$

These equations are completed by the boundary conditions

$$\begin{aligned} \underline{v}_3|_{z=0} &= v_{0,3} - v_3^P|_{z=0}, \quad \partial_z \underline{v}_3|_{z=0} = -\partial_1(v_{0,1} - v_1^P) - \partial_2(v_{0,2} - v_2^P), \\ \underline{\omega}|_{z=0} &= \partial_1(v_{0,2} - v_2^P) - \partial_2(v_{0,1} - v_1^P). \end{aligned} \quad (3-34)$$

System (3-33)–(3-34) is the formulation in terms of vertical velocity and vorticity of a Stokes–Coriolis system with zero source term and inhomogeneous Dirichlet data. Formal solutions are given by

$$\begin{pmatrix} \hat{v}_3(\xi, z) \\ \hat{w}(\xi, z) \end{pmatrix} = \sum_{i=1}^3 e^{-\lambda_i(\xi)z} C_i(\xi) V_i^-(\xi), \quad (3-35)$$

where coefficients C_i obey the system

$$\begin{pmatrix} 1 & 1 & 1 \\ \lambda_1 & \lambda_2 & \lambda_3 \\ \Omega_1 & \Omega_2 & \Omega_3 \end{pmatrix} \begin{pmatrix} C_1 \\ C_2 \\ C_3 \end{pmatrix} = \begin{pmatrix} \hat{v}_3|_{z=0} \\ -\partial_z \hat{v}_3|_{z=0} \\ -\hat{w}|_{z=0} \end{pmatrix}. \quad (3-36)$$

The determinant D_3 of this system is

$$D_3 := (\lambda_2 - \lambda_1)(\Omega_3 - \Omega_1) - (\lambda_3 - \lambda_1)(\Omega_2 - \Omega_1),$$

so that $D_3 \rightarrow \bar{D}_3 \in \mathbb{C}^*$ as $\xi \rightarrow 0^*$.

After tedious computation, we find

$$\begin{aligned} C_1 &= \frac{1}{D_3} \left((\lambda_2 \Omega_3 - \lambda_3 \Omega_2) \hat{v}_3|_{z=0} + (\Omega_3 - \Omega_2) \partial_z \hat{v}_3|_{z=0} + (\lambda_2 - \lambda_3) \hat{w}|_{z=0} \right), \\ C_2 &= \frac{1}{D_3} \left((\lambda_3 \Omega_1 - \lambda_1 \Omega_3) \hat{v}_3|_{z=0} + (\Omega_1 - \Omega_3) \partial_z \hat{v}_3|_{z=0} + (\lambda_3 - \lambda_1) \hat{w}|_{z=0} \right), \\ C_3 &= \frac{1}{D_3} \left((\lambda_1 \Omega_2 - \lambda_2 \Omega_1) \hat{v}_3|_{z=0} + (\Omega_2 - \Omega_1) \partial_z \hat{v}_3|_{z=0} + (\lambda_1 - \lambda_2) \hat{w}|_{z=0} \right). \end{aligned} \quad (3-37)$$

Nevertheless, the expressions in (3-35) are not necessarily well-defined, due to possible singularities at $\xi = 0$. In particular, if we want to apply Lemma 7, we need the coefficient in front of $e^{-\lambda_1(\xi)z}$ to contain somehow some positive power of ξ . Using the asymptotics of Lemma 3, we compute

$$|C_1(\xi)| \leq |\hat{v}_3|_{z=0}| + |\partial_z \hat{v}_3|_{z=0}| + |\hat{w}|_{z=0}|, \quad (3-38)$$

$$|C_2(\xi)| \leq |\xi| |\hat{v}_3|_{z=0}| + |\partial_z \hat{v}_3|_{z=0}| + |\hat{w}|_{z=0}|, \quad (3-39)$$

$$|C_3(\xi)| \leq |\xi| |\hat{v}_3|_{z=0}| + |\partial_z \hat{v}_3|_{z=0}| + |\hat{w}|_{z=0}| \quad (3-40)$$

for small $|\xi|$. The asymptotics is given by:

Lemma 11. *The boundary data $\partial_z \hat{v}_3|_{z=0}$, $\hat{w}|_{z=0}$ in (3-34), as well as $\hat{v}_{0,3}|_{z=0}$ (which appears in $\hat{v}_3|_{z=0}$) “contain a power of ξ at low frequencies”. More precisely, for ξ small enough, they can all be decomposed into terms of the form $\xi \cdot \hat{f}$ for some $f \in L^2_{\text{uloc}}(\mathbb{R}^2)$. As a consequence, for any function $Q \in \mathcal{C}^\infty(\mathbb{R}^2)$,*

$$\begin{aligned} & \left\| \chi(D) Q(D) \exp(-\lambda_1(D)z) \begin{pmatrix} \partial_z \underline{v}_3|_{z=0} \\ \underline{w}|_{z=0} \\ v_{0,3}|_{z=0} \end{pmatrix} \right\|_{L^\infty(\mathbb{R}^2)} \\ & \leq C(1+z)^{-1/3} \left(\|(v_{0,1}, v_{0,2})\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(1+z)^{2/3} F\|_{H^m_{\text{uloc}}(\mathbb{R}^3_+)} \right), \end{aligned}$$

and for $j = 2, 3$,

$$\begin{aligned} & \left\| \chi(D) Q(D) \exp(-\lambda_j(D)z) \begin{pmatrix} \partial_z \underline{v}_3|_{z=0} \\ \underline{\omega}|_{z=0} \\ v_{0,3}|_{z=0} \end{pmatrix} \right\|_{L^\infty(\mathbb{R}^2)} \\ & \leq C e^{-\delta z} \left(\|(v_{0,1}, v_{0,2})\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(1+z)^{2/3} F\|_{H^m_{\text{uloc}}(\mathbb{R}^3_+)} \right). \end{aligned}$$

(2) Concerning the boundary data $v_3^p|_{z=0}$ (which is the other term in $\underline{v}_3|_{z=0}$), we have, for any function $Q \in C^\infty(\mathbb{R}^2)$,

$$\begin{aligned} & \left\| (\chi(D) Q(D) \exp(-\lambda_1(D)z)) v_3^p|_{z=0} \right\|_{L^\infty(\mathbb{R}^2)} \leq C(1+z)^{-1/3} \|F\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}, \\ & \left\| (\chi(D) Q(D) \exp(-\lambda_j(D)z)) v_3^p|_{z=0} \right\|_{L^\infty(\mathbb{R}^2)} \leq C e^{-\delta z} \|F\|_{L^1_{\text{uloc}}(\mathbb{R}^2)}. \end{aligned}$$

Proof. The first part of the statement is obvious for the last two boundary data, namely

$$\partial_z \underline{v}_3|_{z=0} = -\partial_1(v_{0,1} - v_1^p) - \partial_2(v_{0,2} - v_2^p), \quad \text{and} \quad \underline{\omega}|_{z=0} = \partial_1(v_{0,2} - v_2^p) - \partial_2(v_{0,1} - v_1^p).$$

It remains to consider $v_{0,3}$. This is where the assumption on $v_{0,3}$ in the theorem plays a role. Indeed, we have $v_{0,3} = \partial_1 v_1^* + \partial_2 v_2^*$, so that it satisfies the properties of the lemma. The estimate is then a straightforward consequence of [Lemma 7](#).

The former argument does not work with the boundary data $\underline{v}_3^p|_{z=0}$: indeed, if we factor out crudely a power of ξ from the integral defining it, then the convergence of the remaining integral is no longer clear. Therefore we go back to the definition of u_3^p ; we have, using the notations of [\(3-20\)](#),

$$\chi(D) v_3^p|_{z=0} = \int_{\mathbb{R}_+} \sum_{k=0}^2 I_3^k(\cdot, 0, z') dz'.$$

It can be easily checked that the terms with I_3^k for $k = 1, 2$ do not raise any difficulty (in fact, the trace stemming from these two terms contains a power of ξ at low frequencies.) Thus we focus on

$$\int_{\mathbb{R}_+} I_3^0(\cdot, 0, z') dz' = \int_{\mathbb{R}_+} \left(\chi(D) \sum_{i=1}^3 A_i(D) e^{-\lambda_i(D)z'} s_3^0(\cdot, z') + \chi(D) \sum_{i=1}^3 B_i(D) e^{-\lambda_i(D)z'} s_\omega^0(\cdot, z') \right) dz'.$$

Applying $\exp(-\lambda_j(D)z)$, we have to estimate the $L^\infty(\mathbb{R}^2)$ norms of

$$\begin{aligned} & \int_{\mathbb{R}_+} \chi(D) Q(D) \sum_{i=1}^3 A_i(D) e^{-\lambda_i(D)z' - \lambda_j(D)z} s_3^0(\cdot, z') dz', \\ & \int_{\mathbb{R}_+} \chi(D) Q(D) \sum_{i=1}^3 B_i(D) e^{-\lambda_i(D)z' - \lambda_j(D)z} s_\omega^0(\cdot, z') dz'. \end{aligned}$$

We recall that $\widehat{s}_3^0(\xi, z')$ and $\widehat{s}_\omega^0(\xi, z')$ are products of components of $\widehat{F}(\xi, z')$ by homogeneous polynomials of degrees 2 and 3 respectively in ξ , and that the behaviour of A_i, B_i is given in [Lemma 5](#). When $i = j = 1$,

using [Lemma 7](#) and [Lemma 16](#) in [Appendix B](#), the corresponding integral is bounded by

$$\int_{\mathbb{R}_+} \frac{1}{(1+z+z')^{2/3}} \frac{1}{(1+z')^{2/3}} dz' \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}})} \leq C(1+z)^{-1/3} \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}})}.$$

When $i = 2, 3$, the integral is bounded by

$$\int_{\mathbb{R}_+} \frac{\exp(-\delta z')}{(1+z)^{2/3}} \frac{1}{(1+z')^{2/3}} dz' \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}})} \leq C(1+z)^{-2/3} \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}})}.$$

When $j = 2, 3$, the integral is bounded by

$$\int_{\mathbb{R}_+} \frac{\exp(-\delta z)}{(1+z')^{2/3}} \frac{1}{(1+z')^{2/3}} dz' \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}})} \leq C \exp(-\delta z) \|(1+z)^{2/3} F\|_{L^\infty(L^1_{\text{uloc}})}.$$

Gathering all the terms, we obtain the estimate announced in the lemma. \square

Going back to [\(3-35\)](#), we infer that

$$(1+z)^{1/3} \|\chi(D)\underline{v}_3(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} + (1+z)^{2/3} \|\chi(D)\underline{\omega}(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} \\ \leq C(\|(v_{0,1}, v_{0,2})\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(1+z)^{2/3} F\|_{H^m_{\text{uloc}}(\mathbb{R}_+^3)}). \quad (3-41)$$

Then, for further control of the horizontal components (v_1, v_2) , one would like an analogue of [\(3-25\)](#), namely a bound like

$$(1+z)^{1/3} \left\| \frac{D}{|D|^2} \chi(D) \partial_z \underline{v}_3(\cdot, z) \right\|_{L^\infty(\mathbb{R}^2)} + (1+z)^{1/3} \left\| \frac{D}{|D|^2} \chi(D) \underline{\omega}(\cdot, z) \right\|_{L^\infty(\mathbb{R}^2)} \\ \leq C(\|(v_{0,1}, v_{0,2})\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(1+z)^{2/3} F\|_{H^m_{\text{uloc}}(\mathbb{R}_+^3)}).$$

However, such an estimate is not clear. Indeed, in view of [\(3-35\)](#), we have

$$\chi(D) \begin{pmatrix} \partial_z \underline{v}_3(\cdot, z) \\ \underline{\omega}(\cdot, z) \end{pmatrix} = \chi(D) \sum_{i=1}^3 e^{-\lambda_i(D)z} \begin{pmatrix} -\lambda_i(D)C_i \\ -\Omega_i(D)C_i \end{pmatrix}.$$

The term with index $i = 1$ does not raise any difficulty, because $\lambda_1(D)$ and $\Omega_1(D)$ bring extra powers of ξ , which are enough to apply [Lemma 7](#). But the difficulty comes from indices 2 and 3. For instance, they involve terms of the type

$$\chi(D) P_0(D) e^{-\lambda_{2,3}(D)z} \hat{v}_0, \quad \text{with } P_0 \text{ homogeneous of degree } 0,$$

and therefore are not covered by [Lemma 7](#): with the notations of the lemma, one has $\alpha = 0$, which is not enough. Typically, these homogeneous functions of degree zero involve Riesz transforms, meaning $P_0(\xi) = \xi_k \xi_l / |\xi|^2$, $k, l = 1, 2$.

Hence, one must use extra cancellations. We recall that in view of [\(3-6\)](#), we want to exhibit cancellations in $|D|^{-2}(D_1 \partial_z \underline{v}_3 + D_2 \underline{\omega})$ and in $|D|^{-2}(D_2 \partial_z \underline{v}_3 - D_1 \underline{\omega})$. Let us comment briefly on the first term. We compute $(-\xi_1 \lambda_i - \xi_2 \Omega_i) C_i$ for $i = 2, 3$ in terms of the boundary data. Setting $\underline{v}_0 = v_0 - v^p|_{z=0}$, we find that

$$C_2(\xi) = \frac{1}{D_3} (\lambda_3 \Omega_1 - \lambda_1 \Omega_3) \hat{v}_{0,3} + \frac{1}{D_3} [((\Omega_3 - \Omega_1) i \xi_1 - i \xi_2 (\lambda_3 - \lambda_1)) \hat{v}_{0,1} + ((\Omega_3 - \Omega_1) i \xi_2 + i \xi_1 (\lambda_3 - \lambda_1)) \hat{v}_{0,2}].$$

We then use the asymptotic formulas of [Lemma 3](#). In particular,

$$\begin{aligned} (-\xi_1\lambda_2 - \xi_2\Omega_2)((\Omega_3 - \Omega_1)i\xi_1 - i\xi_2(\lambda_3 - \lambda_1)) &= |\xi|^2 + O(|\xi|^3), \\ (-\xi_1\lambda_2 - \xi_2\Omega_2)((\Omega_3 - \Omega_1)i\xi_2 + i\xi_1(\lambda_3 - \lambda_1)) &= -i|\xi|^2 + O(|\xi|^3). \end{aligned}$$

A similar formula holds for C_3 . It follows that there exist $Q_2, Q_3 \in C^\infty(\mathbb{R}^2)^2$ such that

$$\begin{aligned} &\mathcal{F}(\chi(D)|D|^{-2}(D_1\partial_z v_3 + D_2\omega)) \\ &= \chi(\xi) \frac{-\xi_1\lambda_1 - \xi_2\Omega_1}{D_3|\xi|^2} e^{-\lambda_1(\xi)z} C_1(\xi) \\ &\quad + \frac{1}{D_3} [(\lambda_3\Omega_1 - \lambda_1\Omega_3)(-\xi_1\lambda_2 - \xi_2\Omega_2)e^{-\lambda_2z} + (\lambda_1\Omega_2 - \lambda_2\Omega_1)(-\xi_1\lambda_3 - \xi_2\Omega_3)e^{-\lambda_3z}] \hat{v}_{0,3} \\ &\quad + \sum_{i=2,3} \chi(\xi) e^{-\lambda_i z} Q_i(\xi) \cdot \hat{v}_{0,h}(\xi, z). \end{aligned}$$

The first two terms are treated in the same way as [Lemma 11](#), factoring out a power of ξ when necessary, and going back to the definition of v^p . We leave the details to the reader. The inverse Fourier transform of the last term is $\mathcal{F}^{-1}(\chi Q_i e^{-\lambda_i z}) * v_{0,h}$, which is bounded in $L^\infty(\mathbb{R}^2)$ by $e^{-\delta z} \|v_{0,h}\|_{L^2_{\text{uloc}}}$. Similar statements hold for $\chi(D)|D|^{-2}(-\partial_z D_2 v_3 + D_1 \omega)$. It follows that

$$\begin{aligned} (1+z)^{1/3} \|\chi(D)\underline{v}(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} \\ \leq C \left(\|v_{0,1}, v_{0,2}\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L^2_{\text{uloc}}(\mathbb{R}^2)} + \|(1+z)^{2/3} F\|_{H^m_{\text{uloc}}(\mathbb{R}_+^3)} \right). \end{aligned} \quad (3-42)$$

We now address the estimates of $\hat{v}(\xi, z)$ for large frequencies. The arguments are very close to the ones developed after [Lemma 9](#). Using (3-35) and (3-37), for $|\xi| \gg 1$, we find that $\hat{v}_3(\xi, z)$ and $\hat{\omega}(\xi, z)$ can be written as linear combinations of terms of the type

$$R_{ij}(\lambda_1, \lambda_2, \lambda_3, \xi) \exp(-\lambda_i(\xi)z) \hat{g}_j(\xi), \quad 1 \leq i, j \leq 3,$$

where $g_1 = v_3|_{z=0}$, $g_2 = \partial_z v_3|_{z=0}$ and $g_3 = \omega|_{z=0}$ and R_{ij} is a rational expression. Thus, using [Lemmas 3](#) and [5](#), there exists $n \in \mathbb{N}$ such that $|\xi|^{-2n} R_{ij}(\lambda_1, \lambda_2, \lambda_3, \xi)$ is bounded as $|\xi| \rightarrow \infty$ for all i, j . [Lemma 9](#) then gives that for some N sufficiently large,

$$\begin{aligned} \|(1-\chi)(D)v_3(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C e^{-\delta z} \sum_{j=1}^3 \|g_j\|_{H^N_{\text{uloc}}}, \\ \|(1-\chi)(D)\omega(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} &\leq C e^{-\delta z} \sum_{j=1}^3 \|g_j\|_{H^N_{\text{uloc}}}, \end{aligned}$$

and similar estimates hold for $(D/|D|^2)\partial_z v_3$ and $(D/|D|^2)\omega$. Using (3-34) and (3-28)–(3-29), we infer that for some $m \geq 1$ large enough,

$$\|(1-\chi(D))\underline{v}(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} \leq C e^{-\delta z} \left(\|v_0\|_{H^{m+1/2}_{\text{uloc}}(\mathbb{R}^2)} + \|(1+z)^{2/3} F\|_{H^m_{\text{uloc}}(\mathbb{R}_+^3)} \right). \quad (3-43)$$

Gathering (3-42) and (3-43), we deduce that u satisfies the estimate

$$\|(1+z)^{1/3}\underline{v}\|_{L^\infty} \leq C\left(\|v_0\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|(1+z)^{2/3}F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)}\right)$$

for m large enough. Thus, in view of the estimate (3-3) satisfied by v^p , we know $v = \underline{v} + v^p$ is a solution of (2-3) satisfying

$$\|(1+z)^{1/3}v\|_{L^\infty} \leq C\left(\|v_0\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|(1+z)^{2/3}F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)}\right)$$

for m large enough. It remains to go to the higher regularity bound (3-1). First, up to taking a slightly larger m , we clearly have

$$\|(1+z)^{1/3}\nabla v\|_{L^\infty} \leq C\left(\|v_0\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|(1+z)^{2/3}F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)}\right).$$

This follows from direct differentiation of formula (3-7) satisfied by v^p and formula (3-35) satisfied by $\underline{v} = v - v^p$. Clearly, the differentiation is harmless, in particular at low frequencies where it may even add positive powers of ξ . It follows that our solution belongs to $H_{\text{uloc}}^1(\mathbb{R}_+^3)$, and thus enters the framework of local elliptic regularity theory for the Stokes equation. In particular, for any $k \in \mathbb{Z}^3$ with $k_z \leq 2$,

$$\begin{aligned} \|v\|_{H^{m+1}(B(k,1)\cap\Omega_{\text{bl}})} &\leq C\left(\|v_0\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)} + \|v\|_{H^1(B(k,2)\cap\Omega_{\text{bl}})}\right) \\ &\leq C\left(\|v_0\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)} + \|v\|_{H_{\text{uloc}}^1(\mathbb{R}_+^3)}\right) \end{aligned}$$

and for any $k \in \mathbb{Z}^3$ with $k_z > 2$,

$$\begin{aligned} \|v\|_{H^{m+1}(B(k,1)\cap\Omega_{\text{bl}})} &\leq C\left(\|F\|_{H^m(B(k,2)\cap\Omega_{\text{bl}})} + \|v\|_{H^1(B(k,2)\cap\Omega_{\text{bl}})}\right) \\ &\leq C|k_z|^{-1/3}\left(\|(1+z)^{2/3}F\|_{H_{\text{uloc}}^m(\mathbb{R}_+^3)} + \|(1+y)^{1/3}v\|_{H_{\text{uloc}}^1(\mathbb{R}_+^3)}\right). \end{aligned}$$

The bound (3-1) follows.

4. Proof of Theorem 1

4.1. Navier–Stokes–Coriolis system in a half-space. This section is devoted to the well-posedness of system (2-4) under a smallness assumption. Once again, we can assume $M = 0$ with no loss of generality. Following the analysis of the linear case performed in the previous section, we introduce the functional spaces

$$\mathcal{H}^m := \left\{v \in H_{\text{loc}}^m(\mathbb{R}_+^3) : \|(1+y_3)^{1/3}v\|_{H_{\text{uloc}}^m} < +\infty\right\}, \quad m \geq 0,$$

and we set $\|v\|_{\mathcal{H}^m} = C_m \|(1+y_3)^{1/3}v\|_{H_{\text{uloc}}^m}$, where the constant C_m is chosen so that if $u, v \in (\mathcal{H}^m)^3$ for some $m > \frac{3}{2}$, then

$$\|u \otimes v\|_{\mathcal{H}^m} \leq \|u\|_{\mathcal{H}^m} \|v\|_{\mathcal{H}^m}.$$

Clearly \mathcal{H}^m is a Banach space for all $m \geq 0$.

The result proved in this section is the following:

Proposition 12. *Let $m \in \mathbb{N}$, $m \gg 1$. There exists $\delta_0 > 0$ such that for all $v_0 \in H_{\text{uloc}}^{m+1}(\mathbb{R}^2)$ such that $v_{0,3} = \partial_1 v_1^* + \partial_2 v_2^*$, with v_1^*, v_2^* in $L_{\text{uloc}}^2(\mathbb{R}^2)$ and*

$$\|v_0\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} \leq \delta_0, \quad (4-1)$$

the system

$$\begin{cases} v \cdot \nabla v + e \times v + \nabla p - \Delta v = 0 & \text{in } \{y_3 > 0\}, \\ \operatorname{div} v = 0 & \text{in } \{y_3 > 0\}, \\ v|_{y_3=0} = v_0 \end{cases}$$

has a unique solution in \mathcal{H}^{m+1} .

Remark 13. The integer m for which this result holds is the same as the one in [Theorem 2](#).

Proof. [Proposition 12](#) is an easy consequence of the fixed point theorem in \mathcal{H}^{m+1} . For any $v_0 \in H_{\text{uloc}}^{m+1}(\mathbb{R}^2)$ such that $v_{0,3} = \partial_1 v_1^* + \partial_2 v_2^*$, with v_1^*, v_2^* in $L_{\text{uloc}}^2(\mathbb{R}^2)$, we introduce the mapping $T_{v_0} : \mathcal{H}^{m+1} \rightarrow \mathcal{H}^{m+1}$ such that $T_{v_0}(u) = v$ is the solution of (2-3) with $F = u \otimes u$. Notice that $\|(1+z)^{2/3} F\|_{H_{\text{uloc}}^m} \leq \|u\|_{\mathcal{H}^m}^2$. As a consequence, according to [Theorem 2](#), there exists a constant C_0 such that for all $u \in \mathcal{H}^{m+1}$,

$$\|T_{v_0}(u)\|_{\mathcal{H}^{m+1}} \leq C_0 (\|v_0\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2)} + \|(v_1^*, v_2^*)\|_{L_{\text{uloc}}^2(\mathbb{R}^2)} + \|u\|_{\mathcal{H}^{m+1}}^2). \quad (4-2)$$

Let $\delta_0 < 1/(4C_0^2)$, and assume that (4-1) is satisfied. Thanks to the assumption on δ_0 , there exists $R_0 > 0$ such that

$$C_0(\delta_0 + R_0^2) \leq R_0. \quad (4-3)$$

Moreover, $R_0 \in [R_-, R_+]$, where

$$R_{\pm} = \frac{1}{2C_0} (1 \pm \sqrt{1 - 4\delta_0 C_0^2}).$$

Therefore $0 < R_- < (2C_0)^{-1}$, and we can always choose R_0 so that $2R_0 C_0 < 1$. Then according to (4-1), (4-2) and (4-3),

$$\|u\|_{\mathcal{H}^{m+1}} \leq R_0 \implies \|T_{v_0}(u)\|_{\mathcal{H}^{m+1}} \leq R_0.$$

Moreover, if $\|u^1\|_{\mathcal{H}^{m+1}}, \|u^2\|_{\mathcal{H}^{m+1}} \leq R_0$, then setting $w = T_{v_0}(u^1) - T_{v_0}(u^2)$, we have w is a solution of (2-3) with $w|_{z=0} = 0$ and with a source term $F^1 - F^2 = u^1 \otimes u^1 - u^2 \otimes u^2$. Thus, using once again [Theorem 2](#) and the normalization of $\|\cdot\|_{\mathcal{H}^m}$,

$$\|T_{v_0}(u^1) - T_{v_0}(u^2)\|_{\mathcal{H}^{m+1}} \leq C_0 \|F^1 - F^2\|_{\mathcal{H}^m} \leq 2C_0 R_0 \|u^1 - u^2\|_{\mathcal{H}^{m+1}}.$$

Notice that in the inequality above, we have assumed that $\|\cdot\|_{\mathcal{H}^m} \leq \|\cdot\|_{\mathcal{H}^{m+1}}$, which is always possible if the normalization constant C_m is chosen sufficiently small (depending on C_{m+1} , m being large but fixed).

Thus, since $2C_0 R_0 < 1$, we know T_{v_0} is a contraction over the ball of radius R_0 in \mathcal{H}^{m+1} . Using Banach's fixed point theorem, we infer that T_{v_0} has a fixed point in \mathcal{H}^{m+1} . This concludes the proof of [Proposition 12](#). \square

4.2. Navier–Stokes–Coriolis system over a bumped half-plane. We now address the study of the full system (1-4). We follow the steps outlined in the introduction, which we recall here for the reader’s convenience: We first prove that there exists a solution (v^-, p^-) of the system (2-5) for ϕ, ψ in some function spaces to be specified, then construct the solution (v^+, p^+) of (2-4) with $v^+|_{y_3=M} = v^-|_{y_3=M}$. Eventually, we define a mapping \mathcal{F} by $\mathcal{F}(\phi, \psi) := \Sigma(v^+, p^+)e_3|_{y_3=M} - \psi$. We recall that $v = \mathbf{1}_{y_3 \geq M}v^+ + \mathbf{1}_{y_3 < M}v^-$ is a solution of (1-4) if and only if $\mathcal{F}(\phi, \psi) = 0$. The goal is therefore to show that for all ϕ small enough (in a function space to be specified) the equation $\mathcal{F}(\phi, \psi) = 0$ has a unique solution.

Step 1. We study the system (2-5). We introduce the function space

$$\mathcal{V} := \left\{ \phi = (\phi_h, \phi_3) : \phi_h \in H_{\text{uloc}}^2(\partial\Omega_{\text{bl}}), \phi_3 \in H_{\text{uloc}}^1(\partial\Omega_{\text{bl}}), \phi \cdot n|_{\partial\Omega_{\text{bl}}} = 0 \right\} \quad (4-4)$$

for the bottom Dirichlet data, and we set

$$\|\phi\|_{\mathcal{V}} := \|\phi_h\|_{H_{\text{uloc}}^2} + \|\phi_3\|_{H_{\text{uloc}}^1}.$$

As for the stress tensor at $y_3 = M$, since we will need to construct solutions in H_{uloc}^{m+1} (see Proposition 12), we look for ψ in the space $H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$. We then claim that the following result holds:

Lemma 14. *Let $m \geq 1$ be arbitrary. There exists $\delta > 0$ such that for all $\phi \in \mathcal{V}$ and all $\psi \in H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$ with $\|\phi\|_{\mathcal{V}} \leq \delta$ and $\|\psi\|_{H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)} \leq \delta$, system (2-5) has a unique solution*

$$(v^-, p^-) \in H_{\text{uloc}}^1(\Omega_{\text{bl}}^M) \times L_{\text{uloc}}^2(\Omega_{\text{bl}}^M).$$

Moreover, it satisfies the following properties:

- H_{uloc}^{m+1} regularity: for all $M' \in]\text{sup } \gamma, M[$,

$$(v^-, p^-) \in H_{\text{uloc}}^{m+1}(\mathbb{R}^2 \times (M', M)) \times H_{\text{uloc}}^m(\mathbb{R}^2 \times (M', M)),$$

with

$$\|v^-\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}^2 \times (M', M))} + \|p^-\|_{H_{\text{uloc}}^m(\mathbb{R}^2 \times (M', M))} \leq C_{M'}(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)}).$$

- Compatibility condition: there exists $v_1^*, v_2^* \in H_{\text{uloc}}^{1/2}$ such that $v_3^-|_{y_3=M} = \nabla_h \cdot v_h^*$.

Proof. We start with an H_{uloc}^1 a priori estimate. We follow the computations of [Dalibard and Prange 2014], dedicated to the linear Stokes–Coriolis system. We first lift the boundary condition on $\partial\Omega_{\text{bl}}$, introducing

$$v_h^L := \phi_h, \quad v_3^L := \phi_3 - \nabla_h \cdot \phi_h(y_3 - \gamma(y_h)).$$

Then $\tilde{v} := v^- - v^L$ and $\tilde{p} := p^-$ satisfy

$$\begin{aligned} -\Delta \tilde{v} + (v^L + \tilde{v}) \cdot \nabla \tilde{v} + \tilde{v} \cdot \nabla v^L + e_3 \wedge \tilde{v} + \nabla \tilde{p} &= f & \text{in } \Omega_{\text{bl}}^M, \\ \text{div } \tilde{v} &= 0 & \text{in } \Omega_{\text{bl}}^M, \\ \tilde{v}|_{\partial\Omega_{\text{bl}}} &= 0, \end{aligned} \quad (4-5)$$

$$\left(\partial_3 \tilde{v} - \left(\tilde{p} + \frac{|\tilde{v} + v^L|^2}{2} \right) e_3 \right) \Big|_{y_3=M} = \psi - \partial_3 v^L|_{y_3=M} := \tilde{\psi},$$

where $f = -\Delta v^L + v^L \cdot \nabla v^L + e_3 \wedge v^L$.

Notice that thanks to the regularity assumptions on ϕ and v^* , we have $\tilde{\psi} \in L^2_{\text{uloc}}(\mathbb{R}^2)$ and $f \in H^{-1}_{\text{uloc}}(\mathbb{R}^2)$. We then perform energy estimates on the system (4-5), following the strategy of Gérard-Varet and Masmoudi [2010], which is inspired by the work of Ladyžhenskaya and Solonnikov [1980]. The idea is to work with the truncated energies

$$E_k := \int_{\Omega_{\text{bl}}^M \cap \{(y_1, y_2) \in [-k, k]^2\}} \nabla \tilde{v} \cdot \nabla \tilde{v}, \quad (4-6)$$

and to derive an induction inequality on $(E_k)_{k \in \mathbb{N}}$. To that end, we consider a truncation function $\chi_k \in C_0^\infty(\mathbb{R}^2)$ such that $\chi_k \equiv 1$ in $[-k, k]^2$, $\text{Supp } \chi_k \subset [-k-1, k+1]^2$, and $\chi_k, \chi'_k, \chi''_k$ are bounded uniformly in k . Along the lines of [Dalibard and Prange 2014], we multiply (4-5) by the test function

$$\begin{aligned} \varphi &= \begin{pmatrix} \varphi_h \\ \nabla \cdot \Phi_h \end{pmatrix} := \begin{pmatrix} \chi_k \tilde{v}_h \\ -\nabla_h \cdot \left(\chi_k \int_{\gamma(y_h)}^{y_3} \tilde{v}_h(y_h, z) dz \right) \end{pmatrix} \in H^1(\Omega^b) \\ &= \chi_k \tilde{v} - \begin{pmatrix} 0 \\ \nabla_h \chi_k(y_h) \cdot \int_{\gamma(y_h)}^{y_3} \tilde{v}_h(y_h, z) dz \end{pmatrix}. \end{aligned}$$

Since this test function is divergence-free, there is no commutator term stemming from the pressure. In [loc. cit.], an inequality of the following type is derived:

$$E_k \leq C((E_{k+1} - E_k) + (\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H^{-1/2}_{\text{uloc}}}^2)(k+1)^2).$$

This discrete differential inequality is a key a priori estimate, which allows for the construction of a solution. Indeed, introducing an approximate solution \tilde{v}^n for $|y_1, y_2| \leq n$, say with Dirichlet boundary conditions at the lateral boundary, a standard estimate yields that $E_n \leq Cn$, where this time $E_k = \int |\chi_k \nabla \tilde{v}^n|^2$. Combining this information with above induction relation allows one to obtain a uniform bound on the E_k of the type $E_k \leq Ck^2$, from which we deduce a H^1_{uloc} bound on \tilde{v}^n uniformly in n . From there, one obtains an exact solution by compactness. We refer to [loc. cit.] for more details.

Here, there are two noticeable differences with [loc. cit.]:

- The boundary condition at $y_3 = M$ in (4-5) does not involve a Dirichlet-to-Neumann operator, which makes things easier.
- On the other hand, one has to handle the quadratic terms $(v^L + \tilde{v}) \cdot \nabla \tilde{v} + \tilde{v} \cdot \nabla v^L$, which explains the introduction of the $|v|^2$ in the stress tensor at $y_3 = M$.

Therefore we focus on the treatment of these nonlinear terms. The easiest one is

$$\left| \int_{\Omega_{\text{bl}}^M} (\tilde{v} \cdot \nabla v^L) \cdot \varphi \right| \leq C \|\phi\|_{\mathcal{V}} E_{k+1},$$

where the constant C depends only on M and on $\|\gamma\|_{W^{1,\infty}}$. On the other hand,

$$\begin{aligned} \int_{\Omega_{\text{bl}}^M} ((v^L + \tilde{v}) \cdot \nabla \tilde{v}) \cdot (\chi_k \tilde{v}) &= \int_{\Omega_{\text{bl}}^M} \chi_k (v^L + \tilde{v}) \cdot \nabla \frac{|\tilde{v}|^2}{2} \\ &= - \int_{\Omega_{\text{bl}}^M} \frac{|\tilde{v}|^2}{2} (v^L + \tilde{v}) \cdot \nabla \chi_k + \int_{\mathbb{R}^2} \chi_k \left((v_3^L + \tilde{v}_3) \frac{|\tilde{v}|^2}{2} \right) \Big|_{y_3=M}. \end{aligned}$$

The first term on the right-hand side is bounded by $C(E_{k+1} - E_k)^{3/2} + C\|\phi\|_{\mathcal{V}}(E_{k+1} - E_k)$. We group the second one with the boundary terms stemming from the pressure and the Laplacian. The sum of these three boundary terms is

$$\int_{\mathbb{R}^2} \chi_k \left(-\partial_3 \tilde{v} \cdot \tilde{v} + (v_3^L + \tilde{v}_3) \frac{|\tilde{v}|^2}{2} + p^- \tilde{v}_3 \right) \Big|_{y_3=M}.$$

Using the boundary condition in (4-5), the integral above is equal to

$$- \int_{\mathbb{R}^2} \chi_k \tilde{v}|_{y_3=M} \cdot \left(\tilde{\psi} + (\tilde{v} \cdot v^L|_{y_3=M} + \frac{1}{2}|v^L|_{y_3=M}|^2) e_3 \right),$$

which is bounded for any $\delta > 0$ by

$$C\|\phi\|_{\mathcal{V}} E_{k+1} + \delta E_{k+1} + C_\delta (\|\phi\|_{\mathcal{V}}^2 + \|\phi\|_{\mathcal{V}}^4 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2) (k+1)^2.$$

There remains

$$\int_{\Omega_{\text{bl}}^M} ((v^L + \tilde{v}) \cdot \nabla \tilde{v}) \cdot \left(\nabla_h \chi_k(y_h) \cdot \int_{\mathcal{Y}(y_h)}^{y_3} \tilde{v}_h(y_h, z) dz \right),$$

which is bounded by $C(E_{k+1} - E_k)^{3/2} + C\|\phi\|_{\mathcal{V}}(E_{k+1} - E_k)$. Gathering all the terms, we infer that for $\|\phi\|_{\mathcal{V}} \leq 1$,

$$E_k \leq C((E_{k+1} - E_k)^{3/2} + (E_{k+1} - E_k) + \|\phi\|_{\mathcal{V}} E_k + (\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2) (k+1)^2),$$

where the constant C depends only on M and on $\|\gamma\|_{W^{1,\infty}}$. As a consequence, for $\|\phi\|_{\mathcal{V}}$ small enough, we infer that for all $k \geq 1$,

$$E_k \leq C((E_{k+1} - E_k)^{3/2} + (E_{k+1} - E_k) + (\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2) (k+1)^2).$$

Thanks to a backwards induction argument (again, we refer to [Gérard-Varet and Masmoudi 2010] for all details), we infer that

$$E_k \leq C(\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2) k^2 \quad \forall k \in \mathbb{N}$$

for a possibly different constant C . It follows that

$$\|\tilde{v}\|_{H_{\text{uloc}}^1(\Omega_{\text{bl}}^M)} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}})$$

and therefore v^- satisfies the same estimate. From there, we can derive an L_{uloc}^2 estimate for the pressure. Indeed, using the equation and the boundary condition at $y_3 = M$, it follows that for all $y \in \Omega_{\text{bl}}^M$,

$$p^-(y_h, y_3) = \partial_3 v_3^-|_{y_3=M} - \frac{|v^-|_{y_3=M}|^2}{2} - \psi_3(y_h) - \int_{y_3}^M (\Delta v_3^- - v^- \cdot \nabla v_3^-)(y_h, z) dz.$$

Note that by the divergence-free condition, the first-term in the right-hand side can be written as $-\text{div}_h v_h^-|_{y_3=M}$. For $k \in \mathbb{Z}^2$, let $\varphi_k \in H_0^1(\Omega_{\text{bl}}^M)$ such that $\text{Supp } \varphi_k \subset (k + [0, 1]^2) \times \mathbb{R}$. We multiply

the above identity by $\varphi_k(x_h, z)$ and integrate over Ω_{bl}^M . After some integrations by parts, we obtain

$$\begin{aligned} \int_{\Omega_{\text{bl}}^M} p^- \varphi_k &= \int_{\Omega_{\text{bl}}^M} v_h^-|_{y_3=M} \cdot \nabla_h \varphi_k - \int_{\Omega_{\text{bl}}^M} \frac{|v^-|_{y_3=M}|^2}{2} \varphi_k - \int_{\Omega_{\text{bl}}^M} \psi_3 \varphi_k \\ &\quad - \int_{\Omega_{\text{bl}}^M} \left(\int_{y_3}^M (\Delta_h v_3^- + \partial_3^2 v_3^- - v^- \cdot \nabla v_3^-)(y_h, z) dz \right) \varphi_k(y) dy. \end{aligned} \quad (4-7)$$

Using classical trace estimates and Sobolev embeddings, it follows that for all $q \in]1, \infty[$,

$$\|v^-|_{y_3=M}\|_{L^q_{\text{uloc}}(\mathbb{R}^2)} \leq C \|v^-|_{y_3=M}\|_{H^{1/2}_{\text{uloc}}(\mathbb{R}^2)} \leq C \|v^-\|_{H^1_{\text{uloc}}(\Omega_{\text{bl}}^M)}. \quad (4-8)$$

Therefore the top line of the right-hand side of (4-7) is bounded by $C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}})\|\varphi_k\|_{H^1}$ for ϕ, ψ small enough. We now focus on the second line of (4-7). The easiest term is the advection term: we have, since φ_k has a bounded support (uniformly in k),

$$\left| \int_{\Omega_{\text{bl}}^M} \int_{y_3}^M v^- \cdot \nabla v_3^-(x_h, z) dz \varphi_k(y) dy \right| \leq C \|v^-\|_{L^4_{\text{uloc}}} \|\nabla v^-\|_{L^2_{\text{uloc}}} \|\varphi_k\|_{L^4} \leq C \|\varphi_k\|_{H^1} \|v^-\|_{H^1_{\text{uloc}}}^2.$$

We then treat the two terms stemming from the Laplacian separately. For the horizontal derivatives, we merely integrate by parts, recalling that $\varphi_k \in H^1_0(\Omega_{\text{bl}}^M)$, so that

$$\int_{\Omega_{\text{bl}}^M} \int_{y_3}^M \Delta_h v_3^-(y_h, z) dz \varphi_k(y) dy = - \int_{\Omega_{\text{bl}}^M} \int_{y_3}^M \nabla_h v_3^-(y_h, z) \cdot \nabla_h \varphi_k(y) dz dy,$$

and the corresponding term is bounded by $C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}})\|\varphi_k\|_{H^1}$. As for the vertical derivatives, we have

$$\begin{aligned} \int_{\Omega_{\text{bl}}^M} \left(\int_{y_3}^M \partial_3^2 v_3^-(y_h, z) dz \right) \varphi_k(y) dy &= \int_{\Omega_{\text{bl}}^M} (\partial_3 v_3^-(y_h, M) - \partial_3 v_3^-(y)) \varphi_k(y) dy \\ &= - \int_{\Omega_{\text{bl}}^M} (\nabla_h \cdot v_h^-(y_h, M) + \partial_3 v_3^-(y)) \varphi_k(y) dy \\ &= \int_{\Omega_{\text{bl}}^M} v_h^-(y_h, M) \cdot \nabla_h \varphi_k(y) dy - \int_{\Omega_{\text{bl}}^M} \partial_3 v_3^-(y) \varphi_k(y) dy. \end{aligned} \quad (4-9)$$

Both terms of the right-hand side are bounded by $C\|v^-\|_{H^1_{\text{uloc}}}\|\varphi_k\|_{H^1}$.

Taking the estimate (4-7), we infer that there exists a constant C (independent of φ_k and of k) such that for all $\varphi_k \in H^1_0(\Omega_{\text{bl}}^M)$ supported in $(k + [0, 1]^2) \times \mathbb{R}$,

$$\left| \int_{\Omega_{\text{bl}}^M} p^- \varphi_k \right| \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}})\|\varphi_k\|_{H^1_0(\Omega_{\text{bl}}^M)}.$$

We deduce that

$$\|p^-\|_{H^{-1}_{\text{uloc}}(\Omega_{\text{bl}}^M)} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}}).$$

Using the equation on (v^-, p^-) , we also have

$$\|\nabla p^-\|_{H^{-1}_{\text{uloc}}(\Omega_{\text{bl}}^M)} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}}).$$

It then follows from Nečas inequality (see [Boyer and Fabrie 2013, Theorem IV.1.1]) that $p^- \in L^2_{\text{uloc}}(\Omega_{\text{bl}}^M)$, with

$$\|p^-\|_{L^2_{\text{uloc}}(\Omega_{\text{bl}}^M)} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}}).$$

We still have to establish the two properties itemized in Lemma 14. We focus first on the higher-order estimates. Note that using interior regularity results for the Stokes system (see [Galdi 2011]), one has $v^- \in H^N_{\text{uloc}}(\Omega')$ for all open sets $\Omega' \subset \mathbb{R}^2$ such that $\bar{\Omega}' \subset \Omega_{\text{bl}}^M$ and for all $N > 0$. In particular, for all $M_1 < M_2$ in the interval $]\text{sup } \gamma, M[$, we have $v^- \in H^{m+1}_{\text{uloc}}(\mathbb{R}^2 \times (M_1, M_2))$ and $p^- \in H^m_{\text{uloc}}(\mathbb{R}^2 \times (M_1, M_2))$.

We now tackle the regularity for $y_3 > M'$, where $M' \in]\text{sup } \gamma, M[$. Our arguments are somehow standard (and mainly taken from [Boyer and Fabrie 2013]), but since there are a few difficulties related to the nonlinear stress boundary condition at $y_3 = M$, we give details. The idea is to use an induction argument to show that $v^- \in H^l_{\text{uloc}}(\mathbb{R}^2 \times [M', M])$ for all $\text{sup } \gamma < M' < M$ and for $1 \leq l \leq m+1$. Unfortunately, the induction only works for $l \geq 2$: indeed, the implication $h \in H^s(\mathbb{R}^2) \Rightarrow h^2 \in H^s(\mathbb{R}^2)$, which is required to handle the nonlinear boundary condition at $y_3 = M$, is true for $s > 1$ only. Therefore we treat separately the case $l = 2$. In the sequel, we write $\|\phi\| + \|\psi\|$ as a shorthand for $\|\phi\|_{\mathcal{V}} + \|\psi\|_{H^{m-1/2}_{\text{uloc}}}$.

To prove H^2_{uloc} regularity, the first step is to prove a priori estimates for $\partial_1 v^-$, $\partial_2 v^-$ in H^1_{uloc} . To that end, we first localize the equation near a fixed $k \in \mathbb{Z}^2$, then differentiate it with respect to y_j , $j = 1, 2$. Let $\theta \in C^\infty_0(\mathbb{R}^2)$ be equal to 1 in a neighbourhood of $k \in \mathbb{Z}^2$, and such that the size of $\text{Supp } \theta$ is bounded uniformly in k (we omit the k -dependence of θ and of all subsequent functions in order to alleviate the notation). It can be easily checked that the equation satisfied by $w_j := \partial_j(\theta v^-)$ is

$$\begin{aligned} -\Delta w_j + e_3 \wedge w_j + v^- \cdot \nabla w_j + \nabla \partial_j(\theta p^-) &= F_j \quad \text{in } \Omega_\theta, \\ \text{div } w_j &= g_j \quad \text{in } \Omega_\theta, \\ w_j|_{y_3=M'} &\in H^{1/2}(\mathbb{R}^2), \\ (\partial_3 w_j - (\partial_j(\theta p^-) + v^- \cdot w_j - \frac{1}{2}|v^-|^2 \partial_j \theta) e_3)|_{y_3=M} &= \partial_j(\theta \psi), \\ w_j &= 0 \quad \text{on } \partial \text{Supp } \theta \times (M', M), \end{aligned}$$

where $\Omega_\theta := \text{Supp } \theta \times (M', M)$ and

$$\begin{aligned} F_j &= \underbrace{\partial_j(-2\nabla\theta \cdot \nabla v^- - v^- \Delta\theta + (v^- \cdot \nabla\theta)v^- + p^- \nabla\theta)}_{\|\cdot\|_{H^{-1}} \leq C(\|\phi\| + \|\psi\|)} - \partial_j v^- \cdot \nabla(\theta v^-), \\ g_j &= \partial_j(v^- \cdot \nabla\theta) = O(\|\phi\| + \|\psi\|) \quad \text{in } L^2(\mathbb{R}^2 \times (M', M)). \end{aligned}$$

By standard results, see [Galdi 2011, Section II.3], there exists $\bar{w}_j \in H^1(\Omega_\theta)$ such that

$$\begin{aligned} \text{div } \bar{w}_j &= g_j, \quad \bar{w}_j = w_j \quad \text{at } \partial\Omega_\theta \setminus \{y_3 = M\}, \\ \|\bar{w}_j\|_{H^1(\Omega_\theta)} &\leq C(\|g_j\|_{L^2(\Omega_\theta)} + \|w_j\|_{H^{1/2}(\{y_3=M'\})}). \end{aligned}$$

Note that we do not need to correct the trace of w_j at $\{y_3 = M\}$, as there is no Dirichlet boundary condition there. Moreover, we are not sure at this stage that this trace is an $H^{1/2}_{\text{uloc}}$ function. We rather prescribe an artificial smooth data for \bar{w}_j at this boundary, chosen so that it satisfies the good compatibility condition.

Finally, $\tilde{w}_j = w_j - \bar{w}_j$ satisfies

$$\begin{aligned} -\Delta \tilde{w}_j + e_3 \wedge \tilde{w}_j + v^- \cdot \nabla \tilde{w}_j + \nabla \tilde{q}_j &= \tilde{F}_j \quad \text{in } \Omega_\theta, \\ \operatorname{div} \tilde{w}_j &= 0 \quad \text{in } \Omega_\theta, \\ \tilde{w}_j|_{y_3=M'} &= 0, \quad \tilde{w}_j = 0 \quad \text{on } \partial \operatorname{Supp} \theta \times (M', M), \\ (\partial_3 \tilde{w}_j - (\tilde{q}_j + v^- \cdot \tilde{w}_j)e_3)|_{y_3=M} &= \tilde{\psi}_j, \end{aligned}$$

with $\tilde{F}_j = -\partial_j v^- \cdot \nabla(\theta v^-) + O(\|\phi\| + \|\psi\|)$ in H^{-1} , and $\|\tilde{\psi}_j\|_{H^{-1/2}} \leq C(\|\phi\| + \|\psi\|)$. We obtain the estimate

$$\|\nabla \tilde{w}_j\|_{L^2(\Omega_\theta)}^2 \leq C(\|\phi\|^2 + \|\psi\|^2) + \left| \int_{\Omega_\theta} (\partial_j v^- \cdot \nabla(\theta v^-)) \cdot \tilde{w}_j \right| + 2 \int_{\operatorname{Supp} \theta} |v^-|_{y_3=M} |\tilde{w}_j|_{y_3=M}|^2.$$

We first deal with the boundary term:

$$\begin{aligned} \int_{\operatorname{Supp} \theta} |v^-|_{y_3=M} |\tilde{w}_j|_{y_3=M}|^2 &\leq \|v^-|_{y_3=M}\|_{L^2(\operatorname{Supp} \theta)} \|\tilde{w}_j|_{y_3=M}\|_{L^4(\operatorname{Supp} \theta)}^2 \\ &\leq C \|v^-|_{y_3=M}\|_{H_{\text{uloc}}^{1/2}} \|\tilde{w}_j|_{y_3=M}\|_{H^{1/2}(\operatorname{Supp} \theta)}^2 \leq C \|v^-\|_{H^1} \|\tilde{w}_j\|_{H^1}^2 \\ &\leq C(\|\phi\| + \|\psi\|) \|\nabla \tilde{w}_j\|_{L^2}^2. \end{aligned}$$

Hence for ψ and ϕ small enough we can absorb this term in the left-hand side of the energy inequality. As for the quadratic source term, we write

$$\begin{aligned} \partial_j v^- \cdot \nabla(\theta v^-) &= \partial_j v_1^- w_1 + \partial_j v_2^- w_2 + \partial_j v_3^- \theta \partial_3 v^- \\ &= \partial_j v_1^- w_1 + \partial_j v_2^- w_2 + \partial_3 v^- w_{j,3} - v_3^- \partial_j \theta \partial_3 v^-. \end{aligned}$$

For $i = 1, \dots, 3$, $j = 1, 2$, $k = 1, 2$, we have

$$\begin{aligned} \int_{\Omega_\theta} |\partial_i v^-| |w_j| |\tilde{w}_k| &\leq C \|v^-\|_{H_{\text{uloc}}^1(\Omega_{\text{bl}}^M)} \|w_j\|_{L^4(\Omega_\theta)} \|\tilde{w}_k\|_{L^4(\Omega_\theta)} \\ &\leq C(\|\phi\| + \|\psi\|) (\|\tilde{w}_1\|_{H^1(\Omega_\theta)}^2 + \|\tilde{w}_2\|_{H^1(\Omega_\theta)}^2) + C(\|\phi\| + \|\psi\|)^3 \end{aligned}$$

and

$$\left| \int_{\Omega_\theta} v_3^- \partial_j \theta \partial_3 v^- \cdot \tilde{w}_j \right| \leq C \|v_3^-\|_{H_{\text{uloc}}^1} \|\partial_3 v^-\|_{L_{\text{uloc}}^2} \|\tilde{w}_j\|_{H^1(\Omega_\theta)}.$$

Therefore, we obtain, for $\|\phi\| + \|\psi\|$ small enough,

$$\|w_1\|_{H^1(\Omega_\theta)}^2 + \|w_2\|_{H^1(\Omega_\theta)}^2 \leq C(\|\phi\|^2 + \|\psi\|^2).$$

Using the same idea as above to estimate $\partial_j(\theta p^-)$, this gives

$$\|\nabla_h v^-\|_{H_{\text{uloc}}^1(\mathbb{R}^2 \times (M', M))} + \|\nabla_h p^-\|_{L_{\text{uloc}}^2(\mathbb{R}^2 \times (M', M))} \leq (\|\phi\| + \|\psi\|)_{H_{\text{uloc}}^{m-1/2}}.$$

Since v^- is divergence-free, similar estimates hold for $\partial_3 v_3^-$. Thus $v_3^- \in H_{\text{uloc}}^2(\mathbb{R}^2 \times (M', M))$. As for the vertical regularity of v_h^- , we observe that $\partial_3 v^-$ is a solution of the Stokes system with Dirichlet boundary

conditions

$$\begin{aligned} -\Delta \partial_3 v^- + \nabla \partial_3 p^- &= F_3 && \text{in } \mathbb{R}^2 \times (M', M), \\ \operatorname{div} \partial_3 v^- &= 0 && \text{in } \mathbb{R}^2 \times (M', M), \\ \partial_3 v^-|_{y_3=M} &= G, \\ \partial_3 v^-|_{y_3=M'} &= G', \end{aligned}$$

where

$$F_3 = -e_3 \wedge \partial_3 v^- - \partial_3(v_h^- \cdot \nabla_h v^-) - \partial_3(v_3^- \partial_3 v^-) \in H_{\text{uloc}}^{-1}(\mathbb{R}^2), \quad G_h = \psi_h \in H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2),$$

and $G_3 = \partial_3 v_3^-|_{y_3=M} \in H_{\text{uloc}}^{1/2}(\mathbb{R}^2)$, $G' \in H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$. Using the results of Chapter IV in [Galdi 2011], we infer that $\partial_3 v^- \in H_{\text{uloc}}^1(\mathbb{R}^2 \times (M', M))$, $\partial_3 p^- \in L_{\text{uloc}}^2(\mathbb{R}^2 \times (M', M))$, and

$$\|\partial_3 v^-\|_{H_{\text{uloc}}^1(\mathbb{R}^2 \times (M', M))} + \|\partial_3 p^-\|_{L_{\text{uloc}}^2(\mathbb{R}^2 \times (M', M))} \leq C(\|F\|_{H_{\text{uloc}}^{-1}} + \|G\|_{H_{\text{uloc}}^{1/2}} + \|G'\|_{H_{\text{uloc}}^{1/2}}) \leq C(\|\phi\| + \|\psi\|)$$

for ϕ and ψ small enough. Gathering the inequalities, we obtain

$$\|v^-\|_{H_{\text{uloc}}^2(\mathbb{R}^2 \times (M', M))} + \|p^-\|_{H_{\text{uloc}}^1(\mathbb{R}^2 \times (M', M))} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}).$$

Of course, all inequalities above are a priori estimates, but provide H_{uloc}^2 regularity (and a posteriori estimates) through the usual method of translations.

We are now ready for the induction argument. Let $k \in \mathbb{Z}^2$ be fixed. Define a sequence $\vartheta_k^2, \dots, \vartheta_k^{m+1}$ such that $\vartheta_k^l := \theta_1^l(z - M)\theta_2^l(y_h - k)$, where $\theta_1^l \in C_0^\infty(\mathbb{R})$, $\theta_2^l \in C_0^\infty(\mathbb{R}^2)$ are equal to 1 in a neighbourhood of zero. We require furthermore that $\operatorname{Supp} \vartheta_k^{l+1} \subset (\vartheta_k^l)^{-1}(\{1\})$. We then define a $C^{m+1,1}$ domain $\Omega_k \subset \Omega_{\text{bl}}^M$ such that $\operatorname{Supp} \vartheta_k^2 \Subset \overline{\Omega}_k$, and such that $\partial\Omega_k \cap \partial\Omega_{\text{bl}} = \emptyset$ (see Figure 1). Notice also that we choose Ω_k so that $\operatorname{diam}(\Omega_k)$ is bounded uniformly in k (in fact, we can always assume that $\Omega_k = (k, 0) + \Omega_0$ for some fixed domain Ω_0).




Figure 1. The domain Ω_k .

Multiplying (2-5) by ϑ_k^l and dropping the dependence with respect to k , we find that $v^l := \vartheta_k^l v^-$, $p^l := p^- \vartheta_k^l$ is a solution of

$$\begin{cases} -\Delta v^l + \nabla p^l = f^l & \text{in } \Omega_k, \\ \operatorname{div} v^l = g^l & \text{in } \Omega_k, \\ \partial_n v^l - p^l n = \Sigma^l & \text{on } \partial\Omega_k, \end{cases} \quad (4-10)$$

where

$$\begin{aligned} f^l &:= -2\nabla\vartheta_k^l \cdot \nabla v^{l-1} - \Delta\vartheta_k^l v^{l-1} - (e_3 \wedge v^{l-1} + v^{l-1} \cdot \nabla v^{l-1})\vartheta_k^l + p^{l-1} \cdot \nabla\vartheta_k, \\ g^l &= v^{l-1} \cdot \nabla\vartheta_k, \\ \Sigma^l &= \theta_2^l(y_h - k)(\psi + \frac{1}{2}|v^{l-1}|^2 e_3|_{y_3=M}) \quad \text{on } \partial\Omega_k \cap \{y_3 = M\}, \\ \Sigma^l &= 0 \quad \text{on } \partial\Omega_k \cap \{y_3 = M\}^c. \end{aligned} \quad (4-11)$$

Now, Theorem IV.7.1 in [Boyer and Fabrie 2013] implies that for all $l \in \{2, \dots, m\}$, for $\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}$ small enough,

$$(v^l, p^l) \in H^l(\Omega_k) \times H^{l-1}(\Omega_k) \implies (v^{l+1}, p^{l+1}) \in H^{l+1}(\Omega_k) \times H^l(\Omega_k),$$

and

$$\|v^{l+1}\|_{H^{l+1}(\Omega_k)} + \|p^{l+1}\|_{H^l(\Omega_k)} \leq C(\|v^l\|_{H^l(\Omega_k)} + \|p^l\|_{H^{l-1}(\Omega_k)} + \|\psi\|_{H^{l-1/2}(\Omega_k)}).$$

Indeed, assume that $(v^l, p^l) \in H^l(\Omega_k) \times H^{l-1}(\Omega_k)$. Then $f^{l+1} \in H^{l-1}(\Omega_k)$, $g^{l+1} \in H^l(\Omega_k)$, with

$$\|f^{l+1}\|_{H^{l-1}(\Omega_k)} \leq C(\|v^l\|_{H^l} + \|v^l\|_{H^l}^2 + \|p^l\|_{H^{l-1}(\Omega_k)}), \quad \|g^{l+1}\|_{H^l} \leq C\|v^l\|_{H^l}.$$

Moreover, $v^l \in H^{l-1/2}(\partial\Omega_k)$. Since $l \geq 2$, using product laws in fractional Sobolev spaces (see [Strichartz 1967]), we infer that $|v^l|^2|_{y_3=M} \in H^{l-1/2}(\mathbb{R}^2)$, and therefore $\Sigma^{l+1} \in H^{l-1/2}(\mathbb{R}^2)$. From [Boyer and Fabrie 2013, Theorem IV.7.1], we deduce that $(v^{l+1}, p^{l+1}) \in H^{l+1}(\Omega_k) \times H^l(\Omega_k)$, together with the announced estimate. By induction, $v^- \in H_{\text{uloc}}^{m+1}(\Omega_{\text{bl}}^M)$ and $p^- \in H_{\text{uloc}}^m(\Omega_{\text{bl}}^M)$.

There only remains to check the compatibility condition at $y_3 = M$. Notice that

$$v_3^-|_{y_3=M} = \phi_3 + \int_{\gamma(y_h)}^M \partial_3 v_3^- = \phi_3 - \int_{\gamma(y_h)}^M \nabla_h \cdot v_h^- = \phi_3 - \gamma(y_h) \cdot \phi_h + \nabla_h \cdot v_h^*,$$

where

$$v_h^* = - \int_{\gamma(y_h)}^M v_h^- \in H_{\text{uloc}}^{1/2}(\mathbb{R}^2).$$

Since $\phi_3 - \gamma(y_h) \cdot \phi_h = 0$ due to the nonpenetrability condition $\phi \cdot n = 0$, we obtain the desired identity. \square

Step 2. Once (v^-, p^-) is defined thanks to Lemma 14, we define (v^+, p^+) in the half-space $\{y_3 > M\}$ by solving (2-4) with $v^+|_{y_3=M} = v^-|_{y_3=M}$. According to Lemma 14 and to standard trace inequalities,

$$\|v^-|_{y_3=M}\|_{H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}})$$

for some constant C depending only on M and on $\|\gamma\|_{W^{1,\infty}}$. As a consequence, if $C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}) + \|v_h^*\|_{L_{\text{uloc}}^2} \leq \delta_0$, according to Proposition 12 the system (2-4) with $v_0 = v^-|_{y_3=M}$ has a unique solution.

Additionally, $\Sigma(v^+, p^+)e_3|_{y_3=M^+}$ belongs to $H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$. Thus the mapping

$$\begin{aligned} \mathcal{F} : \mathcal{V} \times H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2) &\rightarrow H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2), \\ (\phi, \psi) &\mapsto \Sigma(v^+, p^+)e_3|_{y_3=M^+} - \psi, \end{aligned}$$

is well-defined. Clearly, according to [Lemma 14](#), for $\phi = 0$ and $\psi = 0$, we have $v^- = 0$, $v^+ = 0$ and therefore $\mathcal{F}(0, 0) = 0$.

The strategy is then to apply the implicit function theorem to \mathcal{F} to find a solution of $\mathcal{F}(\phi, \psi) = 0$ for ϕ in a neighbourhood of zero. Therefore we check that \mathcal{F} is \mathcal{C}^1 in a neighbourhood of zero, and that its Fréchet derivative with respect to ψ at $(0, 0)$ is an isomorphism on $H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$.

\mathcal{F} is a \mathcal{C}^1 mapping in a neighbourhood of zero: Let ϕ_0, ψ_0 and ϕ, ψ be in a neighbourhood of zero (in the sense of the functional norms in \mathcal{V} and $H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$). We denote by $v_0^\pm, p_0^\pm, v^\pm, p^\pm$ the solutions of (2-4), (2-5) associated with (ϕ_0, ψ_0) and $(\phi_0 + \phi, \psi_0 + \psi)$ respectively, and we set $w^\pm := v^\pm - v_0^\pm$, $q^\pm = p^\pm - p_0^\pm$.

On the one hand, in Ω_{bl}^M , we know w^- is a solution of the system

$$\begin{aligned} -\Delta w^- + e_3 \wedge w^- + (v_0^- + w^-) \cdot \nabla w^- + w^- \cdot \nabla v_0^- + \nabla q^- &= 0, \\ \operatorname{div} w^- &= 0, \\ w^-|_{\partial\Omega_{\text{bl}}} &= \phi, \\ \left(\partial_3 w^- - q^- e_3 - \frac{2v_0^- \cdot w^- + |w^-|^2}{2} e_3 \right) \Big|_{y_3=M} &= \psi. \end{aligned}$$

Performing estimates similar to the ones of [Lemma 14](#), we infer that for $\|\phi_0\|_{\mathcal{V}} + \|\psi_0\|_{H_{\text{uloc}}^{m-1/2}}$ and $\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}$ small enough,

$$\|w^-\|_{H_{\text{uloc}}^1(\Omega_{\text{bl}})} + \|w^-|_{y_3=M}\|_{H_{\text{uloc}}^{m+1/2}} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}).$$

It follows that

$$w^- = w_L^- + O(\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2)$$

in $H_{\text{uloc}}^1(\Omega_{\text{bl}}^M)$ and in $H_{\text{uloc}}^{m+1}((M', M) \times \mathbb{R}^2)$ for all $M' > \sup \gamma$, where w_L^- solves the same system as w^- minus the quadratic terms $w^- \cdot \nabla w^-$ and $|w^-|^2|_{y_3=M}$.

On the other hand, using [Theorem 2](#), one can show that $w^+ = w_L^+ + O(\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2)$, where

$$\begin{aligned} -\Delta w_L^+ + e_3 \wedge w^+ + v_0^+ \cdot \nabla w_L^+ + w_L^+ \cdot \nabla v_0^+ + \nabla q_L^+ &= 0 \quad \text{in } y_3 > M, \\ \operatorname{div} w_L^+ &= 0 \quad \text{in } y_3 > M, \\ w_L^+|_{y_3=M} &= w_L^-|_{y_3=M}. \end{aligned}$$

Using [Theorem 2](#), we deduce that if $\|(1 + y_3)^{1/3} v_0^+\|_{H_{\text{uloc}}^{m+1}}$ is small enough (which is ensured by the smallness condition on $\|\phi\|, \|\psi\|$), we have

$$\|(1 + y_3)^{1/3} w_L^+\|_{H_{\text{uloc}}^{m+1}(\mathbb{R}_+^3)} \leq C \|w_L^-|_{y_3=M}\|_{H_{\text{uloc}}^{m+1/2}} \leq C(\|\phi\|_{\mathcal{V}} + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}).$$

Therefore, in $H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$,

$$\mathcal{F}(\phi_0 + \phi, \psi_0 + \psi) - \mathcal{F}(\phi_0, \psi_0) = -\psi + \partial_3 w_L^+|_{y_3=M} - (q_L^+ + v_0^+ \cdot w_L^+)|_{y_3=M} e_3 + O(\|\phi\|_{\mathcal{V}}^2 + \|\psi\|_{H_{\text{uloc}}^{m-1/2}}^2).$$

It follows that the Fréchet derivative of \mathcal{F} at (ϕ_0, ψ_0) is

$$\mathcal{L}_{\phi_0, \psi_0} : (\phi, \psi) \mapsto -\psi + \partial_3 w_L^+|_{y_3=M} - (q_L^+ + v_0^+ \cdot w_L^+)|_{y_3=M} e_3.$$

Using the same kind of arguments as above, it is easily proved that w_L^\pm depend continuously on v_0^\pm , and therefore on ϕ_0, ψ_0 . Therefore \mathcal{F} is a C^1 function in a neighbourhood of zero.

$d_\psi \mathcal{F}(0, 0)$ is invertible: Since $d_\psi \mathcal{F}(0, 0) = \mathcal{L}_{0,0}(0, \cdot)$, we consider the systems solved by w_L^\pm with $v_0^\pm = 0$ and $\phi = 0$. We first notice that if $\mathcal{L}_{0,0}(0, \psi) = 0$, then $w_L := \mathbf{1}_{y_3 \leq M} w_L^- + \mathbf{1}_{y_3 > M} w_L^+$ is a solution of the Stokes–Coriolis system in the whole domain Ω_{bl} , with $w_L|_{\partial\Omega_{\text{bl}}} = 0$. Therefore, according to [Dalibard and Prange 2014], $w_L \equiv 0$ and therefore $\psi = 0$. Hence $\ker d_\psi \mathcal{F}(0, 0) = \{0\}$, and $d_\psi \mathcal{F}(0, 0)$ is one-to-one.

On the other hand,

$$(\partial_3 w_L^+ - q_L^+ e_3)|_{y_3=M} = \text{DN}(w_L^-|_{y_3=M}),$$

where DN is the Dirichlet-to-Neumann operator for the Stokes–Coriolis system, introduced in [loc. cit.]. In particular, in order to solve the equation

$$\mathcal{L}_{0,0}(0, \psi_1) = \psi_2$$

for a given $\psi_2 \in H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$, we need to solve the system

$$\begin{aligned} -\Delta w_L^- + e_3 \wedge w_L^- + \nabla q_L^- &= 0, \\ \operatorname{div} w_L^- &= 0, \\ w_L^-|_{\partial\Omega_{\text{bl}}} &= 0, \\ (\partial_3 w_L^- - q_L^- e_3)|_{y_3=M} &= -\psi_2 + \text{DN}(w_L^-|_{y_3=M}). \end{aligned}$$

According to Section 3 in [loc. cit.], the above system has a unique solution $w_L^- \in H_{\text{uloc}}^1(\Omega_{\text{bl}}^M)$. There only remains to prove that $w_L^- \in H_{\text{uloc}}^{m+1}(\{M' < y_3 < M\})$ for all $\sup \gamma < M' < M$. Therefore, we notice that in the domain $\mathbb{R}^2 \times (M', M)$, the horizontal derivatives of w_L^- (up to order m) satisfy a Stokes–Coriolis system similar to the one above (notice that the Dirichlet-to-Neumann operator commutes with ∂_1, ∂_2). It follows that $\nabla_h^\alpha w_L^- \in H_{\text{uloc}}^1(\mathbb{R}^2 \times (M', M))$ for all $|\alpha| \leq m$. In particular, $\nabla_h^\alpha w_L^-|_{y_3=M} \in H_{\text{uloc}}^{1/2}(\mathbb{R}^2)$ and therefore $w_L^-|_{y_3=M} \in H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2)$. It can be checked that $\text{DN} : H_{\text{uloc}}^{m+1/2}(\mathbb{R}^2) \rightarrow H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$. As a consequence, $\psi_1 = \partial_3 w_L^- - q_L^- e_3 \in H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$. Therefore $d_\psi \mathcal{F}(0, 0)$ is an isomorphism of $H^{m-1/2}(\mathbb{R}^2)$.

Using the implicit function theorem, we infer that for all $\phi \in \mathcal{V}$ in a neighbourhood of zero, there exists $\psi \in H_{\text{uloc}}^{m-1/2}(\mathbb{R}^2)$ such that $\mathcal{F}(\phi, \psi) = 0$. Let $v := \mathbf{1}_{y_3 \leq M} v^- + \mathbf{1}_{y_3 > M} v^+$, where v^-, v^+ are the solutions of (2-5)–(2-4) associated with ϕ, ψ . By definition, the jump of v across $\{y_3 = M\}$ is zero, and since $\mathcal{F}(\phi, \psi) = 0$,

$$\Sigma(v^-, p^-) e_3|_{y_3=M} = \psi = \Sigma(v^+, p^+) e_3|_{y_3=M}.$$

Using once again the fact that $|v^+|^2|_{y_3=M} = |v^-|^2|_{y_3=M}$, we deduce that

$$(\partial_3 v^- - p^- e_3)|_{y_3=M} = (\partial_3 v^+ - p^+ e_3)|_{y_3=M}.$$

Thus there is no jump of the stress tensor across $\{y_3 = M\}$, and therefore v is a solution of the Navier–Stokes–Coriolis system in the whole domain Ω_{bl} . This concludes the proof of [Theorem 1](#).

Appendix A: Proofs of Lemmas 7 and 9

Proof of Lemma 7. We begin with a few observations. First, replacing χ by $\chi_1 := Q\chi \in C_c^\infty(\mathbb{R}^2)$, it is enough to prove the lemma with $Q = 1$. Moreover it is clearly sufficient to prove the lemma for $p_k(\xi) = \xi_1^a \xi_2^b$, with $a + b = k$. Notice also that since $\alpha - k \geq -2$, we can always write $\alpha - k = 2m + \alpha_m$, with $\alpha_m \in [-2, 0[$ and $m \in \mathbb{N}$. Then $\xi_1^a \xi_2^b |\xi|^{\alpha-k}$ is a linear combination of terms of the form $\xi_1^{a'} \xi_2^{b'} |\xi|^{\alpha_m}$, with $a' + b' + \alpha_m = \alpha$ and $a', b' \in \mathbb{N}$. Therefore, in the rest of the proof, we take

$$Q \equiv 1, \quad P(\xi) = \xi_1^a \xi_2^b |\xi|^\beta, \quad \text{with } a, b \in \mathbb{N}, \beta \in [-2, 0[, a + b + \beta = \alpha.$$

Some of the arguments of the proof are inspired by the work of Alazard, Burq and Zuily [[Alazard et al. 2016](#)] on the Cauchy problem for gravity water waves in H_{uloc}^s spaces. We introduce a partition of unity $(\varphi_q)_{q \in \mathbb{Z}^2}$, where $\text{Supp } \varphi_q \subset B(q, 2)$ for $q \in \mathbb{Z}^2$ and $\sup_q \|\varphi_q\|_{W^{k,\infty}} < +\infty$ for all k . We also introduce functions $\tilde{\varphi}_q \in C_0^\infty(\mathbb{R}^2)$ such that $\tilde{\varphi}_q \equiv 1$ on $\text{Supp } \varphi_q$, and, say $\text{Supp } \tilde{\varphi}_q \subset B(q, 3)$. Then, for $j = 1, 2, 3$,

$$\begin{aligned} u^j(x_h, z) &= \sum_{q \in \mathbb{Z}^2} \chi(D) P(D) e^{-\lambda_j(D)z} (\varphi_q \underline{v}_0) \\ &= \sum_{q \in \mathbb{Z}^2} \int_{\mathbb{R}^2} K^j(x_h - y_h, z) \varphi_q(y_h) \underline{v}_0(y_h) dy_h = \sum_{q \in \mathbb{Z}^2} \int_{\mathbb{R}^2} K_q^j(x_h, y_h, z) \varphi_q(y_h) \underline{v}_0(y_h) dy_h, \end{aligned} \quad (\text{A-1})$$

where

$$K^j(x_h, z) = \int_{\mathbb{R}^2} e^{ix_h \cdot \xi} \chi(\xi) P(\xi) e^{-\lambda_j(\xi)z} d\xi, \quad K_q^j(x_h, y_h, z) = K^j(x_h - y_h, z) \tilde{\varphi}_q(y_h).$$

We then claim that the following estimates hold: there exists $\delta > 0, C \geq 0$ such that for all $x_h \in \mathbb{R}^2, z > 0$,

$$|K^1(x_h, z)| \leq \frac{C}{(1 + |x_h| + z^{1/3})^{2+\alpha}}, \quad |K^j(x_h, z)| \leq C \frac{e^{-\delta z}}{(1 + |x_h|)^{2+\alpha}} \quad \text{for } j = 2, 3. \quad (\text{A-2})$$

Let us postpone the proof of estimates [\(A-2\)](#) and explain why [Lemma 7](#) follows. Going back to [\(A-1\)](#), we have, for $j = 2, 3$,

$$\begin{aligned} |u^j(x_h, z)| &\leq C e^{-\delta z} \sum_{q \in \mathbb{Z}^2, |q - x_h| \geq 3} \frac{1}{(|q - x_h| - 2)^{2+\alpha}} \int |\varphi_q(y_h) \underline{v}_0(y_h)| dy_h \\ &\quad + C e^{-\delta z} \sum_{q \in \mathbb{Z}^2, |q - x_h| \leq 3} \int |\varphi_q(y_h) \underline{v}_0(y_h)| dy_h \\ &\leq C e^{-\delta z} \|\underline{v}_0\|_{L_{\text{uloc}}^1}. \end{aligned}$$

In a similar fashion,

$$\begin{aligned}
 |u^1(x_h, z)| &\leq C \sum_{q \in \mathbb{Z}^2, |q-x_h| \geq 3} \frac{1}{(|q-x_h|-2+z^{1/3})^{2+\alpha}} \int |\varphi_q(y_h) \underline{v}_0(y_h)| dy_h \\
 &\quad + C \sum_{q \in \mathbb{Z}^2, |q-x_h| \leq 3} \frac{1}{(1+z^{1/3})^{2+\alpha}} \int |\varphi_q(y_h) \underline{v}_0(y_h)| dy_h \\
 &\leq C \|\underline{v}_0\|_{L^1_{\text{uloc}}} (1+z)^{-\alpha/3}.
 \end{aligned}$$

The estimates of [Lemma 7](#) follow for $z \geq 1$.

We now turn to the proof of estimates [\(A-2\)](#). Once again we start with the estimates for K^2, K^3 , which are simpler. Since λ_2, λ_3 are continuous and have nonvanishing real part on the support of χ , there exists a constant $\delta > 0$ such that $\text{Re}(\lambda_j(\xi)) \geq \delta$ for all $\xi \in \text{Supp } \chi$ and for $j = 2, 3$. Clearly, for $|x_h| \leq 1$ we have simply

$$|K^j(x_h, z)| \leq e^{-\delta z} \|\chi P\|_{L^1}.$$

We thus focus on the set $|x_h| \geq 1$. Let $\chi_j(\xi, z) := \chi(\xi) \exp(-\lambda_j(\xi)z)$. Then $\chi_j \in L^\infty(\mathbb{R}_+, \mathcal{S}(\mathbb{R}^2))$, and for all $n_1, n_2, n_3 \in \mathbb{N}$, there exists a constant $\delta_n > 0$ such that

$$|(1+|\xi|^{n_3})\partial_1^{n_1}\partial_2^{n_2}\chi_j(\xi, z)| \leq C_n \exp(-\delta_n z).$$

Estimate [\(A-2\)](#) for K^2, K^3 then follows immediately from the following lemma (whose proof is given after the current one):

Lemma 15. *Let $P(\xi) = \xi_1^{a_1} \xi_2^{a_2} |\xi|^\beta$, with $a_1, a_2 \in \mathbb{N}$, $\beta \in [-2, 0[$, and set $\alpha := a_1 + a_2 + \beta$. Then there exists $C > 0$ such that for any $\zeta \in \mathcal{S}(\mathbb{R}^2)$, for all $x_h \in \mathbb{R}^2$, $|x_h| \geq 1$,*

$$|P(D)\zeta(x_h)| \leq \frac{C}{|x_h|^{2+\alpha}} (\|\zeta\|_1 + \||y_h|^{a_1+a_2+2}\partial_1^{a_1}\partial_2^{a_2}\zeta\|_\infty).$$

We now address the estimates on K^1 . When $|x_h| \leq 1$, $z \leq 1$, we have simply $|K^1(x_h, z)| \leq \|P\chi\|_1$, and the estimate follows. When $z \leq 1$ and $|x_h| \geq 1$, we apply [Lemma 15](#) with $\zeta(\xi) = \mathcal{F}^{-1}(\chi(\xi) \exp(-\lambda_1(\xi)z))$. Notice that

$$\|\zeta\|_1 \lesssim \|\chi(\xi) \exp(-\lambda_1(\xi)z)\|_{W^{3,1}},$$

and

$$\||y_h|^{a_1+a_2+2}\partial_1^{a_1}\partial_2^{a_2}\zeta\|_\infty \lesssim \|\xi_1^{a_1}\xi_2^{a_2}\chi(\xi) \exp(-\lambda_1(\xi)z)\|_{W^{2+a_1+a_2,1}}.$$

Since the right-hand sides of the above inequalities are bounded (recall that $\lambda_1(\xi) = |\xi|^3 \Lambda_1(\xi)$ with $\Lambda_1 \in C^\infty(\mathbb{R}^2)$; see [Remark 4](#)), it follows that estimate [\(A-2\)](#) is true for $z \leq 1$ and $|x_h| \geq 1$.

We now focus on the case $z \geq 1$. We first change variables in the integral defining K^1 and we set $\xi' = z^{1/3}\xi$, $x'_h = x_h/z^{1/3}$. Since P is homogeneous, this leads to

$$K^1(x_h, z) = \frac{1}{z^{(2+\alpha)/3}} \int_{\mathbb{R}^2} e^{ix'_h \cdot \xi'} P(\xi') \chi\left(\frac{\xi'}{z^{1/3}}\right) \exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right) d\xi'.$$

Since $\lambda_1/|\xi|^3$ is continuous and does not vanish on the support of χ , there exists a positive constant δ' such that $\lambda_1(\xi) \geq \delta'|\xi|^3$ on $\text{Supp } \chi$. Therefore, for $|x'_h| \leq 1$, we have

$$|K^1(x_h, z)| \leq \frac{1}{z^{(2+\alpha)/3}} \left\| \exp(-\delta'|\xi|^3) P(\xi') \right\|_{L^1},$$

and the estimate for K^1 on the set $|x_h| \leq z^{1/3}$ is proved.

For $|x'_h| \geq 1$, we split the integral in two. Let $\varphi \in C_0^\infty$ such that $\varphi \equiv 1$ in a neighbourhood of zero. Then

$$\begin{aligned} K^1(x_h, z) &= \frac{1}{z^{(2+\alpha)/3}} \int_{\mathbb{R}^2} e^{ix'_h \cdot \xi'} P(\xi') \varphi(\xi') \chi\left(\frac{\xi'}{z^{1/3}}\right) \exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right) d\xi' \\ &\quad + \frac{1}{z^{(2+\alpha)/3}} \int_{\mathbb{R}^2} e^{ix'_h \cdot \xi'} P(\xi') (1 - \varphi(\xi')) \chi\left(\frac{\xi'}{z^{1/3}}\right) \exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right) d\xi' \\ &=: K_1^1 + K_2^1. \end{aligned}$$

We first consider the term K_2^1 . Because of the truncation $1 - \varphi$, we have removed all singularity coming from P close to $\xi = 0$. Therefore, performing integrations by parts, we have, for any $n \in \mathbb{N}$, for $j = 1, 2$,

$$x_j'^n K_2^1(x_h, z) = \frac{1}{z^{(2+\alpha)/3}} \int_{\mathbb{R}^2} e^{ix'_h \cdot \xi'} D_{\xi_j'}^n \left[P(\xi') (1 - \varphi(\xi')) \chi\left(\frac{\xi'}{z^{1/3}}\right) \exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right) \right] d\xi'.$$

When the $D_{\xi_j'}$ derivative hits $P(1 - \varphi)$, we end up with an integral bounded by

$$C_n \int_{\mathbb{R}^2} |\xi'|^\alpha \mathbf{1}_{\xi' \in \text{Supp}(1-\varphi)} \exp(-\delta'|\xi'|^3) d\xi' \leq C_n.$$

When the derivative hits $\chi(\xi'/z^{1/3})$ the situation is even better, as a power of $z^{1/3}$ is gained with each derivative. Therefore the worst terms occur when the derivative hits the exponential. Remember that $\lambda_1(\xi) = |\xi|^3 \Lambda_1(\xi)$, where $\Lambda_1 \in C^\infty(\mathbb{R}^2)$ with $\Lambda_1(0) = 1$ and Λ_1 does not vanish on \mathbb{R}^2 . Therefore, for all $\xi' \in \mathbb{R}^2$, $z > 0$,

$$\exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right) = \exp\left(-|\xi'|^3 \Lambda_1\left(\frac{\xi'}{z^{1/3}}\right)\right).$$

We infer that for any $0 \leq n \leq 3 + \lfloor \alpha \rfloor$, on $\text{Supp } \chi(\cdot/z^{1/3})$, we have

$$\left| P(\xi') \nabla_{\xi_j'}^n \exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right) \right| \leq C_n \exp\left(-\frac{\delta'}{2}|\xi|^3\right). \quad (\text{A-3})$$

We deduce eventually that

$$|K_2^1(x_h, z)| \leq C \frac{1}{z^{(2+\alpha)/3}} \frac{1}{(1 + |x'_h|^{2+\alpha})} \leq \frac{C}{(|x_h| + z^{1/3})^{2+\alpha}}.$$

For the term K_1^1 , we use once again [Lemma 15](#), with

$$\zeta := \mathcal{F}^{-1}\left(\varphi(\xi') \chi\left(\frac{\xi'}{z^{1/3}}\right) \exp\left(-\lambda_1\left(\frac{\xi'}{z^{1/3}}\right)z\right)\right).$$

Using the same type of estimate as (A-3) above, we obtain

$$|K_2^1(x_h, z)| \leq C \frac{1}{z^{(2+\alpha)/3}} \frac{1}{|x'_h|^{2+\alpha}} \leq \frac{C}{|x_h|^{2+\alpha}}.$$

This concludes the proof of Lemma 7. \square

Proof of Lemma 15. We have

$$P(D)\zeta = D_1^{a_1} D_2^{a_2} \text{Op}(|\xi|^\beta)\zeta.$$

Thus we first compute $\text{Op}(|\xi|^\beta)\zeta$. We first focus on the case $\beta \in]-2, 0[$. We follow the ideas of Droniou and Imbert [2006, Theorem 1], recalling the main steps of the proof. The function $\xi \in \mathbb{R}^2 \mapsto |\xi|^\beta$ is radial and locally integrable, and thus belongs to \mathcal{S}' . Its Fourier transform in $\mathcal{S}'(\mathbb{R}^2)$ is also radial and homogeneous of degree $-\beta-2 \in]-2, 0[$. Therefore it coincides (up to a constant) with $|\cdot|^{-\beta-2}$ in $\mathcal{S}'(\mathbb{R}^2 \setminus \{0\})$, and since the latter function is locally integrable, we end up with $\mathcal{F}^{-1}(|\xi|^\beta) = C|x_h|^{-\beta-2}$ in $\mathcal{S}'(\mathbb{R}^N)$. Hence

$$P(D)\zeta(x_h) = C \partial_1^{a_1} \partial_2^{a_2} \int_{\mathbb{R}^2} \frac{1}{|y_h|^{\beta+2}} \zeta(x_h - y_h) dy_h.$$

Notice that in the present case, we do not need to have an exact formula for $P(D)\zeta$, but merely some information about its decay at infinity. As a consequence we take a shortcut in the proof of [Droniou and Imbert 2006]. We take a cut-off function $\chi \in C_0^\infty(\mathbb{R}^2)$ such that $\chi \equiv 1$ in a neighbourhood of zero, and we write

$$\begin{aligned} P(D)\zeta(x_h) &= C \int_{\mathbb{R}^2} \frac{\chi(y_h)}{|y_h|^{\beta+2}} \partial_1^{a_1} \partial_2^{a_2} \zeta(x_h - y_h) dy_h \\ &\quad + C \sum_{\substack{0 \leq i_1 \leq a_1 \\ 0 \leq i_2 \leq a_2}} C_{i_1, i_2} \int_{\mathbb{R}^2} \partial_1^{i_1} \partial_2^{i_2} (1 - \chi(y_h)) \partial_1^{a_1 - i_1} \partial_2^{a_2 - i_2} \left(\frac{1}{|y_h|^{\beta+2}} \right) \zeta(x_h - y_h) dy_h \\ &=: I_1 + I_2. \end{aligned}$$

We now choose χ in the following way. Let $n = \lfloor |x_h| \rfloor \in \mathbb{N}$, and take $\chi = \chi_n = \eta(\cdot/n)$, where $\text{Supp } \eta \subset B(0, \frac{1}{2})$ and $\eta \equiv 1$ in a neighbourhood of zero. Notice in that case that if $y_h \in \text{Supp } \chi_n$, then $|x_h - y_h| \geq |x_h|/2$. Therefore, for the first term, we have

$$|x_h|^{2+\alpha} |I_1| \leq (n+1)^\beta \left(\int_{|y_h| \leq n/2} |y_h|^{-\beta-2} dy_h \right) \| |y_h|^{2+a_1+a_2} \partial_1^{a_1} \partial_2^{a_2} \zeta \|_{L^\infty} \leq C \| |y_h|^{2+a_1+a_2} \partial_1^{a_1} \partial_2^{a_2} \zeta \|_{L^\infty}.$$

Using the assumptions on η and χ_n and the estimate

$$\left| \partial_1^{a_1 - i_1} \partial_2^{a_2 - i_2} \left(\frac{1}{|y_h|^{\beta+2}} \right) \right| \leq \frac{C}{|y_h|^{\alpha+2-i_1-i_2}} \leq \frac{C}{n^{\alpha+2-i_1-i_2}} \quad \forall y_h \in \text{Supp}(1 - \chi_n),$$

we infer that

$$|I_2| \leq C \|\zeta\|_{L^1} n^{-\alpha-2} \leq C \|\zeta\|_{L^1} |x_h|^{-\alpha-2}.$$

Gathering all the terms, we obtain the inequality announced in the lemma. To conclude the proof, we still have to consider the case $\beta = -2$: in such a case, $|\xi|^\beta$ corresponds to inverting the Laplacian over \mathbb{R}^2 . Hence, the kernel $|x_h - y_h|^{-\beta-2}$ has to be replaced by $\frac{1}{2\pi} \ln(|x_h - y_h|)$. This does not modify the previous reasoning. \square

Proof of Lemma 9. The proof is somewhat simpler than the one of Lemma 7. As indicated in the remark following Lemma 9, notice that for $n > 1$, for all $\xi \in \mathbb{R}^2$, $z > 0$,

$$\left| (1 + |\xi|^2)^{-n} (1 - \chi(\xi)) P(\xi) e^{-\lambda_j(\xi)z} \right| \leq \|P\|_{L^\infty(B_r^c)} \frac{e^{-\delta z}}{(1 + |\xi|^2)^n},$$

and the right-hand side of the above inequality is in $L^1(\mathbb{R}^2)$ for all z . As a consequence, for $j = 1, \dots, 3$, $n > 1$, the kernel

$$K_{n,j}(x_h, z) := \int_{\mathbb{R}^2} e^{ix_h \cdot \xi} (1 + |\xi|^2)^{-n} (1 - \chi)(\xi) P(\xi) \exp(-\lambda_j(\xi)z) d\xi$$

is well-defined and satisfies

$$\|K_{n,j}(\cdot, z)\|_{L^\infty(\mathbb{R}^2)} \leq C_n \|P\|_{L^\infty(B_r^c)} e^{-\delta z}.$$

Furthermore, if $a_1, a_2 \in \mathbb{N}$ with $a_1 + a_2 \leq 3$,

$$x_1^{a_1} x_2^{a_2} K_{n,j}(x_h, z) = \int_{\mathbb{R}^2} e^{ix_h \cdot \xi} D_1^{a_1} D_2^{a_2} \left((1 + |\xi|^2)^{-n} (1 - \chi)(\xi) P(\xi) \exp(-\lambda_j(\xi)z) \right) d\xi.$$

Hence, up to taking a larger n and a smaller δ ,

$$|K_{n,j}(x_h, z)| \leq C_n \|P\|_{W^{3,\infty}(B_r^c)} e^{-\delta z} (1 + |x_h|)^{-3},$$

and in particular, $K_{n,j} \in L_z^\infty(L_{x_h}^2)$. Thus for any $f \in L_{\text{uloc}}^2$,

$$\|(1 + |D|^2)^{-n} (1 - \chi(D)) P(D) \exp(-\lambda_j(D)z) f\|_{L^\infty} = \|K_{n,j} * f\|_{L^\infty} \leq C e^{-\delta z} \|f\|_{L_{\text{uloc}}^2}.$$

Taking $f = (1 + |D|^2)^n \underline{v}_0 = (1 - \Delta_h)^n \underline{v}_0$ for some $\underline{v}_0 \in H_{\text{uloc}}^{2n}$, we obtain the result announced in Lemma 9.

Appendix B. Estimates on a few integrals

Lemma 16. *There exists a positive constant C such that for all $z \geq 0$,*

$$\int_0^\infty \frac{1}{(1 + |z - z'|)^{2/3} (1 + z')^{2/3}} dz' \leq \frac{C}{(1 + z)^{1/3}},$$

$$\int_0^\infty \frac{1}{(1 + |z - z'|)(1 + z')^{2/3}} dz' \leq \frac{C \ln(2 + z)}{(1 + z)^{2/3}},$$

and for all $\gamma, \delta > 0$ such that $\delta < 1$ and $\gamma + \delta > 1$, there exists a constant $C_{\gamma,\delta}$ such that

$$\int_0^\infty \frac{1}{(1 + z + z')^\gamma} \frac{1}{(1 + z')^\delta} dz' \leq \frac{C_{\gamma,\delta}}{(1 + z)^{\gamma+\delta}} \quad \forall z \geq 0.$$

Proof. The first two inequalities are obvious if z is small (say, $z \leq \frac{1}{2}$), simply by writing

$$\frac{1}{1 + |z - z'|} \leq \frac{C}{1 + z'}.$$

Hence we focus on $z' \geq \frac{1}{2}$. In that case, changing variables in the first integral, we have

$$\int_0^\infty \frac{1}{(1 + |z - z'|)^{2/3} (1 + z')^{2/3}} dz' = \frac{1}{z^{1/3}} \int_0^\infty \frac{1}{(z^{-1} + |1 - t|)^{2/3}} \frac{1}{(z^{-1} + t)^{2/3}} dt \leq \frac{1}{z^{1/3}} \int_0^\infty \frac{1}{|1 - t|^{2/3}} \frac{1}{t^{2/3}} dt,$$

which proves the first inequality. The second one is treated in a similar fashion:

$$\int_0^\infty \frac{1}{1+|z-z'|} \frac{1}{(1+z)^{2/3}} dz' = z^{-1} \int_0^\infty \frac{1}{z^{-1}+|1-t|} \frac{1}{(z^{-1}+t)^{2/3}} dt \leq z^{-1} \int_0^\infty \frac{1}{z^{-1}+|1-t|} \frac{1}{t^{2/3}} dt.$$

It is easily checked that

$$\int_{1/2}^{3/2} \frac{1}{z^{-1}+|1-t|} dt \leq C \ln(2+z).$$

The second estimate follows. The last estimate is proved by similar arguments and is left to the reader. \square

Acknowledgements

The authors have been partially funded by the ANR project Dyficolti ANR-13-BS01-0003-01. Anne-Laure Dalibard has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant agreement No. 63765).

References

- [Achdou et al. 1998a] Y. Achdou, P. Le Tallec, F. Valentin, and O. Pironneau, “Constructing wall laws with domain decomposition or asymptotic expansion techniques”, *Comput. Methods Appl. Mech. Engrg.* **151**:1–2 (1998), 215–232. [MR](#) [Zbl](#)
- [Achdou et al. 1998b] Y. Achdou, B. Mohammadi, O. Pironneau, and F. Valentin, “Domain decomposition & wall laws”, pp. 1–14 in *Recent developments in domain decomposition methods and flow problems* (Kyoto, 1996; Anacapri, 1996), edited by H. Fujita et al., GAKUTO Internat. Ser. Math. Sci. Appl. **11**, Gakkōtoshō, Tokyo, 1998. [MR](#) [Zbl](#)
- [Achdou et al. 1998c] Y. Achdou, O. Pironneau, and F. Valentin, “Effective boundary conditions for laminar flows over periodic rough boundaries”, *J. Comput. Phys.* **147**:1 (1998), 187–218. [MR](#) [Zbl](#)
- [Alazard et al. 2016] T. Alazard, N. Burq, and C. Zuily, “Cauchy theory for the gravity water waves system with non-localized initial data”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **33**:2 (2016), 337–395. [MR](#) [Zbl](#)
- [Amirat et al. 2001] Y. Amirat, D. Bresch, J. Lemoine, and J. Simon, “Effect of rugosity on a flow governed by stationary Navier–Stokes equations”, *Quart. Appl. Math.* **59**:4 (2001), 769–785. [MR](#) [Zbl](#)
- [Basson and Gérard-Varet 2008] A. Basson and D. Gérard-Varet, “Wall laws for fluid flows at a boundary with random roughness”, *Comm. Pure Appl. Math.* **61**:7 (2008), 941–987. [MR](#) [Zbl](#)
- [Bocquet and Barrat 2007] L. Bocquet and J.-L. Barrat, “Flow boundary conditions from nano- to micro-scales”, *Soft Matter* **3**:6 (2007), 685–693.
- [Bonnivard and Bucur 2012] M. Bonnivard and D. Bucur, “The uniform rugosity effect”, *J. Math. Fluid Mech.* **14**:2 (2012), 201–215. [MR](#) [Zbl](#)
- [Boyer and Fabrie 2013] F. Boyer and P. Fabrie, *Mathematical tools for the study of the incompressible Navier–Stokes equations and related models*, Applied Mathematical Sciences **183**, Springer, 2013. [MR](#) [Zbl](#)
- [Bresch and Milisic 2010] D. Bresch and V. Milisic, “High order multi-scale wall-laws, Part I: The periodic case”, *Quart. Appl. Math.* **68**:2 (2010), 229–253. [MR](#) [Zbl](#)
- [Bucur et al. 2008] D. Bucur, E. Feireisl, Š. Nečasová, and J. Wolf, “On the asymptotic limit of the Navier–Stokes system on domains with rough boundaries”, *J. Differential Equations* **244**:11 (2008), 2890–2908. [MR](#) [Zbl](#)
- [Casado-Díaz et al. 2003] J. Casado-Díaz, E. Fernández-Cara, and J. Simon, “Why viscous fluids adhere to rugose walls: a mathematical explanation”, *J. Differential Equations* **189**:2 (2003), 526–537. [MR](#) [Zbl](#)
- [Dalibard and Gérard-Varet 2011] A.-L. Dalibard and D. Gérard-Varet, “Effective boundary condition at a rough surface starting from a slip condition”, *J. Differential Equations* **251**:12 (2011), 3450–3487. [MR](#) [Zbl](#)
- [Dalibard and Prange 2014] A.-L. Dalibard and C. Prange, “Well-posedness of the Stokes–Coriolis system in the half-space over a rough surface”, *Anal. PDE* **7**:6 (2014), 1253–1315. [MR](#) [Zbl](#)

- [Droniou and Imbert 2006] J. Droniou and C. Imbert, “Fractal first-order partial differential equations”, *Arch. Ration. Mech. Anal.* **182**:2 (2006), 299–331. [MR](#) [Zbl](#)
- [Galdi 2011] G. P. Galdi, *An introduction to the mathematical theory of the Navier–Stokes equations: steady-state problems*, 2nd ed., Springer, 2011. [MR](#) [Zbl](#)
- [Gérard-Varet 2003] D. Gérard-Varet, “Highly rotating fluids in rough domains”, *J. Math. Pures Appl.* (9) **82**:11 (2003), 1453–1498. [MR](#) [Zbl](#)
- [Gérard-Varet 2009] D. Gérard-Varet, “The Navier wall law at a boundary with random roughness”, *Comm. Math. Phys.* **286**:1 (2009), 81–110. [MR](#) [Zbl](#)
- [Gérard-Varet and Dormy 2006] D. Gérard-Varet and E. Dormy, “Ekman layers near wavy boundaries”, *J. Fluid Mech.* **565** (2006), 115–134. [MR](#) [Zbl](#)
- [Gérard-Varet and Masmoudi 2010] D. Gérard-Varet and N. Masmoudi, “Relevance of the slip condition for fluid flows near an irregular boundary”, *Comm. Math. Phys.* **295**:1 (2010), 99–137. [MR](#) [Zbl](#)
- [Jäger and Mikelić 2001] W. Jäger and A. Mikelić, “On the roughness-induced effective boundary conditions for an incompressible viscous flow”, *J. Differential Equations* **170**:1 (2001), 96–122. [MR](#) [Zbl](#)
- [Jäger and Mikelić 2003] W. Jäger and A. Mikelić, “Couette flows over a rough boundary and drag reduction”, *Comm. Math. Phys.* **232**:3 (2003), 429–455. [MR](#) [Zbl](#)
- [Ladyženskaja and Solonnikov 1980] O. A. Ladyženskaja and V. A. Solonnikov, “Determination of solutions of boundary value problems for stationary Stokes and Navier–Stokes equations having an unbounded Dirichlet integral”, *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **96** (1980), 117–160. In Russian; translated in *J. Soviet Math.* **21**:5 (1983), 728–851. [MR](#)
- [Lauga et al. 2007] E. Lauga, M. Brenner, and H. Stone, “Microfluidics: the no-slip boundary condition”, pp. 1219–1240 in *Handbook of Experimental Fluid Mechanics*, edited by C. Tropea et al., Springer, Berlin, 2007.
- [Mikelić et al. 2013] A. Mikelić, Š. Nečasová, and M. Neuss-Radu, “Effective slip law for general viscous flows over an oscillating surface”, *Math. Methods Appl. Sci.* **36**:15 (2013), 2086–2100. [MR](#) [Zbl](#)
- [Neuss et al. 2006] N. Neuss, M. Neuss-Radu, and A. Mikelić, “Effective laws for the Poisson equation on domains with curved oscillating boundaries”, *Appl. Anal.* **85**:5 (2006), 479–502. [MR](#) [Zbl](#)
- [Pedlosky 1987] J. Pedlosky, *Geophysical fluid dynamics*, 2nd ed., Springer, New York, 1987. [Zbl](#)
- [Prange 2013] C. Prange, “Asymptotic analysis of boundary layer correctors in periodic homogenization”, *SIAM J. Math. Anal.* **45**:1 (2013), 345–387. [MR](#) [Zbl](#)
- [Strichartz 1967] R. S. Strichartz, “Multipliers on fractional Sobolev spaces”, *J. Math. Mech.* **16** (1967), 1031–1060. [MR](#) [Zbl](#)
- [Vinogradova and Yakubov 2006] O. Vinogradova and G. Yakubov, “Surface roughness and hydrodynamic boundary conditions”, *Phys. Rev. E* **73**:4 (2006), art. id. 045302.
- [Wheeler and Horgan 1976] L. T. Wheeler and C. O. Horgan, “A two-dimensional Saint-Venant principle for second-order linear elliptic equations”, *Quart. Appl. Math.* **34**:3 (1976), 257–270. [MR](#) [Zbl](#)
- [Wheeler et al. 1975] L. T. Wheeler, M. J. Turteltaub, and C. O. Horgan, “A Saint-Venant principle for the gradient in the Neumann problem”, *Z. Angew. Math. Phys.* **26** (1975), 141–153. [MR](#) [Zbl](#)
- [Ybert et al. 2007] C. Ybert, C. Barentin, C. Cottin-Bizonne, P. Joseph, and L. Bocquet, “Achieving large slip with superhydrophobic surfaces: scaling laws for generic geometries”, *Phys. of Fluids* **19**:12 (2007), art. id. 123601. [Zbl](#)

Received 3 Nov 2015. Revised 12 May 2016. Accepted 13 Oct 2016.

ANNE-LAURE DALIBARD: dalibard@ann.jussieu.fr

Sorbonne Universités, UPMC Université Paris 06, CNRS, UMR 7598, Laboratoire Jacques-Louis Lions, 4 place Jussieu, 75005 Paris, France

DAVID GÉRARD-VARET: david.gerard-varet@imj-prg.fr

Université Paris Diderot, Sorbonne Paris Cité, Institut de Mathématiques de Jussieu-Paris Rive Gauche, UMR 7586, F-75205 Paris, France

Analysis & PDE

msp.org/apde

EDITORS

EDITOR-IN-CHIEF

Patrick Gérard

patrick.gerard@math.u-psud.fr

Université Paris Sud XI

Orsay, France

BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbb@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or msp.org/apde for submission instructions.

The subscription price for 2017 is US \$265/year for the electronic version, and \$470/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

APDE peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

ANALYSIS & PDE

Volume 10 No. 1 2017

Nonlinear boundary layers for rotating fluids	1
ANNE-LAURE DALIBARD and DAVID GÉRARD-VARET	
Partial data inverse problems for the Hodge Laplacian	43
FRANCIS J. CHUNG, MIKKO SALO and LEO TZOU	
On an isoperimetric-isodiametric inequality	95
ANDREA MONDINO and EMANUELE SPADARO	
Nonradial type II blow up for the energy-supercritical semilinear heat equation	127
CHARLES COLLOT	