



Innovations in Incidence Geometry

Algebraic, Topological and Combinatorial



The exterior splash in $PG(6, q)$: transversals

Susan G. Barwick and Wen-Ai Jackson



The exterior splash in $\text{PG}(6, q)$: transversals

Susan G. Barwick and Wen-Ai Jackson

Let π be an order- q -subplane of $\text{PG}(2, q^3)$ that is exterior to ℓ_∞ . Then the exterior splash of π is the set of $q^2 + q + 1$ points on ℓ_∞ that lie on an extended line of π . Exterior splashes are projectively equivalent to scattered linear sets of rank 3, covers of the circle geometry $CG(3, q)$, and hyper-reguli in $\text{PG}(5, q)$. We use the Bruck–Bose representation in $\text{PG}(6, q)$ to investigate the structure of π , and the interaction between π and its exterior splash. We show that the point set of $\text{PG}(6, q)$ corresponding to π is the intersection of nine quadrics, and that there is a unique tangent plane at each point, namely the intersection of the tangent spaces of the nine quadrics. In $\text{PG}(6, q)$, an exterior splash \mathbb{S} has two sets of cover planes (which are hyper-reguli) and we show that each set has three unique transversal lines in the cubic extension $\text{PG}(6, q^3)$. These transversal lines are used to characterise the carriers and the sublines of \mathbb{S} .

1. Introduction

In [Barwick and Jackson 2012; 2014], we studied order- q -subplanes of $\text{PG}(2, q^3)$ and determined their representation in the Bruck–Bose representation in $\text{PG}(6, q)$. A full characterisation in $\text{PG}(6, q)$ was given for order- q -subplanes that are secant or tangent to ℓ_∞ in $\text{PG}(2, q^3)$. In [Rottey et al. 2015], this was generalised to study subplanes of $\text{PG}(2, q^n)$ in $\text{PG}(2n, q)$. The cases when the subplane is secant or tangent to ℓ_∞ yield nice geometric characterisations. However, the case of an order- q -subplane π of $\text{PG}(2, q^3)$ that is exterior to ℓ_∞ yields a complex structure denoted $[\pi]$ in $\text{PG}(6, q)$. Our main motivation in this article is to investigate the geometric properties of the structure $[\pi]$. The splash of π gives crucial information about the geometrical properties of $[\pi]$, and so we also study the interplay in $\text{PG}(6, q)$ between $[\pi]$ and its splash.

The splash of a subplane π of $\text{PG}(2, q^n)$ is defined to be the set of points on ℓ_∞ that lie on an extended line of π . In [Barwick and Jackson 2015] it was shown that

MSC2010: 51E20.

Keywords: Bruck–Bose representation, subplanes, exterior splash, linear sets.

the splash of a tangent order- q -subplane of $\text{PG}(2, q^3)$ is a linear set. In [Lavrauw and Zanella 2015] the notion of splash was generalised from subplanes to subgeometries, and to general field extensions. It was shown that a splash is a linear set, and conversely, a linear set is a splash.

In this article we let π be a subplane of $\text{PG}(2, q^3)$ of order q that is exterior to ℓ_∞ . The lines of π meet ℓ_∞ in a set \mathbb{S} of size $q^2 + q + 1$, which we call the *exterior splash* of π . Properties of the exterior splash of $\text{PG}(2, q^3)$ were studied in [Barwick and Jackson 2016]. The sets of points in an exterior splash has arisen in many different situations, namely scattered \mathbb{F}_q -linear sets of rank 3, covers of the circle geometry $\text{CG}(3, q)$, hyper-reguli in $\text{PG}(5, q)$, and Sherk surfaces of size $q^2 + q + 1$. Scattered linear sets are surveyed in [Lavrauw 2016]. An important result is that all scattered \mathbb{F}_q -linear sets of rank 3 are projectively equivalent [Lavrauw and Zanella 2015].

This article proceeds as follows. In Section 2 we introduce the notation we use for the Bruck–Bose representation of $\text{PG}(2, q^3)$ in $\text{PG}(6, q)$, as well as presenting some other preliminary results.

We next introduce coordinates; as all scattered \mathbb{F}_q -linear sets of rank 3 are projectively equivalent, we will work with an exterior splash equivalent to the set of points

$$\{(x, x^q) : x \in \text{GF}(q^3) \setminus \{0\}\}.$$

In Section 3 we coordinatise an order- q -subplane \mathcal{B} in $\text{PG}(2, q^3)$ that is exterior to ℓ_∞ , with this exterior splash. This order- q -subplane will be used in many of the proofs in this article.

In Section 4, we study the structure of an order- q -subplane in $\text{PG}(6, q)$. We show that it contains $q^2 + q + 1$ twisted cubics and is the intersection of nine quadrics. Further, we show that there is a unique tangent plane at each point, which is the intersection of the tangent spaces of these nine quadrics.

We next study the exterior splash \mathbb{S} of ℓ_∞ in the Bruck–Bose representation in $\text{PG}(5, q)$. By results of Bruck [1973], \mathbb{S} has two switching sets denoted \mathbb{X}, \mathbb{Y} , which we call covers of \mathbb{S} . The three sets $\mathbb{S}, \mathbb{X}, \mathbb{Y}$ are called hyper-reguli in [Ostrom 1993]. In Section 5, we look at the exterior splash

$$\{(x, x^q) : x \in \text{GF}(q^3) \setminus \{0\}\},$$

and working in $\text{PG}(6, q)$, find coordinates for the two covers \mathbb{X}, \mathbb{Y} . In Section 6, we show that each of the sets $\mathbb{S}, \mathbb{X}, \mathbb{Y}$ has a unique triple of conjugate transversal lines in the cubic extension $\text{PG}(5, q^3)$. Theorem 6.5 characterises the carriers of an exterior splash as the only planes of the regular spread that meet all nine transversal lines. Theorem 6.6 shows that the nine transversal lines are common to the set of $q - 1$ disjoint splashes of ℓ_∞ that have common carriers. We interpret this result in terms of replacing hyper-reguli to create André planes. In Section 7 we use the transversal lines to characterise the order- q -sublines of an exterior splash in terms of how the corresponding 2-reguli meet the cover planes.

2. The Bruck–Bose representation

2A. The Bruck–Bose representation of $\text{PG}(2, q^3)$ in $\text{PG}(6, q)$. We introduce the notation we will use for the Bruck–Bose representation of $\text{PG}(2, q^3)$ in $\text{PG}(6, q)$. We work with the finite field \mathbb{F}_q of order q . A *2-spread* of $\text{PG}(5, q)$ is a set of $q^3 + 1$ planes that partition $\text{PG}(5, q)$. A *2-regulus* of $\text{PG}(5, q)$ is a set of $q + 1$ mutually disjoint planes π_1, \dots, π_{q+1} with the property that if a line meets three of the planes, then it meets all $q + 1$ of them. A 2-regulus \mathcal{R} has a set of $q^2 + q + 1$ mutually disjoint *ruling lines* that meet every plane of \mathcal{R} . A 2-regulus is uniquely determined by three mutually disjoint planes, or four (ruling) lines (mutually disjoint and lying in general position). A 2-spread \mathcal{S} is *regular* if for any three planes in \mathcal{S} , the 2-regulus containing them is contained in \mathcal{S} . See [Hirschfeld and Thas 1991] for more information on 2-spreads.

The following construction of a regular 2-spread of $\text{PG}(5, q)$ will be needed. Embed $\text{PG}(5, q)$ in $\text{PG}(5, q^3)$ and let g be a line of $\text{PG}(5, q^3)$ disjoint from $\text{PG}(5, q)$. Let g^q, g^{q^2} be the conjugate lines of g ; both of these are disjoint from $\text{PG}(5, q)$. Let P_i be a point on g ; then the plane $\langle P_i, P_i^q, P_i^{q^2} \rangle$ meets $\text{PG}(5, q)$ in a plane. As P_i ranges over all the points of g , we get $q^3 + 1$ planes of $\text{PG}(5, q)$ that partition $\text{PG}(5, q)$. These planes form a regular 2-spread \mathcal{S} of $\text{PG}(5, q)$. The lines g, g^q, g^{q^2} are called the (conjugate skew) *transversal lines* of the 2-spread \mathcal{S} . Conversely, given a regular 2-spread in $\text{PG}(5, q)$, there is a unique set of three (conjugate skew) transversal lines in $\text{PG}(5, q^3)$ that generate \mathcal{S} in this way.

We will use the linear representation of a finite translation plane \mathcal{P} of dimension at most three over its kernel, due independently to André [1954] and Bruck and Bose [1964; 1966]. Let Σ_∞ be a hyperplane of $\text{PG}(6, q)$ and let \mathcal{S} be a 2-spread of Σ_∞ . We use the phrase *a subspace of $\text{PG}(6, q) \setminus \Sigma_\infty$* to mean a subspace of $\text{PG}(6, q)$ that is not contained in Σ_∞ . Consider the following incidence structure: the *points* of $\mathcal{A}(\mathcal{S})$ are the points of $\text{PG}(6, q) \setminus \Sigma_\infty$; the *lines* of $\mathcal{A}(\mathcal{S})$ are the 3-spaces of $\text{PG}(6, q) \setminus \Sigma_\infty$ that contain an element of \mathcal{S} ; and *incidence* in $\mathcal{A}(\mathcal{S})$ is induced by incidence in $\text{PG}(6, q) \setminus \Sigma_\infty$. Then the incidence structure $\mathcal{A}(\mathcal{S})$ is an affine plane of order q^3 . We can complete $\mathcal{A}(\mathcal{S})$ to a projective plane $\mathcal{P}(\mathcal{S})$; the points on the line at infinity ℓ_∞ have a natural correspondence to the elements of the 2-spread \mathcal{S} . The projective plane $\mathcal{P}(\mathcal{S})$ is the Desarguesian plane $\text{PG}(2, q^3)$ if and only if \mathcal{S} is a regular 2-spread of $\Sigma_\infty \cong \text{PG}(5, q)$ (see [Bruck 1969]). For the remainder of this article, we use \mathcal{S} to denote a regular 2-spread of $\Sigma_\infty \cong \text{PG}(5, q)$.

We use the following notation. If T is a point of ℓ_∞ in $\text{PG}(2, q^3)$, we use $[T]$ to refer to the plane of \mathcal{S} corresponding to T . More generally, if X is a set of points of $\text{PG}(2, q^3)$, then we let $[X]$ denote the corresponding set in $\text{PG}(6, q)$. If P is an affine point of $\text{PG}(2, q^3)$, we generally simplify the notation and also use P to refer to the corresponding affine point in $\text{PG}(6, q)$, although in some cases, to avoid confusion, we use $[P]$.

When \mathcal{S} is a regular 2-spread, we can relate the coordinates of $\mathcal{P}(\mathcal{S}) \cong \text{PG}(2, q^3)$ and $\text{PG}(6, q)$ as follows. Let τ be a primitive element in \mathbb{F}_{q^3} with primitive polynomial $x^3 - t_2x^2 - t_1x - t_0$. Every element $\alpha \in \mathbb{F}_{q^3}$ can be uniquely written as $\alpha = a_0 + a_1\tau + a_2\tau^2$ with $a_0, a_1, a_2 \in \mathbb{F}_q$. Points in $\text{PG}(2, q^3)$ have homogeneous coordinates (x, y, z) with $x, y, z \in \mathbb{F}_{q^3}$, not all zero. Let the line at infinity ℓ_∞ have equation $z = 0$; so the affine points of $\text{PG}(2, q^3)$ have coordinates $(x, y, 1)$. Points in $\text{PG}(6, q)$ have homogeneous coordinates $(x_0, x_1, x_2, y_0, y_1, y_2, z)$ with $x_0, x_1, x_2, y_0, y_1, y_2, z \in \mathbb{F}_q$. Let Σ_∞ have equation $z = 0$. Let $P = (\alpha, \beta, 1)$ be a point of $\text{PG}(2, q^3)$. We can write $\alpha = a_0 + a_1\tau + a_2\tau^2$ and $\beta = b_0 + b_1\tau + b_2\tau^2$ with $a_0, a_1, a_2, b_0, b_1, b_2 \in \mathbb{F}_q$. We want to map the element α of \mathbb{F}_{q^3} to the vector (a_0, a_1, a_2) , and we use the following notation to do this:

$$[\alpha] = (a_0, a_1, a_2).$$

This gives some notation for the Bruck–Bose map, denoted ϵ , from an affine point $P = (\alpha, \beta, 1) \in \text{PG}(2, q^3) \setminus \ell_\infty$ to the corresponding affine point $[P] \in \text{PG}(6, q) \setminus \Sigma_\infty$, namely

$$\epsilon(\alpha, \beta, 1) = [(\alpha, \beta, 1)] = ([\alpha], [\beta], 1) = (a_0, a_1, a_2, b_0, b_1, b_2, 1).$$

More generally, if $z \in \mathbb{F}_q$, then $\epsilon(\alpha, \beta, z) = ([\alpha], [\beta], z) = (a_0, a_1, a_2, b_0, b_1, b_2, z)$.

Consider the case when $z = 0$, that is, a point on ℓ_∞ in $\text{PG}(2, q^3)$ has coordinates $L = (\alpha, \beta, 0)$ for some $\alpha, \beta \in \mathbb{F}_{q^3}$. In $\text{PG}(6, q)$, the point $\epsilon(\alpha, \beta, 0) = ([\alpha], [\beta], 0)$ is one point in the spread element $[L]$ corresponding to L . Moreover, the spread element $[L]$ consists of all the points $\{([\alpha x], [\beta x], 0) : x \in \mathbb{F}'_{q^3}\}$. Hence the regular 2-spread \mathcal{S} consists of the planes $\{[kx], [x], 0\} : x \in \mathbb{F}'_{q^3}\}$ for $k \in \mathbb{F}'_{q^3} \cup \{\infty\}$.

With this coordinatisation for the Bruck–Bose map, we can calculate the coordinates of the transversal lines of the regular 2-spread \mathcal{S} .

Lemma 2.1 [Barwick and Jackson 2012]. *Let $p_0 = t_1 + t_2\tau - \tau^2 = -\tau^q\tau^{q^2}$, $p_1 = t_2 - \tau = \tau^q + \tau^{q^2}$, $p_2 = -1$, and $A = (p_0, p_1, p_2)$. Then in the cubic extension $\text{PG}(6, q^3)$, one transversal line of the regular 2-spread \mathcal{S} contains the two points $A_1 = (p_0, p_1, p_2, 0, 0, 0, 0) = (A, [0], 0)$ and $A_2 = (0, 0, 0, p_0, p_1, p_2, 0) = ([0], A, 0)$.*

2B. Some useful homographies. In order to simplify the notation in some of the following coordinate-based proofs, we define some homographies which will be useful. We can represent an element $x = x_0 + x_1\tau + x_2\tau^2 \in \mathbb{F}_{q^3}$ as a point $[x] = (x_0, x_1, x_2)$ in $\text{PG}(2, q)$. For $k \in \mathbb{F}'_{q^3}$, consider the homography ζ_k in $\text{PGL}(3, q)$ with matrix M_k that maps $[x]$ to $[kx]$. Let $k \in \mathbb{F}'_{q^3}$ and write $k = k_0 + k_1\tau + k_2\tau^2$, then $M_k = k_0M_1 + k_1M_\tau + k_2M_{\tau^2}$, and hence

$$M_k A = k A \quad \text{and} \quad M_k A^{q^2} = k^{q^2} A^{q^2}, \quad (1)$$

where $A = (p_0, p_1, p_2)^t$ is defined in [Lemma 2.1](#). We use ζ_k to define the homography θ_k of $\text{PG}(5, q)$, $k \in \mathbb{F}_{q^3}$:

$$\theta_k : ([x], [y]) \rightarrow ([kx], [y]) = (M_k[x], [y]).$$

From the matrix M_τ , we construct three more homographies of $\text{PG}(2, q)$ with matrices U_0, U_1, U_2 that help with the notation in the proof of [Theorem 7.4](#). For $i = 0, 1, 2$, (with p_i as in [Lemma 2.1](#)), let

$$U_i = (p_0I + p_1M_\tau + p_2M_\tau^2)^{q^i} = \begin{pmatrix} p_0^{q^i} & \tau^{q^i} p_0^{q^i} & \tau^{2q^i} p_0^{q^i} \\ p_1^{q^i} & \tau^{q^i} p_1^{q^i} & \tau^{2q^i} p_1^{q^i} \\ p_2^{q^i} & \tau^{q^i} p_2^{q^i} & \tau^{2q^i} p_2^{q^i} \end{pmatrix}.$$

Then

$$U_i \begin{pmatrix} a_0 \\ a_1 \\ a_2 \end{pmatrix} = (a_0 + a_1\tau^{q^i} + a_2\tau^{2q^i}) \begin{pmatrix} p_0^{q^i} \\ p_1^{q^i} \\ p_2^{q^i} \end{pmatrix}, \quad a_0, a_1, a_2 \in \mathbb{F}_{q^3}.$$

Note that if $a_0, a_1, a_2 \in \mathbb{F}_q$, and $\alpha = a_0 + a_1\tau + a_2\tau^2$, then $[\alpha] = (a_0, a_1, a_2)^t$, and we write the matrix equation as $U_i[\alpha] = \alpha^{q^i} A^{q^i}$.

2C. Sublines in the Bruck–Bose representation. An *order- q -subplane* of $\text{PG}(2, q^3)$ is a subplane of $\text{PG}(2, q^3)$ of order q . Equivalently, it is an image of $\text{PG}(2, q)$ under $\text{PGL}(3, q^3)$. An *order- q -subline* of $\text{PG}(2, q^3)$ is a line of an order- q -subplane of $\text{PG}(2, q^3)$. An *order- q -subline* of $\text{PG}(1, q^3)$ is defined to be one of the images of $\text{PG}(1, q) = \{(a, 1) : a \in \mathbb{F}_q\} \cup \{(1, 0)\}$ under $\text{PGL}(2, q^3)$.

In [[Barwick and Jackson 2012; 2014](#)], we determine the representation of order- q -subplanes and order- q -sublines of $\text{PG}(2, q^3)$ in the Bruck–Bose representation in $\text{PG}(6, q)$, and we quote the results for order- q -sublines which are needed in this article. We first introduce some terminology to simplify the statements. Recall that \mathcal{S} is a regular 2-spread in the hyperplane at infinity Σ_∞ in $\text{PG}(6, q)$.

Definition 2.2. (i) An *\mathcal{S} -special conic* is a nondegenerate conic \mathcal{C} contained in a plane of \mathcal{S} , such that the extension of \mathcal{C} to $\text{PG}(6, q^3)$ meets the transversals of \mathcal{S} .

(ii) An *\mathcal{S} -special twisted cubic* is a twisted cubic \mathcal{N} in a 3-space of $\text{PG}(6, q) \setminus \Sigma_\infty$ about a plane of \mathcal{S} , such that the extension of \mathcal{N} to $\text{PG}(6, q^3)$ meets the transversals of \mathcal{S} .

Theorem 2.3 [[Barwick and Jackson 2012](#)]. *Let b be an order- q -subline of $\text{PG}(2, q^3)$.*

(i) *If $b \subset \ell_\infty$, then in $\text{PG}(6, q)$, b corresponds to a 2-regulus of \mathcal{S} . Conversely every 2-regulus of \mathcal{S} corresponds to an order- q -subline of ℓ_∞ .*

- (ii) If b meets ℓ_∞ in a point, then b corresponds to a line of $\text{PG}(6, q) \setminus \Sigma_\infty$. Conversely every line of $\text{PG}(6, q) \setminus \Sigma_\infty$ corresponds to an order- q -subline of $\text{PG}(2, q^3)$ tangent to ℓ_∞ .
- (iii) If b is disjoint from ℓ_∞ , then in $\text{PG}(6, q)$, b corresponds to an \mathcal{S} -special twisted cubic. Further, a twisted cubic \mathcal{N} of $\text{PG}(6, q)$ corresponds to an order- q -subline of $\text{PG}(2, q^3)$ if and only if \mathcal{N} is \mathcal{S} -special.

In [Barwick and Jackson 2012], we also determine the representation of secant and tangent order- q -subplanes of $\text{PG}(2, q^3)$ in $\text{PG}(6, q)$. The representation of an exterior order- q -subplane in $\text{PG}(6, q)$ is more complex to describe. One of the motivations of this work is to investigate this representation in more detail. Some aspects of the representation are discussed in more detail in Section 4.

2D. Properties of exterior splashes. We need some group theoretic results about order- q -subplanes and exterior splashes; the first appears in [Barwick and Jackson 2016].

Theorem 2.4. *Let $G = \text{PGL}(3, q^3)$ be the collineation group acting on $\text{PG}(2, q^3)$. The subgroup G_ℓ fixing a line ℓ is transitive on the order- q -subplanes that are exterior to ℓ , and is transitive on the exterior splashes of ℓ .*

This theorem can be proved by generalising the arguments in [Barwick and Jackson 2015]. In particular, it involves looking at two important subgroups of G . The first subgroup fixes an order- q -subplane, and the following property will be very useful.

Theorem 2.5. *The group $K = \text{PGL}(3, q^3)_\pi$ acting on $\text{PG}(2, q^3)$ and fixing an order- q -subplane π is transitive on the points of π .*

The second important subgroup is $I = G_{\pi, \ell}$ which fixes an order- q -subplane π , and a line ℓ exterior to π . By [Barwick and Jackson 2016], I fixes exactly three lines: ℓ , and its conjugates m, n with respect to π ; and I fixes exactly three points: $E_1 = \ell \cap m$, $E_2 = \ell \cap n$, $E_3 = m \cap n$, which are conjugate with respect to π . Further I identifies two fixed points $E_1 = \ell \cap m$, $E_2 = \ell \cap n$ on ℓ which are called the *carriers* of the exterior splash \mathbb{S} of π . This is consistent with the definition of carriers of a circle geometry $\text{CG}(3, q)$; see [Barwick and Jackson 2016]. In [Lunardon et al. 2014], scattered linear sets of pseudoregulus type are considered, and they use the term “transversal points”. The fixed points and fixed lines of I are used to define an important class of conics in an order- q -subplane π with respect to an exterior line ℓ . A conic of π whose extension to $\text{PG}(2, q^3)$ contains the three points E_1, E_2, E_3 is called a (π, ℓ) -*carrier conic* of π . A dual conic of π whose extension to $\text{PG}(2, q^3)$ contains the three lines ℓ, m, n is called a (π, ℓ) -*carrier-dual conic*. Note that carrier-conics/dual conics were called special-conics/dual

conics in [Barwick and Jackson 2016]; we change the name here so that the term “special” is reserved for objects in $\text{PG}(6, q)$.

3. Coordinatising an exterior order- q -subplane

Recall from Theorem 2.4 that the group of homographies of $\text{PG}(2, q^3)$ is transitive on pairs (π, ℓ) where π is an order- q -subplane exterior to the line ℓ . So if we want to use coordinates to prove a result about exterior order- q -subplanes, we can without loss of generality prove it for a particular exterior order- q -subplane. In this section we calculate the coordinates for an exterior order- q -subplane \mathcal{B} of $\text{PG}(2, q^3)$ whose exterior splash has a simple form. Set

$$K = \begin{pmatrix} -\tau & 1 & 0 \\ -\tau^q & 1 & 0 \\ \tau\tau^q & -\tau - \tau^q & 1 \end{pmatrix}, \quad K' = \begin{pmatrix} -1 & 1 & 0 \\ -\tau^q & \tau & 0 \\ -\tau^{2q} & \tau^2 & \tau - \tau^q \end{pmatrix}. \quad (2)$$

Let σ be the homography of $\text{PG}(2, q^3)$ with matrix K . Note that as KK' is a \mathbb{F}_{q^3} -multiple of the identity matrix, it follows that K' is a matrix for the inverse homography σ^{-1} . Thus, if we write the points X of $\text{PG}(2, q^3)$ as column vectors, and the lines ℓ of $\text{PG}(2, q^3)$ as row vectors, then $\sigma(X) = KX$ and $\sigma(\ell) = \ell K'$. The order- q -subplane $\pi_0 = \text{PG}(2, q)$ is secant to ℓ_∞ . We show that the subplane $\sigma(\pi_0)$ is exterior to ℓ_∞ and has the desired simple form as exterior splash.

Theorem 3.1. *In $\text{PG}(2, q^3)$, let $\pi_0 = \text{PG}(2, q)$, let σ be the homography with matrix K given in (2), and let $\mathcal{B} = \sigma(\pi_0)$. Then \mathcal{B} is an order- q -subplane exterior to ℓ_∞ with exterior splash $\mathbb{S} = \{(k, 1, 0) : k \in \mathbb{F}_{q^3}, k^{q^2+q+1} = 1\}$ and carriers $E_1 = (1, 0, 0)$ and $E_2 = (0, 1, 0)$.*

Proof. Note that σ maps $\pi_0 = \text{PG}(2, q)$ to \mathcal{B} and the line $\ell = [-\tau\tau^q, \tau + \tau^q, -1]$ to $\ell_\infty = [0, 0, 1]$. By [Barwick and Jackson 2016], π_0 is exterior to ℓ and has carriers $E = (1, \tau, \tau^2)$ and $E^q = (1, \tau^q, \tau^{2q})$ on ℓ . Hence \mathcal{B} is exterior to ℓ_∞ and has carriers $\sigma(E) = (0, 1, 0)$ and $\sigma(E^q) = (1, 0, 0)$ on ℓ_∞ . By considering the action of σ on the lines $[l, m, n]$ ($l, m, n \in \mathbb{F}_q$, not all zero) of π_0 , we calculate the lines of \mathcal{B} are $\ell_{l,m,n} = [-l - \tau^q m - \tau^{2q} n, l + \tau m + \tau^2 n, n(\tau - \tau^q)]$, with $l, m, n \in \mathbb{F}_q$, not all zero. The exterior splash of \mathcal{B} consists of the points $Q_{l,m,n} = \ell_{l,m,n} \cap \ell_\infty = (l + \tau m + \tau^2 n, (l + \tau m + \tau^2 n)^q, 0)$. Writing $y = l + \tau m + \tau^2 n$, gives $Q_{l,m,n} \equiv (y, y^q, 0) \equiv (y^{1-q}, 1, 0)$ and writing $y = \tau^{-j}$ for some $j \in \{0, \dots, q^3 - 2\}$ yields $Q_{l,m,n} \equiv (\tau^{j(q-1)}, 1, 0)$. Note that if we write $j = n(q^2 + q + 1) + i$ where $0 \leq i < q^2 + q + 1$, then $\tau^{j(q-1)} = \tau^{i(q-1)}$. So we may assume that $Q_{l,m,n} = (\tau^{i(q-1)}, 1, 0)$ with $0 \leq i < q^2 + q + 1$. It is useful to observe that

$$\mathbb{S} = \{(k, 1, 0) : k \in \mathbb{F}_{q^3}, k^{q^2+q+1} = 1\} \equiv \{(\tau^{(q-1)i}, 1, 0) : 0 \leq i < q^2 + q + 1\}$$

as the solutions to $k^{q^2+q+1} = 1$ are $\tau^{i(q-1)}$, $0 \leq i < q^2 + q + 1$. \square

4. The structure of the subplane in $\text{PG}(6, q)$

If π is an exterior order- q -subplane of $\text{PG}(2, q^3)$, then in the Bruck–Bose representation in $\text{PG}(6, q)$, π corresponds to a set of $q^2 + q + 1$ affine points denoted $[\pi]$. It is difficult to characterise the structure of $[\pi]$. We note that as π contains $q^2 + q + 1$ order- q -sublines that are exterior to ℓ_∞ , then by [Theorem 2.3](#), $[\pi]$ contains $q^2 + q + 1$ \mathcal{S} -special twisted cubics, each lying in a 3-space through a distinct plane of the exterior splash of π . In this section we aim to determine more about the structure of $[\pi]$.

4A. The intersection of nine quadrics. We show that the structure $[\pi]$ of $\text{PG}(6, q)$ corresponding to an exterior order- q -subplane π of $\text{PG}(2, q^3)$ is the intersection of nine quadrics in $\text{PG}(6, q)$. This is analogous to [\[Barwick and Jackson 2015, Theorem 9.2\]](#) which shows that a *tangent* order- q -subplane of $\text{PG}(2, q^3)$ corresponds to a structure in $\text{PG}(6, q)$ that is the intersection of nine quadrics.

Theorem 4.1. *Let π be an exterior order- q -subplane in $\text{PG}(2, q^3)$. The corresponding set $[\pi]$ in $\text{PG}(6, q)$ is the intersection of nine quadrics.*

Proof. By [Theorem 2.4](#), we can without loss of generality prove this for the order- q -subplane \mathcal{B} coordinatised in [Section 3](#). We use the homographies σ, σ^{-1} with matrices K, K' respectively, given in [\(2\)](#). A point $P = (x, y, 1) \in \text{PG}(2, q^3)$ belongs to \mathcal{B} if its preimage $K'P = (-x + y, -\tau^q x + \tau y, -\tau^{2q} x + \tau^2 y + (\tau - \tau^q))$ belongs to $\pi_0 = \text{PG}(2, q)$. Suppose firstly that $-x + y \neq 0$, then

$$K'P \equiv \left(1, \frac{-\tau^q x + \tau y}{-x + y}, \frac{-\tau^{2q} x + \tau^2 y + (\tau - \tau^q)}{-x + y} \right).$$

This belongs to $\pi_0 = \text{PG}(2, q)$ if and only if the second and third coordinates belong to \mathbb{F}_q , that is,

$$\left(\frac{-\tau^q x + \tau y}{-x + y} \right)^q = \frac{-\tau^q x + \tau y}{-x + y}, \quad (3)$$

$$\left(\frac{-\tau^{2q} x + \tau^2 y + (\tau - \tau^q)}{-x + y} \right)^q = \frac{-\tau^{2q} x + \tau^2 y + (\tau - \tau^q)}{-x + y}. \quad (4)$$

Writing $x = x_0 + x_1\tau + x_2\tau^2$ and $y = y_0 + y_1\tau + y_2\tau^2$, where $x_i, y_i \in \mathbb{F}_q$ and $i = 1, 2, 3$, then equating powers of $1, \tau, \tau^2$, yields three quadratic equations from each condition, a total of six, each of which represents a quadric in $\text{PG}(6, q)$.

Secondly, suppose $-\tau^q x + \tau y \neq 0$, then

$$K'P \equiv \left(\frac{-x + y}{-\tau^q x + \tau y}, 1, \frac{-\tau^{2q} x + \tau^2 y + (\tau - \tau^q)}{-\tau^q x + \tau y} \right).$$

As before, this lies in π_0 if and only if

$$\left(\frac{-x + y}{-\tau^q x + \tau y} \right)^q = \frac{-x + y}{-\tau^q x + \tau y}, \quad (5)$$

$$\left(\frac{-\tau^{2q} x + \tau^2 y + (\tau - \tau^q)}{-\tau^q x + \tau y} \right)^q = \frac{-\tau^{2q} x + \tau^2 y + (\tau - \tau^q)}{-\tau^q x + \tau y}, \quad (6)$$

leading to a further six quadrics in $\text{PG}(6, q)$. The equations (3) and (5) give the same triple of quadrics. Hence the point P lies in \mathcal{B} if and only if the point $[P]$ lies on a total of nine quadrics in $\text{PG}(6, q)$. Finally, note that if both $-x + y = 0$ and $-\tau^q x + \tau y = 0$, then $x = y = 0$ and the point P has coordinates $(0, 0, 1)$. This satisfies all the quadratic equations from (3), (4), (6), and so in $\text{PG}(6, q)$, $[P]$ lies on each of the nine quadrics. \square

4B. Tangent planes at points of an exterior subplane. We now consider a point P lying in an exterior order- q -subplane π of $\text{PG}(2, q^3)$. In the Bruck–Bose representation in $\text{PG}(6, q)$, P corresponds to an affine point which we also denote by P . We show that in $\text{PG}(6, q)$, there is a unique *tangent plane* \mathcal{T}_P at P to the structure $[\pi]$. We show that there are two equivalent ways to define this tangent plane. Recall from [Theorem 2.3](#) that the order- q -sublines of π correspond to twisted cubics in $\text{PG}(6, q)$. [Theorem 4.2](#) shows that we can define \mathcal{T}_P by looking at the tangent lines at P to these twisted cubics. Then [Theorem 4.3](#) shows that we can define \mathcal{T}_P by looking at the tangent space of P with respect to the nine quadrics defined by $[\pi]$.

Theorem 4.2. *Let π be an exterior order- q -subplane of $\text{PG}(2, q^3)$, and let P be a point of π . Label the lines of π through P by ℓ_0, \dots, ℓ_q . In $\text{PG}(6, q)$, ℓ_i corresponds to a twisted cubic $[\ell_i]$. Let m_i be the unique tangent line to $[\ell_i]$ through P . Then the lines m_0, \dots, m_q lie in a plane \mathcal{T}_P , called the tangent plane of $[\pi]$ at P .*

Proof. By [Theorems 2.4](#) and [2.5](#), we can without loss of generality prove this for the order- q -subplane \mathcal{B} coordinatised in [Section 3](#), and the point $P = (0, 0, 1)$ of \mathcal{B} . First consider the order- q -subplane $\pi_0 = \text{PG}(2, q)$. The point $P = (0, 0, 1)$ lies in π_0 , and the lines of π_0 through P have coordinates $\ell'_m = [m, 1, 0]$, $m \in \mathbb{F}_q \cup \{\infty\}$. Points on the line ℓ'_m distinct from P have coordinates $P'_x = (1, -m, x)$ for $x \in \mathbb{F}_q$. We map the plane π_0 to \mathcal{B} using the homography σ with matrix K given in [\(2\)](#). As $\sigma(P) = P$, the lines of \mathcal{B} through P are $\ell_m = \sigma(\ell'_m)$, $m \in \mathbb{F}_q \cup \{\infty\}$. Points on the line ℓ_m distinct from P have coordinates

$$P_x = \sigma(P'_x) = (-\tau - m, -\tau^q - m, \tau\tau^q + (\tau + \tau^q)m + x),$$

for $x \in \mathbb{F}_q$.

To convert this to a coordinate in $\text{PG}(6, q)$, we need to multiply by an element of \mathbb{F}_{q^3} so that the last coordinate lies in \mathbb{F}_q . Let $F(x) = \tau\tau^q + (\tau + \tau^q)m + x$ (the third coordinate in P_x). As $F(x) \in \mathbb{F}_{q^3}$, we have $F(x)^{q^2+q+1} \in \mathbb{F}_q$. So in $\text{PG}(6, q)$, we

have the point $P_x = ([-(\tau + m)F(x)^{q^2+q}], [-(\tau^q + m)F(x)^{q^2+q}], F(x)^{q^2+q+1})$.

By [Theorem 2.3](#), the line ℓ_m of $\text{PG}(2, q^3)$ corresponds to a twisted cubic $[\ell_m] = \{P_x : x \in \mathbb{F}_q\} \cup \{P\}$ of $\text{PG}(6, q)$. Consider the unique tangent to $[\ell_m]$ through P , and let I_m be the intersection of this tangent with Σ_∞ . We will show that the points I_m , $m \in \mathbb{F}_q \cup \{\infty\}$, form a line. To calculate the coordinates of I_m , we let $Q_x = P P_x \cap \Sigma_\infty$. To calculate $I_m = Q_\infty$, we use the homogeneous coordinate technique of dividing by the largest power of x , and then substituting $x = \infty$, that is, replacing $1/x$ by 0. We use the notation $\lim_{x \rightarrow \infty}$ to describe this technique.

$$\begin{aligned} I_m &= \lim_{x \rightarrow \infty} P P_x \cap \Sigma_\infty = \lim_{x \rightarrow \infty} ([-(\tau + m)F(x)^{q^2+q}], [-(\tau^q + m)F(x)^{q^2+q}], 0) \\ &= ([-(\tau + m)], -[\tau^q + m], 0). \end{aligned}$$

Hence the points I_m , $m \in \mathbb{F}_q \cup \{\infty\}$, form a line $\ell = \langle ([1], [1], 0), ([\tau], [\tau^q], 0) \rangle$ in Σ_∞ . Hence the tangent lines m_0, \dots, m_q to the twisted cubics of $[\pi]$ through P form a plane $\mathcal{T}_P = \langle \ell, P \rangle$ through P , as required. \square

Theorem 4.3. *Let π be an exterior order- q -subplane of $\text{PG}(2, q^3)$, and let P be a point of π . In $\text{PG}(6, q)$, consider the intersection of the tangent spaces at P of the nine quadrics corresponding to $[\pi]$. Then this intersection is equal to the tangent plane \mathcal{T}_P of $[\pi]$ at P as defined in [Theorem 4.2](#).*

Proof. By [Theorems 2.4](#) and [2.5](#), we can without loss of generality prove this for the order- q -subplane \mathcal{B} coordinatised in [Section 3](#), and the point $P = (0, 0, 1)$ of \mathcal{B} . In $\text{PG}(6, q)$, consider the nine quadrics corresponding to $[\mathcal{B}]$ which are given in equations [\(4\)](#), [\(5\)](#) and [\(6\)](#). We want to find the set of lines through P that meet each of these nine quadrics twice at P . Every line ℓ of $\text{PG}(6, q)$ through P has the form $\ell = RP$ for some point $R = ([u], [v], 0) \in \Sigma_\infty$, $u, v \in \mathbb{F}_{q^3}$. So the points of ℓ are of the form $P_s = P + sR = ([su], [sv], 1)$ where $s \in \mathbb{F}_q$. Substituting the point P_s into the quadrics of [\(4\)](#) gives

$$(-\tau^{2q}su + \tau^{2q}sv + (\tau - \tau^q))^q(-su + sv) = (-\tau^{2q}su + \tau^2sv + (\tau - \tau^q))(-su + sv)^q.$$

This expression is a polynomial of degree two in s . The line $\ell = PR$ is tangent to the three quadrics of [\(4\)](#) if this expression has a repeated root $s = 0$, that is, if the coefficient of s is equal to zero. That is,

$$(\tau - \tau^q)^q(-u + v) = (\tau - \tau^q)(-u + v)^q,$$

and so $k = (-u + v)/(\tau - \tau^q)$ is in \mathbb{F}_q . Rearranging gives $v = k(\tau - \tau^q) + u$. Substituting the point P_s into the quadrics of [\(5\)](#) gives no constraints. Substituting the point P_s into the quadrics of [\(6\)](#) and simplifying gives that the constraint $m = (-\tau^q u + \tau v)/(\tau - \tau^q)$ lies in \mathbb{F}_q , and so $v = (m(\tau - \tau^q) + \tau^q u)/\tau$. Equating this with the expression for v obtained from [\(4\)](#) gives $u = m - k\tau$, and so $v = m - k\tau^q$.

Hence the line $\ell = PR$ is tangent to all nine quadrics when R has form

$$R = ([u], [v], 0) = ([m - k\tau], [m - k\tau^q], 0) = m([1], [1], 0) - k([\tau], [\tau^q], 0).$$

Thus the tangent space to $[\mathcal{B}]$ at P is the plane through P and the line

$$\ell = \langle ([1], [1], 0), ([\tau], [\tau^q], 0) \rangle$$

of Σ_∞ . This is the same as the tangent plane \mathcal{T}_P to $[\mathcal{B}]$ at P calculated in the proof of [Theorem 4.2](#). \square

5. Coordinatising the exterior splash and its covers

Let \mathbb{S} be an exterior splash of $\text{PG}(1, q^3)$. In the Bruck–Bose representation, \mathbb{S} corresponds to a set of $q^2 + q + 1$ planes of the regular 2-spread \mathcal{S} in $\Sigma_\infty \cong \text{PG}(5, q)$. To simplify the notation, we use the same symbol \mathbb{S} to denote both the points of the exterior splash on ℓ_∞ , and the planes of the exterior splash contained in \mathcal{S} . In [\[Barwick and Jackson 2016\]](#), we show that an exterior splash is projectively equivalent to a cover of the circle geometry $\text{CG}(3, q)$. Hence by Bruck [\[1973\]](#), there are two *switching sets* \mathbb{X}, \mathbb{Y} for \mathbb{S} . That is, \mathbb{X} and \mathbb{Y} consist of $q^2 + q + 1$ planes each, such that the planes of the three sets \mathbb{S}, \mathbb{X} and \mathbb{Y} each cover the same set of points. Further, planes from different sets meet in unique points, and planes in the same set are disjoint. The three sets $\mathbb{S}, \mathbb{X}, \mathbb{Y}$ are called *hyper-reguli* in [\[Culbert and Ebert 2005; Ostrom 1993\]](#). In this article, we call the families \mathbb{X} and \mathbb{Y} *covers* of the exterior splash \mathbb{S} .

In this section we take the order- q -subplane \mathcal{B} coordinatised in [Section 3](#), with exterior splash \mathbb{S} , and use [\[Ostrom 1993\]](#) to calculate the coordinates of the two covers of \mathbb{S} . We will characterise the two covers in terms of the subplane \mathcal{B} .

We call one cover of \mathbb{S} the *tangent cover with respect to* \mathcal{B} , and denote it by $\mathbb{T}_{\mathcal{B}}$, or if there is only one subplane under consideration, we shorten this to \mathbb{T} . The nomenclature for tangent covers comes from [Theorem 5.3](#) which shows that the tangent planes \mathcal{T}_P of $[\mathcal{B}]$ meet Σ_∞ in lines that lie in distinct planes of the cover \mathbb{T} .

We call the other cover of \mathbb{S} the *conic cover with respect to* \mathcal{B} , and denote it by $\mathbb{C}_{\mathcal{B}}$, or \mathbb{C} . The nomenclature for the conic cover comes from [\[Barwick and Jackson 2017\]](#) which shows that the planes in the cover \mathbb{C} are related to the $(\mathcal{B}, \ell_\infty)$ -carrier conics of \mathcal{B} .

A certain type of embedding is looked at in [\[Lavrauw et al. 2015\]](#). Specialising their results to $\text{PG}(5, q)$, their embedding $\mathcal{Q}_{2,q}$ is equivalent to the set $\mathbb{S} \cup \mathbb{C} \cup \mathbb{T}$. They determine the collineation group stabilising $\mathcal{Q}_{2,q}$. In particular they demonstrate: a collineation of $\text{PG}(5, q)$ that fixes $\mathcal{Q}_{2,q}$ and permutes the families $\mathbb{S}, \mathbb{C}, \mathbb{T}$; and a collineation fixing $\mathcal{Q}_{2,q}$ that permutes the planes in each family. Further, [\[Lavrauw et al. 2015\]](#) determines the equation of $\mathcal{Q}_{2,q}$. In [Lemma 5.1](#) we describe

the homogeneous coordinates for the planes in \mathbb{S} , \mathbb{C} , \mathbb{T} in the format we will work with, and in [Lemma 5.2](#) we calculate the matrix of a homography that fixes the planes in \mathbb{S} , permutes the planes of \mathbb{T} , and permutes the planes of \mathbb{C} (this is the map $\varphi_{0,0}(\tau, \tau)$ of [\[Lavrauw et al. 2015\]](#)).

Lemma 5.1. *Let \mathbb{S} be the exterior splash of the exterior order- q -subplane \mathcal{B} coordinatised in [Section 3](#). Let $\mathcal{K} = \{k = \tau^{i(q-1)} : 0 \leq i < q^2 + q + 1\}$. In $\text{PG}(6, q)$, \mathbb{S} and its two covers \mathbb{T} , \mathbb{C} have planes given by*

$$\mathbb{S} = \{[S_k] = \{([kx], [x], 0) : x \in \mathbb{F}'_{q^3}\} : k \in \mathcal{K}\},$$

$$\mathbb{T} = \{[T_k] = \{([kx], [x^q], 0) : x \in \mathbb{F}'_{q^3}\} : k \in \mathcal{K}\},$$

$$\mathbb{C} = \{[C_k] = \{([kx], [x^{q^2}], 0) : x \in \mathbb{F}'_{q^3}\} : k \in \mathcal{K}\}.$$

Proof. The points of ℓ_∞ in $\text{PG}(2, q^3)$ have coordinates $S_k = (k, 1, 0)$ for $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. Hence in the Bruck–Bose representation of ℓ_∞ in $\Sigma_\infty \cong \text{PG}(5, q)$, planes of the regular 2-spread \mathcal{S} are given by $[S_k] = \{([kx], [x]) : x \in \mathbb{F}'_{q^3}\}$, for $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. Consider the homography β (of order 3) of $\Sigma_\infty \cong \text{PG}(5, q)$ defined by

$$\beta : ([x], [y]) \rightarrow ([x], [y^q]). \quad (7)$$

We consider the action of β on the planes of $[S_k]$. For each $k \in \mathbb{F}_{q^3} \cup \{\infty\}$, define the planes $[T_k]$, $[C_k]$ by $\beta([S_k]) = [T_k]$ and $\beta([T_k]) = [C_k]$. That is, $[T_k] = \{([kx], [x^q]) : x \in \mathbb{F}'_{q^3}\}$, and $[C_k] = \{([kx], [x^{q^2}]) : x \in \mathbb{F}'_{q^3}\}$.

We now consider the exterior order- q -subplane \mathcal{B} coordinatised in [Section 3](#) which by [Theorem 3.1](#) has exterior splash $\mathbb{S} = \{S_k = (k, 1, 0) : k \in \mathcal{K}\} \subset \ell_\infty$, and carriers $S_\infty = (1, 0, 0)$, $S_0 = (0, 1, 0)$. Note that in $\text{PG}(5, q)$, the carriers of \mathcal{B} lie in each of the three sets of planes, as $[S_0] = [T_0] = [C_0]$ and $[S_\infty] = [T_\infty] = [C_\infty]$. In $\text{PG}(5, q)$, we have $\mathbb{S} = \{[S_k] : k \in \mathcal{K}\}$. Let $\mathbb{T} = \{[T_k] : k \in \mathcal{K}\}$ and $\mathbb{C} = \{[C_k] : k \in \mathcal{K}\}$, then $\beta : \mathbb{S} \mapsto \mathbb{T} \mapsto \mathbb{C}$. By [\[Ostrom 1993\]](#), the sets \mathbb{S} , \mathbb{T} , \mathbb{C} cover the same set of points. Moreover, planes in the same set are disjoint, and planes from different sets meet in one point. That is, \mathbb{T} and \mathbb{C} are the two covers of \mathbb{S} . \square

The next lemma calculates the action of a useful homography of $\text{PG}(6, q)$ (this is the map $\varphi_{0,0}(\tau, \tau)$ of [\[Lavrauw et al. 2015\]](#)). Recall that τ is a zero of the primitive polynomial $x^3 - t_2x^2 - t_1x - t_0$.

Lemma 5.2. *Let \mathbb{S} be the exterior splash of the exterior order- q -subplane \mathcal{B} coordinatised in [Section 3](#) with covers \mathbb{C} and \mathbb{T} coordinatised in [Lemma 5.1](#). Consider the homography $\Theta \in \text{PGL}(7, q)$ with 7×7 matrix*

$$\begin{pmatrix} M & 0 & 0 \\ 0 & M & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \text{where } M = \begin{pmatrix} 0 & 0 & t_0 \\ 1 & 0 & t_1 \\ 0 & 1 & t_2 \end{pmatrix}.$$

Then in $\text{PG}(6, q)$, Θ fixes each plane of the regular 2-spread \mathcal{S} , maps the cover plane $[C_k] \in \mathbb{C}$ to $[C_{\tau^{1-q}k}] \in \mathbb{C}$, and the cover plane $[T_k] \in \mathbb{T}$ to $[T_{\tau^{1-q^2}k}] \in \mathbb{T}$, $k \in \mathcal{K}$.

Proof. It is straightforward to show that Θ fixes the planes of the regular 2-spread \mathcal{S} (so it also fixes the planes of the exterior splash \mathcal{S}). In fact $\langle \Theta \rangle$ acts regularly on the set of points, and on the set of lines, of each spread element. Note that M is the matrix M_τ defined in [Section 2B](#), and so $M[x] = [\tau x]$. Consider the action of Θ on a point of the cover plane $[C_k] \in \mathbb{C}$ coordinatised in [Lemma 5.1](#). We have

$$([kx], [x^{q^2}], 0)^\Theta = ([\tau kx], [\tau x^{q^2}], 0) \equiv ([\tau^{1-q}k(\tau^q x)], [(\tau^q x)^{q^2}], 0)$$

which lies in the cover plane $[C_{\tau^{1-q}k}]$ of \mathbb{C} . Similarly a point $([kx], [x^q], 0)$ in the cover plane $[T_k] \in \mathbb{T}$ maps under Θ to the point $([\tau^{1-q^2}k(\tau^{q^2} x)], [(\tau^{q^2} x)^q], 0)$ which lies in the cover plane $[T_{\tau^{1-q^2}k}]$ of \mathbb{T} . \square

Theorem 5.3. *Let P be a point of an exterior order- q -subplane π . In $\text{PG}(6, q)$, the tangent plane \mathcal{T}_P at P to $[\pi]$ meets Σ_∞ in a line that lies in a plane of the tangent cover \mathbb{T} of $[\pi]$. Moreover, distinct points of π correspond to distinct cover planes of \mathbb{T} .*

Proof. By [Theorems 2.4](#) and [2.5](#), we can without loss of generality prove this result for the order- q -subplane \mathcal{B} coordinatised in [Section 3](#) and the point $P = (0, 0, 1) \in \mathcal{B}$. In $\text{PG}(6, q)$, let \mathcal{T}_P be the tangent plane at P . The line $\ell = \mathcal{T}_P \cap \Sigma_\infty$ was calculated in the proof of [Theorem 4.2](#) to be

$$\ell = \{a([1], [1], 0) + b([\tau], [\tau^q], 0) : a, b \in \mathbb{F}_q\}.$$

The points of ℓ all lie in the plane $[T_1] = \{[x], [x^q], 0\} \mid x \in \mathbb{F}'_3\}$, which by [Lemma 5.1](#) is a plane of the tangent cover \mathbb{T} of \mathcal{B} . The collineation of [Lemma 5.2](#) is transitive on the cover planes of \mathbb{T} , hence each cover plane contains a line of a distinct tangent plane. Hence there is a one-to-one correspondence between points of π and cover planes of \mathbb{T} . \square

6. Transversal lines of covers

Recall that a regular 2-spread in $\text{PG}(5, q)$ has three (conjugate skew) transversals in $\text{PG}(5, q^3)$ which meet each (extended) plane of \mathcal{S} . In this section we consider an exterior splash $\mathcal{S} \subset \mathcal{S}$, and show in [Lemma 6.1](#) that the transversals of the 2-spread \mathcal{S} are the only lines of $\text{PG}(5, q^3)$ that meet every extended plane of \mathcal{S} . We then consider the two sets of cover planes \mathbb{T} and \mathbb{C} . [Corollary 6.2](#) shows that each can be uniquely extended to regular 2-spread, and we calculate the coordinates of the corresponding transversal lines in [Theorem 6.3](#). [Theorem 6.5](#) shows that the nine transversals of \mathcal{S} , \mathbb{C} and \mathbb{T} can be used to characterise the carriers of the exterior splash \mathcal{S} . [Theorem 6.6](#), looks at the transversal lines in the situation when ℓ_∞ is partitioned into exterior splashes with common carriers.

6A. The exterior splash and its covers have unique transversals. If \mathcal{X} is a set in $\text{PG}(6, q)$ (such as a line, a plane, or a conic), then we denote its natural extension to $\text{PG}(6, q^3)$ by \mathcal{X}^* . Let \mathcal{S} be the regular 2-spread in Σ_∞ of the Bruck–Bose representation in $\text{PG}(6, q)$. If we extend the planes of \mathcal{S} to $\text{PG}(6, q^3)$, yielding \mathcal{S}^* , then there are exactly three transversal lines to \mathcal{S}^* , that is, three lines that meet every plane of \mathcal{S}^* . These three lines are conjugate and skew. We now consider an exterior splash $\mathbb{S} \subset \mathcal{S}$ and extend the planes of \mathbb{S} to $\text{PG}(6, q^3)$, yielding \mathbb{S}^* . We show that there are exactly three lines of $\text{PG}(6, q^3)$ that meet every plane of \mathbb{S}^* , namely the three transversals of \mathcal{S} .

Lemma 6.1. *Let \mathcal{S} be a regular 2-spread in $\text{PG}(5, q)$, and let $\mathbb{S} \subset \mathcal{S}$ be an exterior splash. In the cubic extension $\text{PG}(5, q^3)$, there are exactly three transversals to \mathbb{S} , namely the three transversals of \mathcal{S} . Hence \mathbb{S} lies in a unique regular 2-spread, namely \mathcal{S} .*

Proof. The three conjugate transversal lines of the regular 2-spread \mathcal{S} , denoted $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}$, are also transversals of \mathbb{S} . Suppose there is a fourth transversal line ℓ of \mathbb{S} . Then the four lines $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}, \ell$ are pairwise skew. Further, these four lines are ruling lines of a unique 2-regulus \mathcal{R} of $\Sigma_\infty^* \cong \text{PG}(5, q^3)$, which contains the set of extended planes \mathbb{S}^* . Now consider two planes $[L], [M] \in \mathbb{S}$; the corresponding points L, M of ℓ_∞ in $\text{PG}(2, q^3)$ lie in two order- q -sublines contained in \mathbb{S} by [Lavrauw and Van de Voorde 2010, Corollary 15]. Hence by Theorem 2.3, $[L], [M]$ lie in two 2-reguli $\mathcal{R}_1, \mathcal{R}_2$ which are contained in \mathbb{S} . Let P be a point in $[L]$, then there are unique lines m_1, m_2 through P that are ruling lines of $\mathcal{R}_1, \mathcal{R}_2$ respectively. Now $\mathcal{R}_1, \mathcal{R}_2$ lie in \mathbb{S} , and so lie in \mathcal{R} , so the extended lines $m_i^*, i = 1, 2$, are two ruling lines of \mathcal{R} that meet in a point P , a contradiction. Hence the line ℓ cannot exist. That is, there are only three transversal lines to \mathbb{S} , and these are necessarily the transversals of \mathcal{S} . \square

As $\mathbb{S}, \mathbb{C}, \mathbb{T}$ are projectively equivalent by [Lavrauw et al. 2015, Theorem 16], an analogous result holds for the two covers of \mathbb{S} .

Corollary 6.2. *In $\text{PG}(5, q)$, let \mathbb{S} be an exterior splash with covers \mathbb{T} and \mathbb{C} . Then in the cubic extension $\text{PG}(5, q^3)$,*

- (i) *the cover \mathbb{T} has exactly three transversal lines in $\text{PG}(5, q^3) \setminus \text{PG}(5, q)$, denoted $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$, and so \mathbb{T} lies in a unique regular 2-spread,*
- (ii) *the cover \mathbb{C} has exactly three transversal lines in $\text{PG}(5, q^3) \setminus \text{PG}(5, q)$, denoted $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$, and so \mathbb{C} lies in a unique regular 2-spread.*

Later we will need the coordinates of the point of intersection of the transversal lines with the corresponding cover planes, and we calculate these next.

Theorem 6.3. *Let \mathcal{B} be the order- q -subplane coordinatised in Section 3 with exterior splash \mathbb{S} and covers \mathbb{C}, \mathbb{T} . Let $p_0 = t_1 + t_2\tau - \tau^2 = -\tau^q\tau^{q^2}$, $p_1 = t_2 - \tau =$*

$\tau^q + \tau^{q^2}$, $p_2 = -1$, and $\eta = p_0 + p_1\tau + p_2\tau^2$. Let $A_1 = (p_0, p_1, p_2, 0, 0, 0, 0)$, $A_2 = (0, 0, 0, p_0, p_1, p_2, 0)$. Then in $\text{PG}(6, q^3)$,

- (i) one transversal line of \mathbb{S} is $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$, and $g_{\mathbb{S}} \cap [S_k]^* = kA_1 + A_2$,
- (ii) one transversal line of \mathbb{T} is $g_{\mathbb{T}} = \langle A_1, A_2^{q^2} \rangle$, and $g_{\mathbb{T}} \cap [T_k]^* = kA_1 + \eta^{1-q^2} A_2^{q^2}$,
- (iii) one transversal line of \mathbb{C} is $g_{\mathbb{C}} = \langle A_1, A_2^q \rangle$, and $g_{\mathbb{C}} \cap [C_k]^* = kA_1 + \eta^{1-q} A_2^q$.

Proof. We use the coordinatisation in $\text{PG}(5, q)$ of the exterior splash \mathbb{S} of \mathcal{B} and the two covers \mathbb{T}, \mathbb{C} given in Lemma 5.1. Lemma 2.1 shows that $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$ is a transversal line for the regular 2-spread \mathcal{S} , where $A_1 = (p_0, p_1, p_2, 0, 0, 0) = (A, [0])$ and $A_2 = (0, 0, 0, p_0, p_1, p_2) = ([0], A)$. Hence $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$ is a transversal line for the exterior splash \mathbb{S} . The planes of the regular 2-spread \mathcal{S} are $[S_k] = \{([kx], [x]) : x \in \mathbb{F}'_{q^3}\}$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. We first show that the extended plane $[S_k]^*$ meets the line $g_{\mathbb{S}}$ in the point $kA_1 + A_2$. Consider the point $P = p_0([k], [1]) + p_1([k\tau], [\tau]) + p_2([k\tau^2], [\tau^2])$ of $\text{PG}(5, q^3)$, and note that $P \in [S_k]^*$. Using the matrix M_k defined in Section 2B, we have

$$P = p_0(M_k[1], [1]) + p_1(M_k[\tau], [\tau]) + p_2(M_k[\tau^2], [\tau^2]) = (M_k A, A) = (kA, A)$$

by (1). Hence $P = kA_1 + A_2$ which lies in $g_{\mathbb{S}} = \langle A_1, A_2 \rangle$, that is, P is the intersection of $g_{\mathbb{S}}$ and $[S_k]^*$ proving part (i).

Consider the homography β defined in (7), acting on $\text{PG}(5, q^3)$. The proof of Lemma 5.1 shows that β maps $g_{\mathbb{S}}$ to $g_{\mathbb{T}}$, and maps $g_{\mathbb{T}}$ to $g_{\mathbb{C}}$. Each element $y \in \mathbb{F}'_{q^3}$ can be considered as a point $[y]$ in $\text{PG}(2, q)$. The collineation of $\text{PG}(2, q)$ mapping the point $[y]$ to $[y^q]$ is a homography, and can be represented using a matrix N with entries in \mathbb{F}_q . We omit the transpose notation, and write $N[y] = [y^q]$. Hence we can write the collineation β as $\beta([x], [y]) = ([x], N[y])$. Clearly $\beta(A_1) = A_1$, and we show that $\beta(A_2) = A_2^{q^2}$. Recall the point $A = (p_0, p_1, p_2) = p_0[1] + p_1[\tau] + p_2[\tau^2]$, so $NA = p_0[1] + p_1[\tau^q] + p_2[\tau^{2q}]$. Using the matrix M_k from Section 2B, it is straightforward to write this as $NA = (p_0^{q^2} I + p_1^{q^2} M_{\tau} + p_2^{q^2} M_{\tau^2})^q [1]$. Now

$$(p_0^{q^2} I + p_1^{q^2} M_{\tau} + p_2^{q^2} M_{\tau^2}) [1] = A^{q^2} \quad \text{and} \quad (p_0^{q^2} I + p_1^{q^2} M_{\tau} + p_2^{q^2} M_{\tau^2}) A^{q^2} = \eta^{q^2} A^{q^2}$$

by (1). So repeated use of (1) yields $NA = \eta^{q^2(q-1)} A^{q^2} = \eta^{1-q^2} A^{q^2}$. Further, as N is over \mathbb{F}_q , we have

$$NA = \eta^{1-q^2} A^{q^2}, \quad NA^q = \eta^{q-1} A, \quad NA^{q^2} = \eta^{q^2-q} A^q. \quad (8)$$

Hence $\beta(kA_1 + A_2) = kA_1 + \eta^{1-q^2} A_2^{q^2}$. As $\beta : g_{\mathbb{S}} \mapsto g_{\mathbb{T}}$, we have $g_{\mathbb{T}} \cap [T_k]^* = kA_1 + \eta^{1-q^2} A_2^{q^2}$ and $g_{\mathbb{T}} = \langle A_1, A_2^{q^2} \rangle$, proving part (ii). Similarly, calculating

$$\beta(kA_1 + \eta^{1-q^2} A_2^{q^2}) = kA_1 + \eta^{1-q^2+q^2-q} A_2^q = kA_1 + \eta^{1-q} A_2^q,$$

and using $\beta : g_{\mathbb{T}} \mapsto g_{\mathbb{C}}$, we get $g_{\mathbb{C}} \cap [C_k]^* = kA_1 + \eta^{1-q} A_2^q$ and $g_{\mathbb{C}} = \langle A_1, A_2^q \rangle$. \square

We can use the transversals of the covers \mathbb{T} and \mathbb{C} to generalise the notion of \mathcal{S} -special conics and twisted cubics in $\text{PG}(6, q)$ defined in [Definition 2.2](#). We define \mathbb{C} -special here, \mathbb{T} -special is similarly defined.

- Definition 6.4.** (i) A \mathbb{C} -special conic is a nondegenerate conic \mathcal{C} contained in a plane of \mathbb{C} , such that the extension of \mathcal{C} to $\text{PG}(6, q^3)$ meets the transversals of \mathbb{C} .
- (ii) A \mathbb{C} -special twisted cubic is a twisted cubic \mathcal{N} in a 3-space of $\text{PG}(6, q) \setminus \Sigma_\infty$ about a plane of \mathbb{C} , such that the extension of \mathcal{N} to $\text{PG}(6, q^3)$ meets the transversals of \mathbb{C} .

6B. Characterising the carriers in $\text{PG}(6, q)$. Letting \mathcal{S} be a regular 2-spread of $\text{PG}(5, q)$, and \mathbb{S} be an exterior splash contained in \mathcal{S} , with covers \mathbb{C} and \mathbb{T} , we can then characterise the carriers of \mathbb{S} in terms of the nine transversals of \mathbb{S} , \mathbb{C} and \mathbb{T} .

Theorem 6.5. *Let \mathcal{S} be a regular 2-spread of $\text{PG}(5, q)$, and let $\mathbb{S} \subset \mathcal{S}$ be an exterior splash with covers \mathbb{C} , \mathbb{T} , whose corresponding triples of transversal lines are $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}$, $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$, and $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$, respectively. Then the carriers of \mathbb{S} are the only two planes of \mathcal{S} whose extension to $\text{PG}(5, q^3)$ meets all nine transversal lines.*

Proof. By [Theorem 2.4](#), we can without loss of generality show this for the exterior splash \mathbb{S} of the exterior order- q -subplane \mathcal{B} coordinatised in [Section 3](#), with carriers $E_1 = (1, 0, 0)$, $E_2 = (0, 1, 0)$. In $\text{PG}(6, q)$, the transversal lines $g_{\mathbb{S}}, g_{\mathbb{S}}^q, g_{\mathbb{S}}^{q^2}$ each meet the carriers $[E_1], [E_2]$ of \mathbb{S} . We use the notation for planes $[S_k] \in \mathbb{S}$, $[T_k] \in \mathbb{T}$ and $[C_k] \in \mathbb{C}$ from [Lemma 5.1](#). By [Corollary 6.2](#), in the cubic extension $\text{PG}(5, q^3)$, the transversal lines $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ meet each plane $[T_k]$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$; and the transversal lines $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ meet each plane $[C_k]$, $k \in \mathbb{F}_{q^3} \cup \{\infty\}$. The carriers of \mathbb{S} satisfy $[E_2] = [S_0] = [T_0] = [C_0]$ and $[E_1] = [S_\infty] = [T_\infty] = [C_\infty]$. Hence in the cubic extension $\text{PG}(5, q^3)$, all nine transversal lines meet the carriers of \mathbb{S} .

We now show that no other plane of the regular 2-spread \mathcal{S} meets all nine transversal lines. We use the homography with matrix M_k defined in [Section 2B](#). A plane of the regular 2-spread \mathcal{S} distinct from $[E_1], [E_2]$ has the form $[S_k] = \{([kx], [x], 0) : x \in \mathbb{F}'_3\}$, for some $k \in \mathbb{F}'_3$. This plane is spanned by the three points

$$\begin{aligned} S_{0,k} &= ([k], [1], 0) = (M_k(1, 0, 0), (1, 0, 0)), \\ S_{1,k} &= ([k\tau], [\tau], 0) = (M_k(0, 1, 0), (0, 1, 0)) \\ S_{2,k} &= ([k\tau^2], [\tau^2], 0) = (M_k(0, 0, 1), (0, 0, 1)). \end{aligned}$$

Hence the extension $[S_k]^*$ to $\text{PG}(5, q^3)$ contains the points

$$S_{k,j} = c_0 S_{0,j} + c_1 S_{1,j} + c_2 S_{2,j},$$

where $c_i \in \mathbb{F}_{q^3}$, not all zero. By [Theorem 6.3](#), a general point X on the transversal line $g_{\mathbb{T}}$ has coordinates $X = rA_1 + A_2^{q^2} = (rp_0, rp_1, rp_2, p_0^{q^2}, p_1^{q^2}, p_2^{q^2})$, for some $r \in \mathbb{F}_{q^3} \cup \{\infty\}$. Now $S_{j,k} = X$ if and only if $c_i = p_i^{q^2}$, $i = 0, 1, 2$, and $M_k(c_0, c_1, c_2) = r(p_0, p_1, p_2)$. That is, $M_k A^{q^2} = rA$. However, $M_k A^{q^2} = k^{q^2} A^{q^2}$, by [\(1\)](#), so there are no solutions to c_0, c_1, c_2 . Hence the transversal line $g_{\mathbb{T}}$ does not meet any further plane of the regular 2-spread \mathcal{S} , and so $g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ do not meet any further plane of \mathcal{S} . A similar argument shows that the lines $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ do not meet any further plane of the regular 2-spread \mathcal{S} . \square

6C. Transversal lines of exterior splashes with common carriers. As exterior splashes are equivalent to covers of the circle geometry $\text{CG}(3, q)$, there are $q - 1$ disjoint exterior splashes on ℓ_{∞} with common carriers E_1, E_2 . We show that in $\text{PG}(6, q)$, the covers of these disjoint exterior splashes have common transversals.

Theorem 6.6. *Let $\mathbb{S}_0, \dots, \mathbb{S}_{q-1}$ be $q - 1$ disjoint exterior splashes on ℓ_{∞} with common carriers E_1, E_2 , and let exterior splash \mathbb{S}_j have covers $\mathbb{C}_j, \mathbb{T}_j$. Then the covers $\mathbb{C}_0, \dots, \mathbb{C}_{q-1}$ have common transversal lines $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$, and the covers $\mathbb{T}_0, \dots, \mathbb{T}_{q-1}$ have common transversal lines $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$.*

Proof. By [Theorem 2.4](#), we can without loss of generality prove this for the order- q -subplane \mathcal{B} coordinatised in [Section 3](#). Let $\mathcal{K} = \{k \in \mathbb{F}'_{q^3} : k^{q^2+q+1} = 1\} = \{k = \tau^{i(q-1)} : 0 \leq i < q^2 + q + 1\}$. Recall that \mathcal{B} has carriers $E_1 = (1, 0, 0)$, $E_2 = (0, 1, 0)$, and exterior splash $\mathbb{S}_0 = \{S_{k,0} = (k, 1, 0) : k \in \mathcal{K}\}$. Let $\mathcal{K}_j = \tau^j \mathcal{K}$, for $j = 0, \dots, q-2$, be the $q-1$ cosets of \mathcal{K} in \mathbb{F}'_{q^3} . Let $\mathbb{S}_j = \{S_{k,j} = (k, 1, 0) : k \in \mathcal{K}_j\}$, $0 \leq j \leq q-2$. Consider the homography ξ acting on ℓ_{∞} that maps the point $(x, y, 0)$ to $(\tau x, y, 0)$. Then ξ fixes E_1, E_2 , maps \mathbb{S}_j to \mathbb{S}_{j+1} ($0 \leq j \leq q-3$), and maps \mathbb{S}_{q-2} to \mathbb{S}_0 . Hence $\mathbb{S}_0, \dots, \mathbb{S}_{q-1}$ are the $q - 1$ disjoint exterior splashes on ℓ_{∞} with carriers $(1, 0, 0)$ and $(0, 1, 0)$.

In $\Sigma_{\infty} \cong \text{PG}(5, q)$, we have planes $[S_{k,j}] = \{([kx], [x]) : x \in \mathbb{F}'_{q^3}\} \in \mathbb{S}$, and define the planes $[T_{k,j}] = \{([kx], [x^q]) : x \in \mathbb{F}'_{q^3}\}$, and $[C_{k,j}] = \{([kx], [x^{q^2}]) : x \in \mathbb{F}'_{q^3}\}$, for $k \in \mathcal{K}_j$. So $\mathbb{S}_j = \{[S_{k,j}], k \in \mathcal{K}_j\}$, and define $\mathbb{T}_j = \{[T_{k,j}], k \in \mathcal{K}_j\}$ and $\mathbb{C}_j = \{[C_{k,j}], k \in \mathcal{K}_j\}$. Note that $\mathbb{T}_0, \mathbb{C}_0$ are the covers of the exterior splash \mathbb{S}_0 of \mathcal{B} . Now consider the map θ_{τ} of $\text{PG}(5, q)$ acting on Σ_{∞} defined in [Section 2B](#); it maps \mathbb{S}_j to \mathbb{S}_{j+1} , \mathbb{T}_j to \mathbb{T}_{j+1} , and \mathbb{C}_j to \mathbb{C}_{j+1} . Hence \mathbb{T}_j and \mathbb{C}_j are covers for \mathbb{S}_j . By [Theorem 6.3](#), the transversal line of \mathbb{T}_0 is $g_{\mathbb{T}} = \langle A_1, A_2^{q^2} \rangle$. Using [\(1\)](#), we see that the homography θ_{τ} fixes $g_{\mathbb{T}}$, and so $g_{\mathbb{T}}$ is a transversal for all \mathbb{T}_j . So $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$ are transversal lines of \mathbb{T}_j for each $j = 0, \dots, q-2$. Similarly, $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ are transversal lines of \mathbb{C}_j for each $j = 0, \dots, q-2$. \square

Remark 6.7. We can interpret this result using the terminology of [\[Culbert and Ebert 2005\]](#). We can partition the planes of a regular 2-spread into $q - 1$ disjoint hyper-reguli with common carriers. Each hyper-regulus has two replacement hyper-reguli, which correspond to our conic and tangent covers. If we replace all $q - 1$

hyper-reguli of \mathcal{S} with hyper-reguli of the *same type* (that is, all belonging to \mathbb{C} , or all belonging to \mathbb{T}), then the resulting 2-spread has transversals either $g_{\mathbb{C}}, g_{\mathbb{C}}^q, g_{\mathbb{C}}^{q^2}$ or $g_{\mathbb{T}}, g_{\mathbb{T}}^q, g_{\mathbb{T}}^{q^2}$, and so is regular. Hence the resulting André plane is Desarguesian. If we replace all the hyper-reguli of \mathcal{S} with a combination of hyper-reguli from each type, then the resulting 2-spread is not regular, and so the resulting André plane is non-Desarguesian.

7. Sublines of an exterior splash

In this section we characterise the order- q -sublines of \mathbb{S} with respect to the covers of \mathbb{S} and their transversal lines.

7A. Background. Let π be an exterior order- q -subplane of $\text{PG}(2, q^3)$ with exterior splash \mathbb{S} on ℓ_{∞} . There are $2(q^2 + q + 1)$ order- q -sublines in an exterior splash which lie in two families of size $q^2 + q + 1$. These families are studied in [Lavrauw and Van de Voorde 2010; Barwick and Jackson 2016].

We first describe properties of the two families given in [Lavrauw and Van de Voorde 2010]; here the two families are called regular and irregular with respect to a plane in one of the covers. That is, let \mathbb{S} be an exterior splash in $\text{PG}(5, q)$, and let α be a plane that meets each plane of \mathbb{S} in a point, so α lies in one of the covers \mathbb{X} or \mathbb{Y} of \mathbb{S} . In $\text{PG}(2, q^3)$, let b be an \mathbb{F}_q -subline contained in \mathbb{S} , so by Theorem 2.3, in $\text{PG}(6, q)$, $[b]$ is a 2-regulus. The subline b is called regular with respect to α if $\alpha \cap [b]$ is a line, otherwise b is irregular. Suppose α lies in the cover \mathbb{X} , and $\alpha \cap [b]$ is a line, then each plane in the cover \mathbb{X} meets $[b]$ in a line, and each plane in the cover \mathbb{Y} meets $[b]$ in a set of points which is not collinear. We adapt the phrases regular and irregular with respect to α in terms of the covers of \mathbb{S} . We say b is both \mathbb{X} -regular and \mathbb{Y} -irregular if each plane in \mathbb{X} meets $[b]$ in a line. In particular, we note that if we start with a scattered \mathbb{F}_q -linear set of rank 3 of $\text{PG}(1, q^3)$, then an \mathbb{F}_q -subline b contained in the linear set can be categorised as both regular and irregular (by choosing α in different covers).

In [Lunardon and Polverino 2004], it is shown that if \mathbb{S} is an exterior splash of ℓ_{∞} in $\text{PG}(2, q^3)$, then there is an order- q -subplane β and point P such that \mathbb{S} is the projection of β from P onto ℓ_{∞} . In [Barwick and Jackson 2016, Theorem 5.2], the projection and splash constructions are compared, and it is shown that in almost all cases, the projection and exterior splash of β are distinct. In [Lavrauw and Van de Voorde 2010], the two families of sublines of \mathbb{S} are characterised in relation to a point P and subplane β which project \mathbb{S} : one family arises from projecting the sublines of β , the other arises from projecting certain conics of β . The latter family are described as irregular in [Lavrauw and Van de Voorde 2010], although it is not specified which cover these sublines are irregular with respect to.

Now we describe properties of the two families given in [Barwick and Jackson 2016]. Here the two families of order- q -sublines of \mathbb{S} are characterised with respect

to geometric objects of an exterior π with exterior splash \mathbb{S} . If A is a point of π , then the pencil of $q + 1$ lines of π through A meets ℓ_∞ in an order- q -subline of \mathbb{S} , called a π -pencil-subline of \mathbb{S} . Recall from [Section 2D](#) that a (π, ℓ_∞) -carrier-dual conic of π is a dual conic that contains the three lines fixed by the subgroup I fixing π and ℓ . If Γ is a (π, ℓ_∞) -carrier-dual conic of π , then the lines of Γ meet ℓ_∞ in an order- q -subline of \mathbb{S} , called a π -dual-conic-subline of \mathbb{S} . Note that in [\[Barwick and Jackson 2016, Theorem 4.4\]](#), we show that it is possible to switch the roles of the two families by considering different associated order- q -subplanes.

7B. A characterisation of the sublines of an exterior splash. We now consider the interaction in $\text{PG}(6, q)$ of the two families of order- q -sublines of \mathbb{S} with the two covers of \mathbb{S} . We show in [Theorem 7.1](#) that each family meets planes from one cover in lines, and planes from the other cover in conics. [Theorem 7.2](#) shows that the converse is true, and so we have a characterisation of the order- q -sublines of \mathbb{S} . This allows us to relate the families from [\[Barwick and Jackson 2016\]](#) and [\[Lavrauw and Van de Voorde 2010\]](#). [Theorem 7.4](#) shows that the conics concerned in each case are special with respect to the conic cover.

Suppose \mathcal{R} is a 2-regulus in $\text{PG}(5, q)$, and consider a plane α that meets \mathcal{R} in a set of $q + 1$ points. Then an easy counting argument shows that these points form either a line or a conic in α . We abbreviate this to “ \mathcal{R} meets α in a line or a conic”.

Theorem 7.1. *Let π be an exterior order- q -subplane with exterior splash \mathbb{S} , conic cover \mathbb{C} , and tangent cover \mathbb{T} .*

- (i) *A π -pencil-subline of \mathbb{S} corresponds in $\text{PG}(6, q)$ to a 2-regulus that meets each plane of \mathbb{T} in a distinct line, and meets each plane of \mathbb{C} in a conic.*
- (ii) *A π -dual-conic-subline of \mathbb{S} corresponds in $\text{PG}(6, q)$ to a 2-regulus that meets each plane of \mathbb{T} in a conic, and meets each plane of \mathbb{C} in a distinct line.*

Proof. Let P be a point in the exterior order- q -subplane π , and let d be the corresponding π -pencil-subline of \mathbb{S} . By [Theorem 2.3](#), in $\text{PG}(6, q)$, $[d]$ is a 2-regulus contained in \mathbb{S} . Consider the tangent plane \mathcal{T}_P to $[\pi]$ at P . By [Theorem 4.2](#), the lines of \mathcal{T}_P through P meet Σ_∞ in points that lie in distinct planes of the 2-regulus $[d]$. Hence $\mathcal{T}_P \cap \Sigma_\infty$ is a ruling line of the 2-regulus $[d]$. By [Theorem 5.3](#), this ruling line $\mathcal{T}_P \cap \Sigma_\infty$ lies in a tangent cover plane. The homography Θ of [Lemma 5.2](#) fixes the planes of $[b]$ and is transitive on the cover planes of \mathbb{T} . Hence each ruling line of $[b]$ meets a unique cover plane of \mathbb{T} .

A straightforward geometric argument shows that planes of \mathbb{T}, \mathbb{C} meet a 2-regulus of \mathbb{S} in a line or a conic. Hence a conic cover plane meets the 2-regulus $[d]$ in a conic. As there are $q^2 + q + 1$ π -pencil-sublines of \mathbb{S} , every line in a plane of \mathbb{T} is a ruling line for some 2-regulus corresponding to a π -pencil-subline. Hence

if $[d']$ is a 2-regulus of \mathbb{S} corresponding to a π -dual-conic-subline, then planes of \mathbb{T} meet $[d']$ in conics, and so planes of \mathbb{C} meet $[d']$ in ruling lines of $[d']$. Moreover, applying the homography of [Lemma 5.2](#) shows that each ruling line of $[d']$ lies in a unique conic cover plane. \square

By [Theorem 2.3](#), there is a one-to-one correspondence between the order- q -sublines of \mathbb{S} in $\text{PG}(2, q^3)$, and the 2-reguli contained in \mathbb{S} in $\text{PG}(6, q)$. Hence the converse of [Theorem 7.1](#) is also true, and so we have a characterisation of order- q -sublines of \mathbb{S} relating to the cover planes of the associated order- q -subplane.

Theorem 7.2. *Let π be an exterior order- q -subplane with exterior splash \mathbb{S} , conic cover \mathbb{C} , and tangent cover \mathbb{T} .*

- (i) *A 2-regulus contained in \mathbb{S} that meets some plane of \mathbb{T} in a line corresponds to a π -pencil-subline of \mathbb{S} .*
- (ii) *A 2-regulus contained in \mathbb{S} that meets some plane of \mathbb{C} in a conic corresponds to a π -pencil-subline of \mathbb{S} .*
- (iii) *A 2-regulus contained in \mathbb{S} that meets some plane of \mathbb{T} in a conic corresponds to a π -dual-conic-subline of \mathbb{S} .*
- (iv) *A 2-regulus contained in \mathbb{S} that meets some plane of \mathbb{C} in a line corresponds to a π -dual-conic-subline of \mathbb{S} .*

This allows us to determine the relationship between the different family naming used in [\[Barwick and Jackson 2016\]](#) and [\[Lavrauw and Van de Voorde 2010\]](#).

Corollary 7.3. *Let π be an exterior order- q -subplane with exterior splash \mathbb{S} , conic cover \mathbb{C} , and tangent cover \mathbb{T} .*

- (i) *Let b be a π -pencil-subline of \mathbb{S} , then b is \mathbb{T} -regular and \mathbb{C} -irregular.*
- (ii) *Let d be a π -dual-conic-subline of \mathbb{S} , then d is \mathbb{C} -regular and \mathbb{T} -irregular.*

In fact, we can give a stronger characterisation of the order- q -sublines of \mathbb{S} , namely that the conics of [Theorem 7.1](#) are *special* with respect to the associated cover. In order to prove that the conics are special, we need to introduce coordinates, and the proof is calculation intensive.

Theorem 7.4. *Let π be an exterior order- q -subplane with exterior splash \mathbb{S} , conic cover \mathbb{C} , and tangent cover \mathbb{T} .*

- (i) *A 2-regulus of \mathbb{S} corresponding to a π -pencil-subline of \mathbb{S} meets each plane of \mathbb{C} in a \mathbb{C} -special conic.*
- (ii) *A 2-regulus of \mathbb{S} corresponding to a π -dual-conic-subline of \mathbb{S} meets each plane of \mathbb{T} in a \mathbb{T} -special conic.*

Proof. By [Theorem 2.4](#), we can without loss of generality prove this for the exterior order- q -subplane \mathcal{B} coordinatised in [Section 3](#). We start with the order- q -subplane $\pi_0 = \text{PG}(2, q)$ and the line $\ell = [-\tau\tau^q, \tau + \tau^q, -1]$ which is exterior to π_0 . Note that using the notation for p_0, p_1, p_2 given in [Theorem 6.3](#), we have $\ell = [p_0^{q^2}, p_1^{q^2}, p_2^{q^2}]$. A line of π_0 has coordinates $[l, m, n]$ for $l, m, n \in \mathbb{F}_q$, and meets ℓ in the point $W'_{l,m,n} = (-n(\tau + \tau^q) - m, l - n\tau\tau^q, m\tau\tau^q + l(\tau + \tau^q))$. We apply the homography σ of [Section 3](#) with matrix K to map π_0 and ℓ to \mathcal{B} and ℓ_∞ , respectively. The point $W'_{l,m,n}$ of ℓ maps to the point $W_{l,m,n} = KW'_{l,m,n} = (l + m\tau + n\tau^2, l + m\tau^q + n\tau^{2q}, 0)$ of ℓ_∞ . Writing $\varepsilon = \varepsilon_{l,m,n} = l + m\tau + n\tau^2$, we have $W_\varepsilon = W_{l,m,n} = (\varepsilon, \varepsilon^q, 0) \equiv (\varepsilon^{1-q}, 1, 0)$. Using the notation from [Lemma 5.1](#), this is the point $S_{\varepsilon^{1-q}} \in \ell_\infty$. In $\text{PG}(6, q)$, W_ε corresponds to the spread plane $[W_\varepsilon] = [W_{l,m,n}] = \{([\varepsilon x], [\varepsilon^q x], 0) \equiv ([\varepsilon^{1-q} x], [x], 0) : x \in \mathbb{F}'_q\} = [S_{\varepsilon^{1-q}}]$.

Fix a point $P = (a, b, c)$ of π_0 , so $a, b, c, \in \mathbb{F}_q$, not all zero. Let

$$\mathcal{L} = \{(l, m, n) : l, m, n \in \mathbb{F}_q \text{ not all zero, and } la + mb + nc = 0\}.$$

The $q + 1$ lines of π_0 through P have coordinates $[l, m, n] \in \mathcal{L}$. These $q + 1$ lines meet the exterior line ℓ of π_0 in a π_0 -pencil-subline which, under the collineation σ , maps to a \mathcal{B} -pencil-subline d of ℓ_∞ . By [Theorem 2.3](#), in $\text{PG}(6, q)$, d corresponds to the 2-regulus $[d]$ which we denote by \mathcal{R} , so $\mathcal{R} = [d] = \{[W_\varepsilon] = [S_{\varepsilon^{1-q}}] : \varepsilon \in \mathcal{W}\}$, where $\mathcal{W} = \{\varepsilon = \varepsilon_{l,m,n} = l + m\tau + n\tau^2 : (l, m, n) \in \mathcal{L}\}$. For each $\alpha \in \mathbb{F}'_q$, consider the set of points $t_\alpha = \{([\varepsilon\alpha], [\varepsilon^q\alpha], 0) : \varepsilon \in \mathcal{W}\}$. As \mathcal{W} is closed under addition, t_α is a line of $\Sigma_\infty \cong \text{PG}(5, q)$; further t_α meets every plane in \mathcal{R} . Hence t_α is a ruling line of the 2-regulus \mathcal{R} .

By [Theorem 7.2\(ii\)](#), the 2-regulus \mathcal{R} meets a cover plane of the conic cover \mathbb{C} in a conic $\mathcal{C}_k = [C_k] \cap \mathcal{R}$ for $k \in \mathcal{K}$. To show that the conic \mathcal{C}_k is \mathbb{C} -special, we need to extend it to $\text{PG}(5, q^3)$, and show that it meets the three transversal lines of \mathbb{C} . To do this, we extend the 2-regulus \mathcal{R} of $\Sigma_\infty \cong \text{PG}(5, q)$ to a 2-regulus \mathcal{R}^* of $\text{PG}(5, q^3)$, so $\mathcal{C}_k^* = [C_k]^* \cap \mathcal{R}^*$. We then use coordinates to show that one of the planes of \mathcal{R}^* contains the transversal line $g_{\mathbb{C}}^{q^2}$ of \mathbb{C} , and then deduce that \mathcal{C}_k^* meets $g_{\mathbb{C}}^{q^2}$.

To extend \mathcal{R} to a 2-regulus \mathcal{R}^* of $\text{PG}(5, q^3)$, we find four lines in $\text{PG}(5, q^3)$ that meet each extended plane of \mathcal{R} . As a 2-regulus is uniquely determined by four ruling lines in general position, we can use these four lines to define the 2-regulus \mathcal{R}^* . The transversal line $g_{\mathbb{S}}$ of the regular 2-spread \mathcal{S} can be used as one of our ruling lines; for the other three ruling lines, we use the extended lines t_1^* , t_τ^* , $t_{\tau^2}^*$, which each meet every plane of \mathcal{R} . So \mathcal{R}^* is the 2-regulus of $\text{PG}(5, q^3)$ determined by the four ruling lines t_1^* , t_τ^* , $t_{\tau^2}^*$, $g_{\mathbb{S}}$ (which are in general position), and further $\mathcal{R}^* \cap \Sigma_\infty = \mathcal{R}$.

We now exhibit a plane γ of \mathcal{R}^* that contains the transversal line $g_{\mathbb{C}}^{q^2}$ of the conic cover \mathbb{C} . Extend the set \mathcal{L} to

$$\mathcal{L}^* = \{(l, m, n) : l, m, n \in \mathbb{F}_{q^3} \text{ not all zero, and } la + mb + nc = 0\}.$$

We use the matrix M_τ defined in [Section 2B](#), and write $M = M_\tau$. The ruling line t_τ^* , $i = 0, 1, 2$, has points $P_{\tau^i, l, m, n}$ with $(l, m, n) \in \mathcal{L}^*$, where $P_{\tau^i, l, m, n} = l(M^i[1], M^i[1], 0) + m(M^i[\tau], M^i[\tau^q], 0) + n(M^i[\tau^2], M^i[\tau^{2q}], 0)$. Recall that the order- q -subline d corresponds to the fixed point $P = (a, b, c) \in \pi_0$. Consider the following $(l, m, n) \in \mathcal{L}^*$:

$$l = c\tau - b\tau^2, \quad m = a\tau^2 - c, \quad n = b - a\tau. \quad (9)$$

Note that for these l, m, n we have

$$l + m\tau + n\tau^2 = 0. \quad (10)$$

For l, m, n as in (9), consider the plane γ spanned by the three points $P_{1, l, m, n} \in t_1^*$, $P_{\tau, l, m, n} \in t_\tau^*$, $P_{\tau^2, l, m, n} \in t_{\tau^2}^*$. We first show that γ is a plane of the 2-regulus \mathcal{R}^* by showing that the fourth ruling line $g_\mathbb{S}$ of \mathcal{R}^* also meets γ . By [Theorem 6.3](#), $g_\mathbb{S} = \langle A_1, A_2 \rangle$, and we show that $g_\mathbb{S}$ meets γ by showing that the point A_2 lies in γ . With l, m, n given by (9), consider the point $F = p_0 P_{1, l, m, n} + p_1 P_{\tau, l, m, n} + p_2 P_{\tau^2, l, m, n}$ of γ . To simplify the notation, we use the point $A = (p_0, p_1, p_2)^t$, and matrix $U_0 = p_0 I + p_1 M + p_2 M^2$ defined in [Section 2B](#), and note that $U_0[\alpha] = \alpha A$. We have

$$\begin{aligned} F &= (lU_0[1] + mU_0[\tau] + nU_0[\tau^2], lU_0[1] + mU_0[\tau^q] + nU_0[\tau^{2q}], 0) \\ &= (lA + m\tau A + n\tau^2 A, lA + m\tau^q A + n\tau^{2q} A, 0). \end{aligned}$$

By (10), $F \equiv ([0], A, 0) = A_2$, and by [Lemma 2.1](#), $g_\mathbb{S} = \langle A_1, A_2 \rangle$, so $F \in g_\mathbb{S} \cap \gamma$. That is, the four ruling lines $t_1^*, t_\tau^*, t_{\tau^2}^*, g_\mathbb{S}$ of the 2-regulus \mathcal{R}^* all meet the plane γ , and so γ is a plane of \mathcal{R}^* .

We now show that the transversal line $g_\mathbb{C}^{q^2}$ of \mathbb{C} lies in the plane γ of \mathcal{R}^* . Let $G = p_0^{q^2} P_{1, l, m, n} + p_1^{q^2} P_{\tau, l, m, n} + p_2^{q^2} P_{\tau^2, l, m, n}$, and note that $G \in \gamma$. We use the matrix

$$U_2 = p_0^{q^2} I + p_1^{q^2} M + p_2^{q^2} M^2$$

defined in [Section 2B](#), and note that $U_2[\alpha] = \alpha^{q^2} A^{q^2}$, so we have

$$\begin{aligned} G &= (lU_2[1] + mU_2[\tau] + n^2 U_2[\tau^2], lU_2[1] + mU_2[\tau^q] + nU_2[\tau^{2q}], 0) \\ &= (lA^{q^2} + m\tau^{q^2} A^{q^2} + n\tau^{2q^2} A^{q^2}, lA^{q^2} + m\tau A^{q^2} + n\tau^2 A^{q^2}, 0). \end{aligned}$$

By (10), $G \equiv (A^{q^2}, [0], 0) = A_1^{q^2}$, so γ contains the points $G = A_1^{q^2}$ and $F = A_2$. Hence by [Theorem 6.3](#), γ contains the transversal line $g_\mathbb{C}^{q^2} = \langle A_1^{q^2}, A_2 \rangle$ of \mathbb{C} .

We showed above that the 2-regulus $[d] = \mathcal{R}$ meets a cover plane $[C_i]$ of \mathbb{C} in a conic \mathcal{C}_i . We want to show that \mathcal{C}_i is a \mathbb{C} -special conic, that is, we want to show that in $\text{PG}(6, q^3)$, the extended conic $\mathcal{C}_i^* = [C_i]^* \cap \mathcal{R}^*$ contains the three points $g_\mathbb{C} \cap [C_i]^*$, $g_\mathbb{C}^q \cap [C_i]^*$, $g_\mathbb{C}^{q^2} \cap [C_i]^*$. We have shown that the transversal line $g_\mathbb{C}^{q^2}$ of \mathbb{C} lies in a plane γ of \mathcal{R}^* . As the extended cover plane $[C_i]^*$ meets the transversal line $g_\mathbb{C}^{q^2}$ in a unique point denoted P_i , we have

$$P_i = [C_i]^* \cap g_\mathbb{C}^{q^2} = [C_i]^* \cap \gamma \in [C_i]^* \cap \mathcal{R}^* = \mathcal{C}_i^*.$$

Hence C_i^* contains the point $g_{\mathbb{C}}^{q^2} \cap [C_i]^*$, and hence it also contains the conjugate points $g_{\mathbb{C}}^q \cap [C_i]^*$, $g_{\mathbb{C}} \cap [C_i]^*$. That is, the conic $C_i = [C_i] \cap \mathcal{R}$ is a \mathbb{C} -special conic, completing the proof of part (i). As \mathbb{C} and \mathbb{T} are projectively equivalent by [Lavrauw et al. 2015, Theorem 16], part (ii) holds by symmetry. \square

8. Conclusion

An investigation into the interaction between an exterior order- q -subplane π of $\text{PG}(2, q^3)$, and its exterior splash on ℓ_{∞} began in [Barwick and Jackson 2016]. The main focus of that paper was to show that exterior splashes are projectively equivalent to scattered \mathbb{F}_q -linear sets of rank 3, covers of circle geometries, Sherk sets of size $q^2 + q + 1$. Further, we investigated the geometric relationship between the order- q -sublines of \mathbb{S} and the points of π . The current article focusses on using the Bruck–Bose representation in $\text{PG}(6, q)$ to continue the study of exterior splashes, in particular their interplay with order- q -subplanes. The notion of special conics and special twisted cubics is closely tied with this interplay.

References

- [André 1954] J. André, “Über nicht-Desarguessche Ebenen mit transitiver Translationsgruppe”, *Math. Z.* **60** (1954), 156–186. [MR](#) [Zbl](#)
- [Barwick and Jackson 2012] S. G. Barwick and W.-A. Jackson, “Sublines and subplanes of $\text{PG}(2, q^3)$ in the Bruck–Bose representation in $\text{PG}(6, q)$ ”, *Finite Fields Appl.* **18**:1 (2012), 93–107. [MR](#) [Zbl](#)
- [Barwick and Jackson 2014] S. G. Barwick and W.-A. Jackson, “A characterisation of tangent subplanes of $\text{PG}(2, q^3)$ ”, *Des. Codes Cryptogr.* **71**:3 (2014), 541–545. [MR](#) [Zbl](#)
- [Barwick and Jackson 2015] S. G. Barwick and W.-A. Jackson, “An investigation of the tangent splash of a subplane of $\text{PG}(2, q^3)$ ”, *Des. Codes Cryptogr.* **76**:3 (2015), 451–468. [MR](#) [Zbl](#)
- [Barwick and Jackson 2016] S. G. Barwick and W.-A. Jackson, “Exterior splashes and linear sets of rank 3”, *Discrete Math.* **339**:5 (2016), 1613–1623. [MR](#) [Zbl](#)
- [Barwick and Jackson 2017] S. G. Barwick and W.-A. Jackson, “The exterior splash in $\text{PG}(6, q)$: carrier conics”, *Adv. Geom.* **17**:4 (2017), 407–422. [MR](#) [Zbl](#)
- [Bruck 1969] R. H. Bruck, “Construction problems of finite projective planes”, pp. 426–514 in *Combinatorial Mathematics and its Applications* (Chapel Hill, N.C., 1967), edited by D. J. A. Welsh, Univ. North Carolina Press, Chapel Hill, N.C., 1969. [MR](#) [Zbl](#)
- [Bruck 1973] R. H. Bruck, “Circle geometry in higher dimensions, II”, *Geometriae Dedicata* **2** (1973), 133–188. [MR](#) [Zbl](#)
- [Bruck and Bose 1964] R. H. Bruck and R. C. Bose, “The construction of translation planes from projective spaces”, *J. Algebra* **1** (1964), 85–102. [MR](#) [Zbl](#)
- [Bruck and Bose 1966] R. H. Bruck and R. C. Bose, “Linear representations of projective planes in projective spaces”, *J. Algebra* **4** (1966), 117–172. [MR](#) [Zbl](#)
- [Culbert and Ebert 2005] C. Culbert and G. L. Ebert, “Circle geometry and three-dimensional subregular translation planes”, *Innov. Incidence Geom.* **1** (2005), 3–18. [MR](#) [Zbl](#)
- [Hirschfeld and Thas 1991] J. W. P. Hirschfeld and J. A. Thas, *General Galois geometries*, The Clarendon Press, Oxford University Press, New York, 1991. [MR](#) [Zbl](#)

- [Lavrauw 2016] M. Lavrauw, “Scattered spaces in Galois geometry”, pp. 195–216 in *Contemporary developments in finite fields and applications*, edited by A. Canteaut et al., World Sci. Publ., Hackensack, NJ, 2016. [MR](#) [Zbl](#)
- [Lavrauw and Van de Voorde 2010] M. Lavrauw and G. Van de Voorde, “On linear sets on a projective line”, *Des. Codes Cryptogr.* **56**:2-3 (2010), 89–104. [MR](#) [Zbl](#)
- [Lavrauw and Zanella 2015] M. Lavrauw and C. Zanella, “Subgeometries and linear sets on a projective line”, *Finite Fields Appl.* **34** (2015), 95–106. [MR](#) [Zbl](#)
- [Lavrauw et al. 2015] M. Lavrauw, J. Sheekey, and C. Zanella, “On embeddings of minimum dimension of $\text{PG}(n, q) \times \text{PG}(n, q)$ ”, *Des. Codes Cryptogr.* **74**:2 (2015), 427–440. [MR](#) [Zbl](#)
- [Lunardon and Polverino 2004] G. Lunardon and O. Polverino, “Translation ovoids of orthogonal polar spaces”, *Forum Math.* **16**:5 (2004), 663–669. [MR](#) [Zbl](#)
- [Lunardon et al. 2014] G. Lunardon, G. Marino, O. Polverino, and R. Trombetti, “Maximum scattered linear sets of pseudoregulus type and the Segre variety $\mathcal{S}_{n,n}$ ”, *J. Algebraic Combin.* **39**:4 (2014), 807–831. [MR](#) [Zbl](#)
- [Ostrom 1993] T. G. Ostrom, “Hyper-reguli”, *J. Geom.* **48**:1-2 (1993), 157–166. [MR](#) [Zbl](#)
- [Rottey et al. 2015] S. Rottey, J. Sheekey, and G. Van de Voorde, “Subgeometries in the André/Bruck–Bose representation”, *Finite Fields Appl.* **35** (2015), 115–138. [MR](#) [Zbl](#)

Received 2 Feb 2015.

SUSAN G. BARWICK:

susan.barwick@adelaide.edu.au

School of Mathematical Sciences, University of Adelaide, Australia

WEN-AI JACKSON:

wen.jackson@adelaide.edu.au

School of Mathematical Sciences, University of Adelaide, Australia

Innovations in Incidence Geometry

msp.org/iig

MANAGING EDITOR

Tom De Medts	Ghent University tom.demedts@ugent.be
Linus Kramer	Universität Münster linus.kramer@wwu.de
Klaus Metsch	Justus-Liebig Universität Gießen klaus.metsch@math.uni-giessen.de
Bernhard Mühlherr	Justus-Liebig Universität Gießen bernhard.m.muehlherr@math.uni-giessen.de
Joseph A. Thas	Ghent University thas.joseph@gmail.com
Koen Thas	Ghent University koen.thas@gmail.com
Hendrik Van Maldeghem	Ghent University hendrik.vanmaldeghem@ugent.be

HONORARY EDITORS

Jacques Tits
Ernest E. Shult †

EDITORS

Peter Abramenko	University of Virginia
Francis Buekenhout	Université Libre de Bruxelles
Philippe Cara	Vrije Universiteit Brussel
Antonio Cossidente	Università della Basilicata
Hans Cuypers	Eindhoven University of Technology
Bart De Bruyn	University of Ghent
Alice Devillers	University of Western Australia
Massimo Giulietti	Università degli Studi di Perugia
James Hirschfeld	University of Sussex
Dimitri Leemans	Université Libre de Bruxelles
Oliver Lorscheid	Instituto Nacional de Matemática Pura e Aplicada (IMPA)
Guglielmo Lunardon	Università di Napoli “Federico II”
Alessandro Montinaro	Università di Salento
James Parkinson	University of Sydney
Antonio Pasini	Università di Siena (emeritus)
Valentina Pepe	Università di Roma “La Sapienza”
Bertrand Rémy	École Polytechnique
Tamás Szonyi	ELTE Eötvös Loránd University, Budapest

PRODUCTION

Silvio Levy (Scientific Editor)
production@msp.org

See inside back cover or msp.org/iig for submission instructions.

The subscription price for 2019 is US \$275/year for the electronic version, and \$325/year (+\$20, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Innovations in Incidence Geometry: Algebraic, Topological and Combinatorial (ISSN pending) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

IIG peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Innovation in Incidence Geometry

Vol. 17 No. 1

2019

- The exterior splash in $PG(6, q)$: transversals 1
SUSAN G. BARWICK and WEN-AI JACKSON
- Ruled quintic surfaces in $PG(6, q)$ 25
SUSAN G. BARWICK
- A characterization of Clifford parallelism by automorphisms 43
RAINER LÖWEN
- Generalized quadrangles, Laguerre planes and shift planes of odd order 47
GÜNTER F. STEINKE and MARKUS STROPPEL
- A new family of 2-dimensional Laguerre planes that admit $PSL_2(\mathbb{R}) \times \mathbb{R}$ as a group of automorphisms 53
GÜNTER F. STEINKE

