

# involve

a journal of mathematics

On the number of pairwise touching simplices

Bas Lemmens and Christopher Parsons



# On the number of pairwise touching simplices

Bas Lemmens and Christopher Parsons

(Communicated by Kenneth S. Berenhaut)

In this note, it is shown that the maximum number of pairwise touching translates of an  $n$ -simplex is at least  $n + 3$  for  $n = 7$ , and for all  $n \geq 5$  such that  $n \equiv 1 \pmod{4}$ . The current best known lower bound for general  $n$  is  $n + 2$ . For  $n = 2^k - 1$  and  $k \geq 2$ , we will also present an alternative construction to give  $n + 2$  touching simplices using Hadamard matrices.

## 1. Introduction

A classic problem in discrete geometry is to determine for a given convex body  $K$  in  $\mathbb{R}^n$  the maximum number of pairwise touching translates of  $K$ . This number is called the *touching number* of  $K$  and is denoted by  $t(K)$ . It is well known that for any convex body  $K$  in  $\mathbb{R}^n$ ,

$$t(K) \leq 2^n,$$

and equality holds if, and only if,  $K$  is a parallelopiped; see [Danzer and Grünbaum 1962; Petty 1971; Soltan 1975]. On the other hand, it is unknown if for each convex body  $K$  in  $\mathbb{R}^n$  the inequality  $t(K) \geq n + 1$  holds when  $n \geq 4$ ; see [Bezdek 2010, Section 2.3].

This paper concerns the touching number of  $n$ -dimensional simplices,  $\Delta_n$ . This number was studied by Koolen, Laurent and Schrijver [Koolen et al. 2000]. They showed, among other things, that  $t(\Delta_n) \geq n + 2$  for all  $n \geq 3$  and  $t(\Delta_3) = 5$ , see Figure 1. Lemmens [2007] gave examples that showed that  $t(\Delta_4) \geq 7$  and  $t(\Delta_5) \geq 9$ .

The main goal of this short note is to present a construction that gives the following small improvement of the lower bound for  $t(\Delta_n)$ .

**Theorem 1.1.** *For  $n = 7$  and  $n \equiv 1 \pmod{4}$ , with  $n \geq 5$ , we have that*


$$t(\Delta_n) \geq n + 3.$$

---

*MSC2010:* primary 52C17; secondary 05B40, 46B20.

*Keywords:* touching number, simplices, equilateral sets,  $\ell_1$ -norm.

Parsons is grateful for the support from School of Mathematics, Statistics and Actuarial Science at the University of Kent and from the HE STEM project “Communicating Mathematical Sciences”.



**Figure 1.** Five pairwise touching tetrahedra.

The problem of determining  $t(\Delta_n)$  is known to be equivalent to finding the maximum size of  $\ell_1$ -norm equilateral sets in a hyperplane [Koolen et al. 2000; Lemmens 2007]. We will discuss the equivalence between these two problems in the next section.

## 2. Equilateral sets

A convex body  $K$  in  $\mathbb{R}^n$  which is centrally symmetric, i.e.,  $x \in K$  if and only if  $-x \in K$ , is the unit ball of a norm  $\|\cdot\|_K$  on  $\mathbb{R}^n$ . Indeed, for  $x \in \mathbb{R}^n$ , we can define the norm by

$$\|x\|_K = \inf\{\lambda > 0 : x \in \lambda K\}.$$

A set  $S$  in a normed space  $(\mathbb{R}^n, \|\cdot\|)$  is called an *equilateral set* if there exists a constant  $\delta > 0$  such that

$$\|s - t\| = \delta \quad \text{for all } s \neq t \text{ in } S.$$

The maximum size of an equilateral set in  $(\mathbb{R}^n, \|\cdot\|)$  is the *equilateral dimension* of  $(\mathbb{R}^n, \|\cdot\|)$  and is denoted by  $e(\mathbb{R}^n, \|\cdot\|)$ . Note that the constant  $\delta > 0$  does not play a role, as we can always scale the equilateral set. Clearly, if  $K$  is a centrally symmetric body in  $\mathbb{R}^n$ , then  $S = \{s_1, \dots, s_p\}$  is an equilateral set in  $(\mathbb{R}^n, \|\cdot\|_K)$  with pairwise distance 2 if, and only if, the set of unit balls with centers  $s_1, \dots, s_p$  is a configuration of  $p$  pairwise touching translates of  $K$ .

The equilateral dimension has been studied for many normed spaces; see, for example, [Alon and Pudlák 2003; Swanepoel 2004a; Swanepoel 2004b]. Particular attention has been given to so-called  $\ell_p$ -norms which are defined as follows. For  $1 \leq p < \infty$ , the  $\ell_p$ -norm on  $\mathbb{R}^n$  is given by  $\|x\|_p = (\sum_i |x_i|^p)^{1/p}$ . For the  $\ell_1$ -norm,

$n = 5$	$n = 6$	$n = 8$
(4, 0, 1, 1, 2)	(4, 0, 1, 1, 1, 1)	(0, 4, 2, 2, 0, 4, 2, 2)
(0, 4, 1, 1, 2)	(0, 4, 1, 1, 1, 1)	(4, 0, 2, 2, 4, 0, 2, 2)
(1, 1, 4, 0, 2)	(1, 1, 4, 0, 1, 1)	(2, 2, 0, 4, 2, 2, 0, 4)
(1, 1, 0, 4, 2)	(1, 1, 0, 4, 1, 1)	(2, 2, 4, 0, 2, 2, 4, 0)
(2, 2, 0, 0, 4)	(1, 1, 1, 1, 4, 0)	(8, 2, 1, 1, 0, 2, 1, 1)
(0, 0, 2, 2, 4)	(1, 1, 1, 1, 0, 4)	(4, 4, 4, 4, 0, 0, 0, 0)
(2, 2, 2, 2, 0)	(2, 2, 2, 2, 0, 0)	(4, 4, 0, 0, 4, 4, 0, 0)
	(2, 2, 0, 0, 2, 2)	(4, 4, 0, 0, 0, 0, 4, 4)
	(0, 0, 2, 2, 2, 2)	(4, 0, 4, 0, 0, 4, 0, 4)
		(4, 0, 0, 4, 0, 4, 4, 0)

**Table 1.** Equilateral sets.

it has been conjectured by Kusner that  $e(\mathbb{R}^n, \|\cdot\|_1) = 2n$  [Guy 1983], but at present this has only been confirmed for  $1 \leq n \leq 4$ ; see [Bandelt et al. 1998; Koolen et al. 2000]. Obviously,  $2n$  is a lower bound for  $e(\mathbb{R}^n, \|\cdot\|_1)$ , as the set of standard basis vectors and their opposites form an equilateral set. The best known upper bound is  $Cn \log n$ , where  $C > 0$  is a constant, which was obtained using probabilistic methods by Alon and Pudlák [2003].

Finding the touching number for the  $n$ -dimensional simplex turns out to be equivalent to determining the maximum size of an  $\ell_1$ -norm equilateral set contained in a hyperplane. More precisely, if  $h(n)$  is the maximum size of an  $\ell_1$ -norm equilateral set in  $H_\alpha = \{x \in \mathbb{R}^n : \sum_i x_i = \alpha\}$  for some  $\alpha \in \mathbb{R}$ , then

$$t(\Delta_n) = h(n+1) \quad \text{for all } n \geq 1; \tag{2-1}$$

see [Koolen et al. 2000; Lemmens 2007]. For example, the  $\ell_1$ -norm equilateral set

$$S = \{(2, 0, 1, 1), (0, 2, 1, 1), (1, 1, 2, 0), (1, 1, 0, 2), (2, 2, 0, 0)\} \tag{2-2}$$

in the hyperplane  $H_4 \subseteq \mathbb{R}^4$  corresponds to the configuration of five pairwise touching translates of a tetrahedron depicted in Figure 1. The examples of equilateral sets in Table 1 were found with the aid of a computer. In particular, we see that  $t(\Delta_7) \geq 10$ , which settles the  $n = 7$  case in Theorem 1.1.

It is interesting to note that in these examples all the nonzero coordinates are powers of 2. We have looked into those type of examples in more detail, which led to the construction in Proposition 4.1. At present, however, we have no clear understanding of why these coordinate values generate large examples.

Before we prove Theorem 1.1, we mention that the inequalities

$$h(n) \leq e(\mathbb{R}^n, \|\cdot\|_1) \leq h(2n - 1)$$

are known to hold for all  $n \geq 1$  [Koolen et al. 2000; Lemmens 2007]. Thus,  $e(\mathbb{R}^n, \|\cdot\|_1)$  grows linearly in  $n$  if, and only if,  $h(n)$  does.

### 3. Proof of Theorem 1.1

For each  $n \equiv 2 \pmod{4}$  with  $n \geq 6$ , we shall construct an  $\ell_1$ -norm equilateral set in  $H_\alpha = \{x \in \mathbb{R}^n : \sum_i x_i = \alpha\}$  of size  $n+2$ , where  $\alpha = (n-2)^2/2$ . The result then follows from Equation (2-1). So let  $n \equiv 2 \pmod{4}$  with  $n \geq 6$ . Define

$$\begin{aligned} v^1 &= (b, 0, a, a, \dots, a, a), \\ v^2 &= (0, b, a, a, \dots, a, a), \\ v^3 &= (a, a, b, 0, \dots, a, a), \\ v^4 &= (a, a, 0, b, \dots, a, a), \\ &\vdots \\ v^{n-1} &= (a, a, a, a, \dots, b, 0), \\ v^n &= (a, a, a, a, \dots, 0, b), \end{aligned}$$

in  $\mathbb{R}^n$ , where  $a = (n-4)/2$  and  $b = n-2$ . Furthermore let

$$v^{n+1} = (\underbrace{y, y, \dots, y}_k, \underbrace{z, z, \dots, z}_{n-k}) \quad \text{and} \quad v^{n+2} = (\underbrace{z, z, \dots, z}_{n-k}, \underbrace{y, y, \dots, y}_k)$$

in  $\mathbb{R}^n$ . We now show that if we take

$$k = \frac{n-2}{2}, \quad y = \frac{n-6}{2}, \quad \text{and} \quad z = \frac{n-2}{2},$$

then  $V = \{v^1, \dots, v^{n+2}\}$  is an  $\ell_1$ -norm equilateral set in  $H_\alpha$ , where  $\alpha = (n-2)^2/2$  and the distance is  $2(n-2)$ .

To verify this we note first that  $b \geq z \geq a \geq y \geq 0$ . For  $i = 1, \dots, n$ , the coefficient sum of  $v^i$  is given by

$$b + (n-2)a = n-2 + \frac{(n-2)(n-4)}{2} = \frac{(n-2)^2}{2}.$$

Similarly the coefficient sum for the vectors  $v^{n+1}$  and  $v^{n+2}$  is equal to

$$(n-k)z + ky = \frac{(n+2)(n-2)}{4} + \frac{(n-2)(n-6)}{4} = \frac{(n-2)^2}{2}.$$

Let  $1 \leq i \neq j \leq n$ . For  $i = 2k-1$  and  $j = 2k$ , the distance between  $v^i$  and  $v^j$  is given by

$$\|v^i - v^j\|_1 = |b - 0| + |0 - b| = 2(n-2),$$

and for all other  $i \neq j$ ,

$$\|v^i - v^j\|_1 = |b - a| + |0 - a| + |a - b| + |a - 0| = 2(b - a) + 2a = 2(n - 2).$$

Also

$$\|v^{n+1} - v^{n+2}\|_1 = k|z - y| + k|y - z| = (n - 2)\left(\frac{n-2}{2} - \frac{n-6}{2}\right) = 2(n - 2).$$

Finally the distance between any of the first  $n$  vectors and the last two is calculated as in either the case of  $v^1$  and  $v^{n+1}$ ,

$$\begin{aligned}\|v^1 - v^{n+1}\|_1 &= |b - y| + |0 - y| + (k - 2)|a - y| + (n - k)|a - z| \\ &= n - 2 + \frac{n-6}{2} + \frac{n+2}{2} \\ &= 2(n - 2),\end{aligned}$$

or, as in the case of  $v^1$  and  $v^{n+2}$ ,

$$\begin{aligned}\|v^1 - v^{n+2}\|_1 &= |b - z| + |0 - z| + (n - k - 2)|a - z| + k|a - y| \\ &= n - 2 + \frac{n-2}{2} + \frac{n-2}{2} \\ &= 2(n - 2).\end{aligned}$$

Thus,  $V$  is an  $\ell_1$ -norm equilateral set in  $H_\alpha$  of size  $n + 2$ . [Table 2](#) shows examples in dimensions  $n = 6, 10$  and  $14$ .

#### 4. Hadamard matrices

In this section, we will give an alternative construction that shows that  $t(\Delta_n) \geq n + 2$  for all  $n = 2^k - 1$  with  $k \geq 2$  using  $\ell_1$ -norm equilateral sets and Hadamard matrices. Recall that an  $n \times n$  matrix  $H = [h_{ij}]$  with entries  $h_{ij} \in \{-1, 1\}$  for all  $i$  and  $j$  is called a *Hadamard matrix* if  $HH^T = nI$ . There exists a simple well-known construction of Hadamard matrices of size  $2^k$ . Define  $H_1 = [1]$  and

$$H_{2^{k+1}} = \begin{bmatrix} H_{2^k} & H_{2^k} \\ H_{2^k} & -H_{2^k} \end{bmatrix}$$

for all  $k \geq 1$ . So,

$$H_2 = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad H_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}, \quad \dots.$$

$n = 6$	$n = 10$	$n = 14$
(4, 0, 1, 1, 1, 1)	(8, 0, 3, 3, 3, 3, 3, 3, 3, 3)	(12, 0, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5)
(0, 4, 1, 1, 1, 1)	(0, 8, 3, 3, 3, 3, 3, 3, 3, 3)	(0, 12, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5)
(1, 1, 4, 0, 1, 1)	(3, 3, 8, 0, 3, 3, 3, 3, 3, 3)	(5, 5, 12, 0, 5, 5, 5, 5, 5, 5, 5, 5)
(1, 1, 0, 4, 1, 1)	(3, 3, 0, 8, 3, 3, 3, 3, 3, 3)	(5, 5, 0, 12, 5, 5, 5, 5, 5, 5, 5, 5)
(1, 1, 1, 1, 4, 0)	(3, 3, 3, 8, 0, 3, 3, 3, 3)	(5, 5, 5, 5, 12, 0, 5, 5, 5, 5, 5, 5)
(1, 1, 1, 1, 0, 4)	(3, 3, 3, 0, 8, 3, 3, 3, 3)	(5, 5, 5, 0, 12, 5, 5, 5, 5, 5, 5)
(2, 2, 2, 2, 0, 0)	(3, 3, 3, 3, 3, 8, 0, 3, 3)	(5, 5, 5, 5, 5, 12, 0, 5, 5, 5, 5, 5)
(0, 0, 2, 2, 2, 2)	(3, 3, 3, 3, 3, 3, 0, 8, 3, 3)	(5, 5, 5, 5, 5, 5, 0, 12, 5, 5, 5, 5, 5)
	(3, 3, 3, 3, 3, 3, 3, 8, 0)	(5, 5, 5, 5, 5, 5, 5, 12, 0, 5, 5, 5)
	(3, 3, 3, 3, 3, 3, 3, 0, 8)	(5, 5, 5, 5, 5, 5, 5, 0, 12, 5, 5, 5)
	(4, 4, 4, 4, 4, 2, 2, 2, 2)	(5, 5, 5, 5, 5, 5, 5, 5, 5, 12, 0, 5, 5)
	(2, 2, 2, 2, 4, 4, 4, 4, 4, 4)	(5, 5, 5, 5, 5, 5, 5, 5, 5, 0, 12, 5, 5)
		(5, 5, 5, 5, 5, 5, 5, 5, 5, 12, 0)
		(5, 5, 5, 5, 5, 5, 5, 5, 5, 0, 12)
		(6, 6, 6, 6, 6, 6, 6, 4, 4, 4, 4, 4)
		(4, 4, 4, 4, 4, 6, 6, 6, 6, 6, 6, 6)

**Table 2.** Equilateral sets of size  $n + 2$ .

Now suppose  $k \geq 2$ . Let  $v^1, \dots, v^{2^k} \in \mathbb{R}^{2^k}$  denote the rows of the Hadamard matrix  $H_{2^k}$ , and define the set

$$V_k = \{v^3\} \cup \{v^i : i = 5, \dots, 2^k\}.$$

Furthermore, let  $W_k = \{w^1, w^2, w^3, w^4\} \in \mathbb{R}^{2^k}$  be given by

$$\begin{aligned} w^1 &= (1, -1, 0, 0, 1, -1, 0, 0, \dots, 1, -1, 0, 0), \\ w^2 &= (-1, 1, 0, 0, -1, 1, 0, 0, \dots, -1, 1, 0, 0), \\ w^3 &= (0, 0, 1, -1, 0, 0, 1, -1, \dots, 0, 0, 1, -1), \\ w^4 &= (0, 0, -1, 1, 0, 0, -1, 1, \dots, 0, 0, -1, 1). \end{aligned}$$

**Proposition 4.1.** *For each  $k \geq 2$ , the set  $V_k \cup W_k$  is an  $\ell_1$ -norm equilateral set of size  $2^k + 1$  in  $H_0 = \{x \in \mathbb{R}^{2^k} : \sum_i x_i = 0\}$ .*

*Proof.* Let  $k \geq 2$ . It is easy to show that each  $u \in V_k \cup W_k$  lies in  $H_0$ . Also note that any two distinct points  $v^i$  and  $v^j$  in  $V_k$  satisfy

$$\|v^i - v^j\|_1 = 2^k,$$

as the rows in  $H_{2^k}$  differ in exactly  $2^{k-1}$  places. The reader can check that

$$\|w^i - w^j\|_1 = 2^k \quad \text{for all } 1 \leq i \neq j \leq 4.$$

So, it remains to show that

$$\|v^i - w^j\|_1 = 2^k \quad \text{for all } v^i \in V_k \text{ and } w^j \in W_k. \quad (4-1)$$

We use induction on  $k$ . Note that if  $k = 2$ , we have that

$$V_2 \cup W_2 = \{(1, 1, -1, -1), (1, -1, 0, 0), (-1, 1, 0, 0), (0, 0, 1, -1), (0, 0, -1, 1)\},$$

which is an  $\ell_1$ -norm equilateral set with distance 4. Now suppose that (4-1) holds for  $k$ . Denote the points in  $V_{k+1}$  by  $\bar{v}^i$  and the points in  $W_{k+1}$  by  $\bar{w}^j$ . Note that for  $j = 1, \dots, 4$ , we have  $\bar{w}^j = (w^j, w^j)$ , where  $w^j \in W_k$ . Also observe that for  $i = 3, 5, \dots, 2^k$ , we have  $\bar{v}^i = (v^i, v^i)$ , and for  $i = 2^k + 1, \dots, 2^{k+1}$ , we have  $\bar{v}^i = (v^{i-2^k}, -v^{i-2^k})$ , where  $v^i \in V_k$ .

So, for  $i = 3, 5, \dots, 2^k$  and  $j = 1, \dots, 4$ , we have that

$$\|\bar{v}^i - \bar{w}^j\|_1 = \sum_{l=1}^{2^{k+1}} |\bar{v}_l^i - \bar{w}_l^j| = 2 \sum_{l=1}^{2^k} |v_l^i - w_l^j| = 2 \cdot 2^k = 2^{k+1}$$

by the induction hypothesis. Also for  $i = 2^k + 1, \dots, 2^{k+1}$  and  $j = 1, \dots, 4$ , we have that

$$\|\bar{v}^i - \bar{w}^j\|_1 = \sum_{l=1}^{2^k} (|v_l^{i-2^k} - w_l^j| + |v_l^{i-2^k} + w_l^j|) = \sum_{l=1}^{2^k} (1 - w_l^j + 1 + w_l^j) = 2^{k+1},$$

as  $v_l^i \in \{-1, 1\}$  and  $-1 \leq w_l^j \leq 1$  for all  $l$ . □

The reader should note that the equilateral set  $V_k \cup W_k$  can be seen as a generalization of the equilateral set  $S$  in (2-2), as  $V_2 \cup W_2 = S - (1, 1, 1, 1)$ . Furthermore, the example in Table 1 with  $n = 8$  is also of this type, if one ignores the point  $(8, 2, 1, 1, 0, 2, 1, 1)$ .

## References

- [Alon and Pudlák 2003] N. Alon and P. Pudlák, “Equilateral sets in  $l_p^n$ ”, *Geom. Funct. Anal.* **13**:3 (2003), 467–482. [MR 2004h:46011](#) [Zbl 1034.46015](#)
- [Bandelt et al. 1998] H.-J. Bandelt, V. Chepoi, and M. Laurent, “Embedding into rectilinear spaces”, *Discrete Comput. Geom.* **19**:4 (1998), 595–604. [MR 99d:51017](#) [Zbl 0973.51012](#)
- [Bezdek 2010] K. Bezdek, *Classical topics in discrete geometry*, Springer, New York, 2010. [MR 2011j:52014](#) [Zbl 1207.52001](#)
- [Danzer and Grünbaum 1962] L. Danzer and B. Grünbaum, “Über zwei Probleme bezüglich konvexer Körper von P. Erdős und von V. L. Klee”, *Math. Z.* **79** (1962), 95–99. [MR 25 #1488](#) [Zbl 0188.27602](#)
- [Guy 1983] R. K. Guy, “An olla-podrida of open problems, often oddly posed”, *Amer. Math. Monthly* **90**:3 (1983), 196–200. [MR 1540158](#)
- [Koolen et al. 2000] J. Koolen, M. Laurent, and A. Schrijver, “Equilateral dimension of the rectilinear space”, *Des. Codes Cryptogr.* **21**:1-3 (2000), 149–164. [MR 2001j:52013](#) [Zbl 0970.51016](#)

- [Lemmens 2007] B. Lemmens, “Variations of a combinatorial problem on finite sets”, *Elem. Math.* **62**:2 (2007), 59–67. [MR 2008e:05017](#) [Zbl 1135.05008](#)
- [Petty 1971] C. M. Petty, “Equilateral sets in Minkowski spaces”, *Proc. Amer. Math. Soc.* **29** (1971), 369–374. [MR 43 #1051](#) [Zbl 0214.20801](#)
- [Soltan 1975] P. S. Soltan, “Analogues of regular simplexes in normed spaces”, *Dokl. Akad. Nauk SSSR* **222**:6 (1975), 1303–1305. In Russian; translated in *Soviet Math. Dokl.* **16**:3 (1975), 787–789. [MR 52 #4127](#) [Zbl 0338.46025](#)
- [Swanepoel 2004a] K. J. Swanepoel, “Equilateral sets in finite-dimensional normed spaces”, pp. 195–237 in *Seminar of mathematical analysis*, edited by D. Girela Álvarez et al., Colección Abierta **71**, Univ. Sevilla Secr. Publ., Seville, 2004. [MR 2005j:46009](#) [Zbl 1071.52008](#)
- [Swanepoel 2004b] K. J. Swanepoel, “A problem of Kusner on equilateral sets”, *Arch. Math. (Basel)* **83**:2 (2004), 164–170. [MR 2005i:52024](#) [Zbl 1062.52017](#)

Received: 2013-12-17 Accepted: 2014-02-23

[b.lemmens@kent.ac.uk](mailto:b.lemmens@kent.ac.uk)

*School of Mathematics, Statistics & Actuarial Science,  
University of Kent, Cornwallis Building, Canterbury,  
CT2 7NF, United Kingdom*

[cmp37@kent.ac.uk](mailto:cmp37@kent.ac.uk)

*School of Mathematics, Statistics & Actuarial Science,  
University of Kent, Cornwallis Building, Canterbury,  
CT2 7NF, United Kingdom*

## EDITORS

### MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

### BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	La Trobe University, Australia P.Cerone@latrobe.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tobrie1@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyj@potsdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemonns	Wake Forest University, USA plemonns@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA jgalian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	UMass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew@dms.蒙特利尔.ca	Józeph H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sgupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rw@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakil@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnson@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

### PRODUCTION

Silvio Levy, Scientific Editor

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2015 is US \$140/year for the electronic version, and \$190/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY



nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

# involve

2015

vol. 8

no. 3

Colorability and determinants of $T(m, n, r, s)$ twisted torus knots for $n \equiv \pm 1 \pmod{m}$	361
MATT DELONG, MATTHEW RUSSELL AND JONATHAN SCHROCK	
Parameter identification and sensitivity analysis to a thermal diffusivity inverse problem	385
BRIAN LEVENTHAL, XIAOJING FU, KATHLEEN FOWLER AND OWEN ESLINGER	
A mathematical model for the emergence of HIV drug resistance during periodic bang-bang type antiretroviral treatment	401
NICOLETA TARFULEA AND PAUL READ	
An extension of Young's segregation game	421
MICHAEL BORCHERT, MARK BUREK, RICK GILLMAN AND SPENCER ROACH	
Embedding groups into distributive subsets of the monoid of binary operations	433
GREGORY MEZERA	
Persistence: a digit problem	439
STEPHANIE PEREZ AND ROBERT STYER	
A new partial ordering of knots	447
ARAZELLE MENDOZA, TARA SARGENT, JOHN TRAVIS SHRONTZ AND PAUL DRUBE	
Two-parameter taxicab trigonometric functions	467
KELLY DELP AND MICHAEL FILIPSKI	
${}_3F_2$ -hypergeometric functions and supersingular elliptic curves	481
SARAH PITMAN	
A contribution to the connections between Fibonacci numbers and matrix theory	491
MIRIAM FARBER AND ABRAHAM BERMAN	
Stick numbers in the simple hexagonal lattice	503
RYAN BAILEY, HANS CHAUMONT, MELANIE DENNIS, JENNIFER MCLOUD-MANN, ELISE McMAHON, SARA MELVIN AND GEOFFREY SCHUETTE	
On the number of pairwise touching simplices	513
BAS LEMMENS AND CHRISTOPHER PARSONS	
The zipper foldings of the diamond	521
ERIN W. CHAMBERS, DI FANG, KYLE A. SYKES, CYNTHIA M. TRAUB AND PHILIP TRETENERO	
On distance labelings of amalgamations and injective labelings of general graphs	535
NATHANIEL KARST, JESSICA OEHRELEIN, DENISE SAKAI TROXELL AND JUNJIE ZHU	