

**CORRECTION: INVARIANCE PRINCIPLES AND ASYMPTOTIC
CONSTANCY OF SOLUTIONS OF PRECOMPACT
FUNCTIONAL DIFFERENTIAL EQUATIONS**

(Tôhoku Math. J. 42 (1990), 217–229)

WILLIAM E. HORNOR

(Received February 25, 1991)

In the proof of Theorem 5.1, with $t^* > 0$, the author should have noted that

$$[0, t^*] \subset [t^{**} - NQ, t^{**} - NP],$$

and should have set

$$F_N = \{t^{**} - NQ, \dots, t^{**} - ((N-k)Q + kP), \dots, t^{**} - NP\},$$

and

$$\hat{F}_N = (F_N \cap [0, t^*]) \cup \{0, t^*\}.$$

By hypothesis (i) of the theorem, we have $V[x(t_0, \psi, g)(t)] = c$ for each $t \in \hat{F}_N$ with the possible exceptions $t=0$ and $t=t^*$. We then conclude that $V[x(t_0, \psi, g)(t)] = c$ for every $t > 0$.

DEPARTMENT OF MATHEMATICS
THE UNIVERSITY OF SOUTHERN MISSISSIPPI
SOUTHERN STATION, BOX 5045
HATTIESBURG, MISSISSIPPI 39406-5045
U.S.A.

