

192. No Free Inverse Semigroup is Orderable^{*)}

By Tôru SAITÔ

Tokyo Gakugei University

(Comm. by Kenjiro SHODA, M. J. A., Dec. 12, 1974)

A semigroup S is said to be orderable if S admits a simple order to make S a simply ordered semigroup. It is known that every free group is orderable ([1] or [4]). On the other hand, it is shown in [3] that a free commutative idempotent semigroup with r free generators is orderable if and only if $r \leq 2$. In this paper we consider the orderability of free inverse semigroups. Since it is easily shown that a free inverse semigroup with r free generators contains a free commutative idempotent subsemigroup with r free generators, it follows from the above result that a free inverse semigroup with a set of free generators consisting of more than three elements is not orderable. But we have the following stronger result.

Theorem. *No free inverse semigroup is orderable.*

Proof. By way of contradiction we assume that S is a free inverse semigroup which is at the same time an ordered semigroup. We denote by \leq the natural partial order on the inverse semigroup S . Let F be a set of free generators of S and let $x \in F$. Then, by [5] Corollary 2.5, the inverse subsemigroup $\langle x \rangle$ generated by x is a free inverse semigroup with a free generator x . Also in S the set E of idempotents of S forms a semilattice with respect to the partial order \leq and, by [6] Theorem 3, the semilattice E is a tree semilattice. Now we have

$$x^2x^{-2} \leq xx^{-1}, \quad xx^{-2}x \leq xx^{-1}$$

and so x^2x^{-2} and $xx^{-2}x$ are comparable. If $x^2x^{-2} \leq xx^{-2}x$, then $x^2x^{-2} \leq xx^{-2}x \leq x^{-1}x$, which contradicts [5] Corollary 2.4. Next we suppose $xx^{-2}x \leq x^2x^{-2}$. Then $xx^{-2}x = (x^2x^{-2})(xx^{-2}x) = x^2x^{-3}x$ and so $x^2 = x(xx^{-2}x)x = x^3x^{-3}x^2$. Hence

$$x^{-2}x^2 = x^{-2}x^3x^{-3}x^2 = (x^{-2}x^2)(xx^{-1})(x^{-2}x^2) \leq xx^{-1},$$

which contradicts again [5] Corollary 2.4.

References

- [1] G. Birkhoff: Review of "Everett, C. J., and Ulam, S., Ordered groups." Math. Rev., 7, 4 (1946).
- [2] A. H. Clifford and G. B. Preston: The algebraic theory of semigroups, Vol. I. Amer. Math. Soc. (1961); Vol. II. Amer. Math. Soc. (1967).

^{*)} Dedicated to Professor Kiiti Morita on his 60th birthday.

- [3] Е. Я. Габович: Об упорядочиваемости свободных полугрупп некоторых многообразий. Матем. Заметки. (to appear).
- [4] К. Iwasawa: On linearly ordered groups. *J. Math. Soc. Japan*, **1**, 1–9 (1948).
- [5] N. R. Reilly: Free generators in free inverse semigroups. *Bull. Austral. Math. Soc.*, **7**, 407–424 (1972).
- [6] T. Saitô: Ordered idempotent semigroups. *J. Math. Soc. Japan*, **14**, 150–169 (1962).
- [7] H. E. Scheiblich: Free inverse semigroups. *Proc. Amer. Math. Soc.*, **38**, 1–7 (1973).