

72. On Two Classes of Subalgebras of $L^1(G)$

By Leonard Y. H. YAP

Department of Mathematics University of Singapore

(Comm. by Kinjirô KUNUGI, M. J. A., May 12, 1972)

1. Introduction. Let G and \hat{G} be two locally compact Abelian groups in Pontrjagin duality. The Fourier transform of a function $f \in L^1(G)$ will be denoted by \hat{f} . For $1 \leq p < \infty$, define

$$A^p(G) = \{f \in L^1(G) : \hat{f} \in L^p(\hat{G})\}, \quad B^p(G) = L^1(G) \cap L^p(G).$$

The space $A^p(G)$ is a Banach algebra with respect to the norm $\|\cdot\|_{A^p(G)}$ defined by $\|f\|_{A^p(G)} = \|f\|_1 + \|\hat{f}\|_p$ and the usual convolution product. The Banach algebra $A^p(G)$ have been studied by Larsen-Liu-Wang [8], Lai [5]–[7], Martin-Yap [9], and others. The space $B^p(G)$ is a Banach algebra with respect to the norm $\|\cdot\|_{B^p(G)}$ defined by $\|f\|_{B^p(G)} = \|f\|_1 + \|f\|_p$ and the usual convolution product. The Banach algebras $B^p(G)$ have been studied by Warner [12], Yap [15], and others. The purpose of this paper is to extend some of the results on $A^p(G)$ and $B^p(G)$ to the spaces

$$A(p, q)(G) = \{f \in L^1(G) : \hat{f} \in L(p, q)(\hat{G})\}$$

and

$$B(p, q)(G) = L^1(G) \cap L(p, q)(G)$$

respectively (see next section for the definition of $L(p, q)(G)$ and some relevant facts about these spaces). In Section 2 we identify the maximal ideal spaces of the algebras $A(p, q)(G)$ and $B(p, q)(G)$, show that they satisfy Ditkin's condition and that the Shilov-Wiener Tauberian theorem holds for these algebras. In Section 3 we prove non-factorization theorems for these algebras.

2. Tauberian theorem for $A(p, q)(G)$ and $B(p, q)(G)$. For the convenience of the reader, we now review briefly what we need from the theory of $L(p, q)$ spaces.

Definition 2.1. Let f be a measurable function defined on (G, λ) , where λ is the Haar measure of G . For $y \geq 0$, we define

$$m(f, y) = \lambda\{x \in G : |f(x)| > y\}.$$

For $x \geq 0$, we define

$$\begin{aligned} f^*(x) &= \inf \{y : y > 0 \text{ and } m(f, y) \leq x\} \\ &= \sup \{y : y > 0 \text{ and } m(f, y) > x\}, \end{aligned}$$

with the conventions $\inf \phi = \infty$ and $\sup \phi = 0$. For $x > 0$, we define

$$f^{**}(x) = x^{-1} \int_0^x f^*(t) dt.$$

We also define

$$\|f\|_{(p,q)}^* = \left\{ \int_0^\infty [x^{1/p} f^*(x)]^q \frac{dx}{x} \right\}^{1/q}, \quad (0 < p < \infty, 0 < q < \infty)$$

$$\|f\|_{(p,\infty)}^* = \sup_{x>0} x^{1/p} f^*(x) \quad (0 < p < \infty)$$

$$L(p, q)(G) = \{f : \|f\|_{(p,q)}^* < \infty\}.$$

It is quite easy to see that we have

$$\int_0^\infty f^*(x)^p dx = \int_G |f(x)|^p d\lambda(x)$$

and hence $L^p(G) = L(p, p)(G)$, $A^p(G) = A(p, p)(G)$, $B^p(G) = B(p, p)(G)$.

If we replace $f^*(x)$ by $f^{**}(x)$ in the definition of $\|f\|_{(p,q)}^*$, the resulting number will be denoted by $\|f\|_{(p,q)}$. For $1 < p < \infty$, $1 \leq q \leq \infty$, it is known that

(i) $\|f\|_{(p,q)}^* \leq \|f\|_{(p,q)} \leq p/(p-1) \cdot \|f\|_{(p,q)}^*$ (see the proof of (3.2) in [13]),

(ii) $(L(p, q), \|\cdot\|_{(p,q)})$ is a Banach space. (see [4, (2.6)], [10, (2.1)].)

Thus we can endow $A(p, q)(G)$ and $B(p, q)(G)$ ($1 < p < \infty$, $1 \leq q \leq \infty$) with the norms

$$\|f\|_{A(p,q)} = \|f\|_1 + \|\hat{f}\|_{(p,q)}, \quad \|f\|_{B(p,q)} = \|f\|_1 + \|f\|_{(p,q)}$$

respectively.

We now single out the following fact for easy reference.

Lemma 2.2. *Let $1 < p < \infty$, $1 \leq q \leq \infty$. Let $\{f_n\}$ be a sequence in $L(p, q)(G)$ and $\|f_n - f\|_{(p,q)} \rightarrow 0$, where $f \in L(p, q)(G)$. Then $\{f_n\}$ has a subsequence which converges pointwise almost everywhere to f .*

Proof. See the proof of (2.3) in Hunt [4, p. 258] and (2.1(i)) above.

We will prove the main result in this section via the concept of Segal algebra whose definition we now recall. A subalgebra $S(G)$ of $L^1(G)$ is called a *Segal algebra* if:

- (S-1) $S(G)$ is dense in $L^1(G)$ in the L^1 -norm topology and if $f \in S(G)$ then $f_a \in S(G)$, where $f_a(x) = f(a^{-1}x)$;
- (S-2) $S(G)$ is a Banach algebra under some norm $\|\cdot\|_S$ which also satisfies $\|f\|_S = \|f_a\|_S$ for all $f \in S(G)$, $a \in G$ (multiplication in $S(G)$ is the usual convolution);
- (S-3) if $f \in (G)$, then for any $\epsilon > 0$ there exists a neighborhood U of the identity element of G such that $\|f_y - f\|_S < \epsilon$ for all $y \in U$.

Proposition 2.3. *For $1 < p < \infty$ and $1 \leq q < \infty$, the space $A(p, q)$ is a Segal algebra with respect to the norm $\|\cdot\|_{A(p,q)}$.*

Proof. Clearly $A(p, q)$ is a subalgebra of L^1 and $f_a \in A(p, q)$ whenever $f \in A(p, q)$, $a \in G$. Since $D = \{f \in L^1 : \hat{f} \text{ has compact support}\}$ is dense in L^1 (see [11, 2.6.6]) and $D \subset A(p, q)$, $A(p, q)$ is dense in L^1 . Thus condition (S-1) is satisfied.

That $A(p, q)$ is a Banach algebra with respect to the norm $\|\cdot\|_{A(p,q)}$ can be proved as in [8, Theorems 1 and 3], using Lemma (2.2) above.

It is clear that $\|f\|_{A(p,q)} = \|f_a\|_{A(p,q)}$ for all $f \in A(p,q)$, $a \in G$. Thus condition (S-2) is fulfilled.

Next we check that $A(p,q)$ satisfies condition (S-3). Let $0 \neq f \in A(p,q)$ and let $\varepsilon > 0$. First we choose a neighborhood U of the identity element e of G such that $\|f_y - f\|_1 < \varepsilon/2$ for all $y \in U$. Define $\varepsilon' = \varepsilon(p-1)/p$. Choose a continuous function ϕ on \hat{G} having compact support such that $\|\phi - \hat{f}\|_{(p,q)}^* < \varepsilon'/8$ (see [13, (4.2)]). Let K denote the support of ϕ , and let $K' = \hat{G} \setminus K$. It follows that

$$(1) \quad \|\hat{f}\chi_{K'}\|_{(p,q)}^* < \varepsilon'/8.$$

Now define

$$N(K, \varepsilon') = \{y \in G : |(y, \gamma) - 1| < \varepsilon'/4 \|\hat{f}\|_{(p,q)}^* \text{ for all } \gamma \in K\}.$$

Then $N(K, \varepsilon')$ is a neighborhood of e in G . We now choose a symmetric neighborhood W of e such that $W \subset U \cap N(K, \varepsilon')$. It follows that

- (i) for $y \in W$ and $\gamma \in K$ we have

$$|\hat{f}_y(\gamma) - \hat{f}(\gamma)| = |(y^{-1}, \gamma) - 1| \cdot |\hat{f}(\gamma)| < \varepsilon'(4 \|\hat{f}\|_{(p,q)}^*)^{-1} |\hat{f}(\gamma)|,$$
 and hence $\|(\hat{f}_y - \hat{f})\chi_K\|_{(p,q)}^* < \varepsilon'/4$;
- (ii) for $y \in W$ and $\gamma \in K'$ we have $|\hat{f}_y(\gamma) - \hat{f}(\gamma)| \leq 2|\hat{f}(\gamma)|$. It follows from (1) that $\|(\hat{f}_y - \hat{f})\chi_{K'}\|_{(p,q)}^* < \varepsilon'/4$.

Thus for $y \in W$ we have $\|\hat{f}_y - \hat{f}\|_{(p,q)}^* < \varepsilon'/2$, and hence $\|f_y - f\|_{A(p,q)} < \varepsilon$ for all $y \in W$.

Proposition 2.4. *For $1 < p < \infty$ and $1 \leq q < \infty$, the space $B(p,q)$ is a Segal algebra with respect to some norm which is equivalent to the norm $\|\cdot\|_{B(p,q)}$.*

Proof. Blozinski [1, (2.9)] shows that if $f \in L^1$ and $g \in L(p,q)$ then $\|f * g\|_{(p,q)} \leq C(p,q) \|f\|_1 \cdot \|g\|_{(p,q)}$, where $C(p,q)$ is a constant depending only on p, q . We assume with no loss of generality that $C(p,q) \geq 1$. It follows that if $f, g \in B(p,q)$ then $\|f * g\|_{B(p,q)} \leq C(p,q) \|f\|_{B(p,q)} \cdot \|g\|_{B(p,q)}$. Thus $\|f\|_{B(p,q)} = C(p,q) \|f\|_{B(p,q)}$ defines a norm in $B(p,q)$ under which $B(p,q)$ is a Banach algebra. Since $B(p,q)$ contains all the continuous functions with compact supports, $B(p,q)$ is dense in L^1 . Thus conditions (S-1) and (S-2) are satisfied.

We now prove that $B(p,q)$ satisfies condition (S-3). Let $0 \neq f \in B(p,q)$ and let $\varepsilon > 0$. First choose a continuous function ϕ with compact support such that $\|\phi - f\|_{(p,q)}^* < \varepsilon'/4$, where $\varepsilon' = \varepsilon(p-1)/pC(p,q)$. Let $K = \text{support of } \phi$. By the uniform continuity of ϕ , there is a neighborhood V of the identity element e in G such that

$$\|\phi - \phi_x\|_\infty < \frac{\varepsilon'}{4} (q/p)^{1/q} (2\lambda(K))^{-1/p}$$

for all $x \in V$. It follows that $\|\phi - \phi_x\|_{(p,q)}^* < \varepsilon'/4$ for all $x \in V$. Next choose a neighborhood W of e such that $W \subset V$ and $\|f - f_x\|_1 < \varepsilon/4C(p,q)$ for all $x \in W$. Thus for $x \in W$ we have

$$\begin{aligned} \|f - f_x\|_{B(p,q)} &= C(p, q) \cdot \|f - f_x\|_1 + C(p, q) \cdot \|f - f_x\|_{(p,q)} \\ &< \varepsilon/4 + C(p, q) [\|f - \phi\|_{(p,q)} + \|\phi - \phi_x\|_{(p,q)} + \|\phi_x - f_x\|_{(p,q)}] \\ &< \varepsilon/4 + C(p, q) \frac{p}{p-1} (\varepsilon'/4 + \varepsilon'/4 + \varepsilon'/4) = \varepsilon. \end{aligned}$$

Theorem 2.5. *Let $S(G) = A(p, q)(G)$ or $B(p, q)(G)$. Then*

- (i) *the maximal ideal space of $S(G)$ can be identified with the dual group \hat{G} of G ;*
- (ii) *the algebra $S(G)$ satisfies Ditkin's condition;*
- (iii) *the Shilov-Wiener Tauberian theorem holds in $S(G)$.*

Proof. Immediate from Propositions (2.3) and (2.4) and the fact that every Segal algebra has properties (i)–(iii) (Yap [16]).

3. Non-factorization in $A(p, q)(G)$ and $B(p, q)(G)$. We recall that an algebra A is said to have the *factorization property* if $A = A \cdot A$, where $A \cdot A = \{xy : x, y \in A\}$. We use A^2 to denote the ideal in A generated by $A \cdot A$. The group algebra $L^1(G)$ is known to have the factorization property (Cohen [2]), but in general $A^p(G)$ and $B^p(G)$ do not satisfy this property (Martin-Yap [9] and Yap [15]). In this section we extend these non-factorization theorems to the algebras $A(p, q)(G)$ and $B(p, q)(G)$.

Lemma 3.1. $A(p, q)^2 \subset A(p/2, q/2)$.

Proof. It suffices to show that if $f, g \in A(p, q)$ then $f * g \in A(p/2, q/2)$. First we define $\alpha = 2(p + q)/q$. Thus $|\hat{f}|^{p/\alpha}, |\hat{g}|^{p/\alpha} \in L(\alpha, \alpha q/p)$ and by O'Neil [10, 3.4] we see that $|\hat{f}\hat{g}|^{p/\alpha} \in L(r, s)$, where

$$1/r = 1/\alpha + 1/\alpha, \quad 1/s = p/\alpha q + p/\alpha q.$$

It follows that $\widehat{f * g} = \hat{f}\hat{g} \in L(p/2, q/2)$, and hence $f * g \in A(p/2, q/2)$.

Theorem 3.2. If G is non-discrete, $1 < p < \infty$, $1 \leq q < \infty$, then $A(p, q)(G)^2 \neq A(p, q)(G)$.

Proof. Suppose $A(p, q)^2 = A(p, q)$, then by Lemma (3.1) we would have $A(p, q) \subset A(p/2^n, q/2^n)$ for $n = 1, 2, 3, \dots$. We will show that this leads to a contradiction. Since G is non-discrete, \hat{G} is non-compact, and we may choose a symmetric neighborhood U of the identity in \hat{G} whose closure \bar{U} is compact, and a sequence $\gamma_1, \gamma_2, \gamma_3, \dots$ in \hat{G} such that

$$\gamma_i U^2 \cap \gamma_j U^2 = \emptyset \quad (i \neq j)$$

Now let N be a positive integer such that $p < 2^N$. Define

$$\alpha = 2^N/p, a_n = n^{-\alpha} \quad (n = 1, 2, 3, \dots)$$

$$g = \chi_U, h = \sum_{k=1}^{\infty} a_k \chi_{\gamma_k U^2}.$$

Thus $g, h \in L^2(\hat{G})$ and so by Rudin [11, Theorem 1.6.3] there is a function $f \in L^1(G)$ such that $\hat{f} = g * h$. It follows that $\hat{f}(\gamma) = g * h(\gamma) = a_k \rho(U)$ for $\gamma \in \gamma_k U$, where ρ denotes the Harr measure of \hat{G} . Direct computations (similar to those in [14, p. 138]) show that $\hat{f} \in L(p, q)$, but $\hat{f} \notin L(p/2^N, q/2^N)$. Hence $f \in A(p, q)$, but $f \notin A(p/2^N, q/2^N)$.

Lemma 3.3. *If $f \in L(p_1, s) \cap L(p_2, s)$, then $f \in L(r, s)$ for all r such*

that $p_1 < r < p_2$.

Proof. Define $\beta = (1/r - 1/p_2)(1/p_1 - 1/p_2)^{-1}$, and note that

$$\begin{aligned} \|f\|_{(r,s)}^{*s} &= \int_0^\infty f^*(x)^s x^{s/r-1} dx \\ &= \int_0^\infty [f^*(x)^\beta x^{\beta(s/p_1-1)}] \cdot [f^*(x)^{(1-\beta)s} x^{(1-\beta)(s/p_2-1)}] dx \\ &\leq \|f\|_{(p_1,s)}^{*s\beta} \cdot \|f\|_{(p_2,s)}^{*(1-\beta)s} \quad (\text{by Hölder's inequality}). \end{aligned}$$

Theorem 3.4. If G is non-discrete and $1 < p < \infty$, $1 \leq q < \infty$, then $B(p, q)(G)^2 \neq B(p, q)(G)$.

Proof. Let $f, g \in B(p, q)$. Since $L^1 = L(1, 1)$, and $L(1, 1) \subset L(1, q)$ (by [13, (3.3)]), it follows that $f, g \in L(1, q)$. Define $r = 2p/(1+p)$. Clearly $1 < r < p$, and so $f, g \in L(r, q)$ by Lemma (3.3). By [13, (3.5)] we have $f * g \in L(p, q/2)$. Thus $B(p, q)^2 \subset B(p, q/2)$. But $B(p, q/2)$ is a proper subset of $B(p, q)$ (see the proof of Case I of Theorem (2.7) in Yap [14]).

Remark 3.5. Theorem (3.4) is valid for all (non-discrete) locally compact unimodular groups and the proof is the same.

Conjecture. For a Segal algebra $S(G)$, $S(G)^2 \neq S(G)$ if $S(G) \neq L^1(G)$.

References

- [1] A. P. Blozinski: On a convolution theorem for $L(p, p)$ spaces (to appear).
- [2] P. J. Cohen: Factorization in group algebras. *Duke Math. J.*, **26**, 199–205 (1959).
- [3] E. Hewitt and K. A. Ross: *Abstract Harmonic Analysis. I.* Berlin (1963).
- [4] R. A. Hunt: On $L(p, p)$ spaces, *l'Enseigne. Math.*, **12**, 249–276 (1966).
- [5] H. C. Lai: On some properties of $A^p(G)$ -algebras. *Proc. Japan Acad.*, **45**, 572–575 (1969).
- [6] —: On the category of $L^1(G)$, $L^p(G)$ in $A^q(G)$. *ibid.*, **45**, 577–581 (1969).
- [7] —: Remark on the $A^p(G)$ -algebras. *ibid.*, **46**, 58–63 (1970).
- [8] R. Larsen, T. S. Liu, and J. K. Wang: On functions with Fourier transforms in L^p . *Michigan Math. J.*, **11**, 369–378 (1964).
- [9] J. C. Martin and L. Y. H. Yap: The algebra of functions with Fourier transforms in L^p . *Proc. Amer. Math. Soc.*, **24**, 217–219 (1970).
- [10] R. O'Neil: Convolution operators and $L(p, q)$ spaces. *Duke Math. J.*, **30**, 129–142 (1963).
- [11] W. Rudin: *Fourier Analysis on Groups.* New York (1962).
- [12] C. R. Warner: Closed ideals in the group algebra $L^1(G) \cap L^2(G)$. *Trans. Amer. Math. Soc.*, **121**, 408–423 (1966).
- [13] L. Y. H. Yap: Some remarks on convolution operators and $L(p, q)$ spaces. *Duke Math. J.*, **36**, 647–658 (1969).
- [14] —: On the impossibility of representing certain functions by convolutions. *Math. Scand.*, **26**, 132–140 (1970).
- [15] —: Ideals in subalgebras of the group algebras. *Studia Math.*, **35**, 165–175 (1970).
- [16] —: Every Segal algebra satisfies Ditkin's condition. *Studia Math.*, **40**, 235–237 (1971).