

142. The Number of Tree Semilattices

By Tôru SAITÔ

Tokyo Gakugei University

(Comm. by Kinjirô KUNUGI, M.J.A., Nov. 12, 1963)

By a *tree semilattice* we mean a semilattice T which satisfies the following condition:

if $a \leq a'$, $b \leq b'$, a and b are non-comparable, then a' and b' are non-comparable.

This semilattice plays an important role in the theory of ordered semigroups ([1], [2]).

For a positive integer n , we denote by $T(n)$ the number of non-isomorphic tree semilattices of order n . In this note we give a method of calculating the number $T(n)$.

Let T be a tree semilattice of order n and let 0 be the zero element of T . We denote $T \setminus 0$ by T' . Clearly $T(1)=1$. If $n > 1$, then T' is decomposed into disjoint tree semilattices, say, i tree semilattices of order 1, j tree semilattices of order 2, k tree semilattices of order 3 and so on. Evidently

$$n - 1 = i + 2j + 3k + \dots$$

Now there is 1 way of selecting i tree semilattices of order 1, ${}_{T(2)}H_j$ ways of selecting j tree semilattices of order 2, ${}_{T(3)}H_k$ ways of selecting k tree semilattices of order 3 and so on. Thus we have the following

Theorem. $T(n)$ satisfies the formal relation

$$\begin{aligned} & (1 + x + x^2 + \dots)(1 + {}_{T(2)}H_1x^2 + {}_{T(2)}H_2x^4 + {}_{T(2)}H_3x^6 + \dots) \\ & (1 + {}_{T(3)}H_1x^3 + {}_{T(3)}H_2x^6 + {}_{T(3)}H_3x^9 + \dots) \\ & = T(1) + T(2)x + T(3)x^2 + T(4)x^3 + T(5)x^4 + \dots \end{aligned}$$

Comparing the corresponding coefficients, we can calculate $T(n)$ recursively. We list the first 10 numbers of $T(n)$.

n	1	2	3	4	5	6	7	8	9	10
$T(n)$	1	1	2	4	9	20	48	115	286	719

Appendix 1. Tree semilattice in the above sense were called flowing semilattice by Tamura [3].

2. In [4], Kimura gave a formula to calculate the number of orderable semilattices (in his sense). Reminding of Theorem 14 in [1], this formula can be obtained by a similar reasoning as in the present paper.

References

- [1] T. Saitô: Ordered idempotent semigroups, *J. Math. Soc. Japan*, **14**, 150-169 (1962).
- [2] —: Ordered completely regular semigroups, to appear in *Pacific J. Math.*
- [3] T. Tamura: On a special semilattice with a minimal condition, *J. Gakugei, Tokushima Univ.*, **7**, 9-17 (1956).
- [4] N. Kimura: Total orderings on a semilattice, *Proc. Japan Acad.*, **34**, 124-125 (1958).