

39. A Generalization of Local Class Field Theory by Using K -groups. I

By Kazuya KATO^{*)}

Department of Mathematics, Faculty of Science, University of Tokyo

(Communicated by Kunihiko KODAIRA, M. J. A., Sept. 12, 1977)

§0. Introduction. This note is a summary of our recent results on a generalization of local class field theory. Details will be published elsewhere.

Let F be a field which is complete with respect to a discrete valuation and with finite residue field. Let K be a field which is complete with respect to a discrete valuation and with residue field F . In this Part I, we shall study abelian extensions of K . The case in which F is a function field of one variable over a finite field and a generalization of our results will be studied in Part II ([1]).

§1. In Part I, let F denote a field which is complete with respect to a discrete valuation and with finite residue field, and let K denote a field which is complete with respect to a discrete valuation and with residue field F , and let K^{ab} denote the maximum abelian extension of K .

Theorem 1. (1) *There exists a canonical homomorphism*

$$\Phi: K_2(K) \longrightarrow \text{Gal}(K^{ab}/K)$$

having the following property: For each finite abelian extension L of K , Φ induces an isomorphism

$$K_2(K)/N_{L/K}K_2(K) \cong \text{Gal}(L/K),$$

where $N_{L/K}$ denotes the norm map in K_2 -theory.

(2) $L \mapsto N_{L/K}K_2(L)$ is a bijection from the set of all finite abelian extensions of K in a fixed algebraic closure of K to the set of all open subgroups of finite indices of $K_2(K)$ with respect to the topology defined later in §4.

This is closely connected with the following result on the Brauer group of K .

Theorem 2. *There exists a canonical isomorphism*

$$\Psi: \text{Br}(K) \xrightarrow{\cong} \text{Hom}_c(K^*, \mathbf{Q}/\mathbf{Z})_{\text{tor}}$$

having the following property, where K^ denotes the multiplicative group of K and $\text{Hom}_c(K^*, \mathbf{Q}/\mathbf{Z})_{\text{tor}}$ denotes the torsion part of the group of all continuous homomorphism $K^* \rightarrow \mathbf{Q}/\mathbf{Z}$ with respect to the topology*

^{*)} The author wishes to express his sincere gratitude to Professor Y. Ihara for his stimulation and for his suggestion to study class field theory of those type of field discussed in Part II §1.

defined later (§ 4): For each central simple algebra A over K , the kernel of $\Psi(\{A\})$ is $Nrd_{A/K}A^*$ where Nrd denotes the reduced norm.

§ 2. The definitions of the p -primary parts of Φ and Ψ in the mixed characteristic case. (Cf. Part II for the case $ch(K)=p$.)

Suppose that $ch(F)=p>0$ and $ch(K)=0$. (ch denotes the characteristic of a field.) Let K_{nr} be the maximum unramified extension of K and F_s be the residue field of K_{nr} , so that F_s is the separable closure of F . Put $G=\text{Gal}(K_{nr}/K)\cong\text{Gal}(F_s/F)$. Let r be any natural number. Consider the following diagram of G -modules:

$$(1) \quad \begin{array}{ccc} K_2(K_{nr})/K_2(K_{nr})^{p^r} & \xrightarrow{g} & H^2(K_{nr}, \mu_{p^r} \otimes \mu_{p^r}) \\ \downarrow t & & \\ F_s^*/F_s^{*p^r} & & \end{array}$$

Here, we use the notation in [3] for the Galois cohomology group, t denotes the tame symbol, g denotes the Galois symbol (cf. [4]), and μ_{p^r} denotes the group of all p^r -th roots of 1. By Proposition 1 (1) below, g is an isomorphism. So, (1) induces a homomorphism

$$(2) \quad H^1(G, H^2(K_{nr}, \mu_{p^r} \otimes \mu_{p^r})) \longrightarrow H^1(G, F_s^*/F_s^{*p^r}).$$

On the other hand, $H^1(G, H^2(K_{nr}, \mu_{p^r} \otimes \mu_{p^r})) \cong H^3(K, \mu_{p^r} \otimes \mu_{p^r})$ by Proposition 1 (2) below, and

$$H^1(G, F_s^*/F_s^{*p^r}) \cong \frac{1}{p^r} \mathbf{Z}/\mathbf{Z}$$

by ordinary local class field theory. So, (2) induces a homomorphism

$$(3) \quad H^3(K, \mu_{p^r} \otimes \mu_{p^r}) \longrightarrow \frac{1}{p^r} \mathbf{Z}/\mathbf{Z},$$

which is in fact an isomorphism.

Now, (3) induces two homomorphisms

$$(4) \quad \begin{array}{ccc} K_2(K)/K_2(K)^{p^r} \otimes \text{Hom}_c(\text{Gal } K^{ab}/K), \mathbf{Z}/p^r & & \\ \xrightarrow{b} & H^2(K, \mu_{p^r} \otimes \mu_{p^r}) \otimes H^1(K, \mathbf{Z}/p^r) & \\ \xrightarrow{c} & H^3(K, \mu_{p^r} \otimes \mu_{p^r}) \longrightarrow \frac{1}{p^r} \mathbf{Z}/\mathbf{Z}, & \end{array}$$

and

$$(5) \quad \begin{array}{ccc} K^*/K^{*p^r} \otimes \text{Br}(K)_{p^r} & & \\ \xrightarrow{b'} & H^1(K, \mu_{p^r}) \otimes H^2(K, \mu_{p^r}) & \\ \xrightarrow{c'} & H^3(K, \mu_{p^r} \otimes \mu_{p^r}) \longrightarrow \frac{1}{p^r} \mathbf{Z}/\mathbf{Z}, & \end{array}$$

where:

Hom_c is the group of continuous homomorphisms,

b is the tensor product of the Galois symbol and the canonical isomorphism $\text{Hom}_c(\text{Gal}(K^{ab}/K), \mathbf{Z}/p^r) \cong H^1(K, \mathbf{Z}/p^r)$,

b' is the tensor product of $K^*/K^{*p^r} \cong H^1(K, \mu_{p^r})$ and $\text{Br}(K)_{p^r} \cong H^2(K, \mu_{p^r})$, where $\text{Br}(K)_{p^r}$ denotes the group $\{w \in \text{Br}(K) \mid p^r w = 0\}$,

c and c' are the cup products.

Consequently, we have a homomorphism from $K_2(K)$ to the pro- p -part of $\text{Gal}(K^{ab}/K)$ by (4) and a homomorphism from the p -primary part of $\text{Br}(K)$ to $\text{Hom}(K^*, \mathbf{Q}_p/\mathbf{Z}_p)$ by (5). These are the definitions of the p -primary parts of Φ and Ψ .

Proposition 1. *Let S be a field which is complete with respect to a discrete valuation and with residue field E . Suppose that $ch(E) = p > 0$, $ch(S) = 0$ and $[E : E^p] = p$. Then,*

(1) *the Galois symbol $K_2(S)/K_2(S)^{p^r} \rightarrow H^2(S, \mu_{p^r} \otimes \mu_{p^r})$ is an isomorphism for each $r \geq 0$.*

(2) *Suppose further that E is separably closed. Then $cd_p(S) = 2$. (Cf. [3] for the notation cd_p .)*

We need Proposition 2 (2) below to prove Proposition 1 (2).

Definition for Proposition 2. For each $i = 0, 1, 2$, we call a field S a B_i -field if and only if for each finite extension T of S and for each finite extension T' of T , the norm map $N_{T'/T} : K_i(T') \rightarrow K_i(T)$ is surjective.

This is an analogy of the concept "C_i-field". We can prove that a C_i-field is a B_i-field for each $i = 0, 1, 2$.

Proposition 2. *Let S be a field which is complete with respect to a discrete valuation and with residue field E . Suppose that E is a B₁-field. Then:*

(1) *For each central simple algebra A over S , $Nrd : A^* \rightarrow S^*$ is surjective.*

(2) *S is a B₂-field.*

Proposition 2 is an analogy of the following well known fact. "A field which is complete with respect to a discrete valuation is B₁ if its residue field is B₀ (i.e. algebraically closed)."

§ 3. The definitions of the "prime to p " parts of Φ and Ψ .

Let n be any natural number which is not divisible by $ch(F)$. Let G and K_{nr} be as in § 2. Then we have

$$(6) \quad \begin{aligned} H^3(K, \mu_n \otimes \mu_n) &\cong H^2(G, H^1(K_{nr}, \mu_n \otimes \mu_n)) \\ &\cong H^2(G, \mu_n) \cong \frac{1}{n} \mathbf{Z}/\mathbf{Z}, \end{aligned}$$

which can be easily deduced by the known facts in [3]. The composite of (6) induces a homomorphism from $K_2(K)$ to the "prime to p " part of $\text{Gal}(K^{ab}/K)$ and a homomorphism from the "prime to p " part of $\text{Br}(K)$ to $\text{Hom}(K^*, \mathbf{Q}/\mathbf{Z})$ in the same way as in § 2. These are the definitions of the "prime to p " parts of Φ and Ψ .

This simple argument cannot be adopted in case of § 2. The main difficulty in our theory lies in the p -primary part in the mixed characteristic case.

§ 4. The topologies of K^* and $K_2(K)$. In case $ch(F) = 0$, we take

the discrete topologies of K^* and $K_2(K)$. In what follows, suppose that $ch(F) = p > 0$.

Let R be the ring of integers of K , and m be the maximal ideal of R . First, we define the canonical topology of R/m^n for each n . Let $W(F)$ be the Witt ring of F (cf. [2]). Choose r such that $r \geq n-1$. Then there exists a unique ring-homomorphism $w_r: W(F) \rightarrow R/m^n$ such that

$$w_r(\bar{a}_0, \bar{a}_1, \bar{a}_2, \dots) \equiv \sum_{i=0}^r p^i a_i^{p^{r-i}} \pmod{m^n}$$

for all $a_i \in R$, where \bar{a}_i denotes the residue class of a_i . By w_r , R/m^n becomes a finitely generated $W(F)$ -module. We define the topology of R/m^n by regarding R/m^n as a quotient $W(F)$ -module of a finite product of $W(F)$. (Here the topology of $W(F)$ is the product topology of the valuation topology of F .) This topology of R/m^n is independent of the choice of r . In this way, R/m^n becomes a topological ring and $(R/m^n)^*$ becomes a topological group for the induced topology.

We define the topology of R^* by regarding R^* as the inverse limit of $(R/m^n)^*$ as $n \rightarrow \infty$. We define the topology of K^* in such a way that R^* becomes an open subgroup of K^* .

Finally, we define the topology of $K_2(K)$ by the following characterization. For each commutative topological group H and for each group-homomorphism $h: K_2(K) \rightarrow H$, h is continuous if and only if the composite map

$$K^* \times K^* \longrightarrow H: (x, y) \longmapsto h(\{x, y\})$$

is continuous.

References

- [1] K. Kato: A generalization of local class field theory by using K -groups. II (to appear in these Proceedings).
- [2] J. P. Serre: *Corps locaux*. Paris, Hermann (1962).
- [3] —: *Cohomologie galoisienne*. Berlin, Springer (1964).
- [4] J. Tate: Symbols in arithmetic. Actes du Congrès International des Mathématiciens 1970, 1, Gauthier-Villars, Paris, 201–211 (1971).