

The rank and generators of Kihara's elliptic curve with torsion $\mathbf{Z}/4\mathbf{Z}$ over $\mathbf{Q}(t)$

By Andrej DUJELLA,^{*} Ivica GUSIĆ^{**} and Petra TADIĆ^{***}

(Communicated by Heisuke HIRONAKA, M.J.A., Sept. 14, 2015)

Abstract: For the elliptic curve E over $\mathbf{Q}(t)$ found by Kihara, with torsion group $\mathbf{Z}/4\mathbf{Z}$ and rank ≥ 5 , which is the current record for the rank of such curves, by using a suitable injective specialization, we determine exactly the rank and generators of $E(\mathbf{Q}(t))$.

Key words: Elliptic curve; specialization homomorphism; rank; torsion; generators; mwrank.

1. Introduction. By Mazur's theorem, we know that the torsion group of an elliptic curve over \mathbf{Q} is one of the following 15 groups: $\mathbf{Z}/n\mathbf{Z}$ with $1 \leq n \leq 10$ or $n = 12$, $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/2m\mathbf{Z}$ with $1 \leq m \leq 4$. The same 15 groups appear as possible torsion groups for elliptic curves over the field of rational functions $\mathbf{Q}(t)$. The current records for the rank of elliptic curves over $\mathbf{Q}(t)$ with prescribed torsion group can be found in the table [3]. Note that in this table for the most of torsion groups only the lower bounds for the rank of record curves are given. Indeed, it seems that only for the torsion group $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/4\mathbf{Z}$ the exact rank over $\mathbf{Q}(t)$ of the record curve can be found in literature. In fact, in [5], Dujella and Peral proved that the corresponding curve, obtained from the so called Diophantine triples, has rank equal to 4 and they provide the generators for the group. The proof uses the method introduced by Gusić and Tadić in [6] for an efficient search for injective specializations. In this paper, we will prove an analogous result for the curve with record rank over $\mathbf{Q}(t)$ with torsion group $\mathbf{Z}/4\mathbf{Z}$ found by Kihara [9] (see Theorem 2.1). Here we will use results from the recent paper [7], where the authors generalize and extend their method from [6]. In particular, by results of [7], now the method can be applied to curves with only one rational 2-torsion point.

Our main tool is [7, Theorem 1.3]. It deals with

2010 Mathematics Subject Classification. Primary 11G05, 14H52.

^{*}) Department of Mathematics, University of Zagreb, Bijenička cesta 30, 10000 Zagreb, Croatia.

^{**}) Faculty of Chemical Engin. and Techn., University of Zagreb, Marulićev trg 19, 10000 Zagreb, Croatia.

^{***}) Department of Economics and Tourism, Juraj Dobrila University of Pula, 52100 Pula, Croatia.

elliptic curves E given by $y^2 = x^3 + A(t)x^2 + B(t)x$, where $A, B \in \mathbf{Z}[t]$, with exactly one nontrivial 2-torsion point over $\mathbf{Q}(t)$. If $t_0 \in \mathbf{Q}$ satisfies the condition that for every nonconstant square-free divisor h of $B(t)$ or $A(t)^2 - 4B(t)$ in $\mathbf{Z}[t]$ the rational number $h(t_0)$ is not a square in \mathbf{Q} , then the specialized curve E_{t_0} is elliptic and the specialization homomorphism at t_0 is injective. If additionally there exist $P_1, \dots, P_r \in E(\mathbf{Q}(t))$ such that $P_1(t_0), \dots, P_r(t_0)$ are the free generators of $E(t_0)(\mathbf{Q})$, then $E(\mathbf{Q}(t))$ and $E(t_0)(\mathbf{Q})$ have the same rank r , and P_1, \dots, P_r are the free generators of $E(\mathbf{Q}(t))$.

We can mention here that by the methods from [7], it is easy to show that the general families of curves with torsion $\mathbf{Z}/10\mathbf{Z}$, $\mathbf{Z}/12\mathbf{Z}$ and $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/8\mathbf{Z}$ given in Kubert's paper [10] all have rank 0 over $\mathbf{Q}(t)$, as expected from the corresponding entries in the above table (see Remark 3.1).

2. Kihara's curve with rank ≥ 5 . In 2004, Kihara [9] constructed a curve over $\mathbf{Q}(t)$ with torsion group $\mathbf{Z}/4\mathbf{Z}$ and rank ≥ 5 . This improved his previous result [8] with rank ≥ 4 . We briefly describe Kihara's construction. The quartic curve H given by the equation

$$(2.1) \quad (x^2 - y^2)^2 + 2a(x^2 + y^2) + b = 0$$

is considered. Forcing five points with coordinates of the form (r, s) , (r, u) , (s, p) , (u, q) , (p, m) to satisfy (2.1) leads to a system of certain quadratic Diophantine equations, for which a parametric solution is found. By the transformation $X = (a^2 - b)y^2/x^2$ and $Y = (a^2 - b)y(b + ax^2 + ay^2)/x^3$, we get from H the elliptic curve E with equation

$$(2.2) \quad Y^2 = X(X^2 + (2a^2 + 2b)X + (a^2 - b)^2).$$

We can write (2.2) in the form

$$Y^2 = X^3 + A(t)X^2 + B(t)X,$$

where

$$\begin{aligned} A(t) = & 4 \cdot (6561t^{52} - 363636t^{51} + 9938430t^{50} \\ & - 173377920t^{49} + 2093381633t^{48} \\ & - 17874840696t^{47} + 104874633374t^{46} \\ & - 354077727932t^{45} - 225490507368t^{44} \\ & + 11426935382596t^{43} - 77979654468618t^{42} \\ & + 278213963503072t^{41} - 111253886611731t^{40} \\ & - 6018676884563976t^{39} + 50437302286983990t^{38} \\ & - 272612407880904180t^{37} \\ & + 1164419469986139655t^{36} \\ & - 4197430502686137512t^{35} \\ & + 13172874796371804604t^{34} \\ & - 36632311181492128960t^{33} \\ & + 91270726486568186066t^{32} \\ & - 205234200473064086512t^{31} \\ & + 418565949588822731196t^{30} \\ & - 776853016569799513688t^{29} \\ & + 1315193607411569449248t^{28} \\ & - 2034605787348730781688t^{27} \\ & + 2881061692467531957212t^{26} \\ & - 3743290609966430672640t^{25} \\ & + 4481632889126213095506t^{24} \\ & - 4982866114496291088464t^{23} \\ & + 5212858449376876320156t^{22} \\ & - 5228984497845291579880t^{21} \\ & + 5133683278868660716535t^{20} \\ & - 4993730543781934338052t^{19} \\ & + 4784499422236344389062t^{18} \\ & - 440888972356409443776t^{17} \\ & + 3784174478544654656557t^{16} \\ & - 2930449238068167436056t^{15} \\ & + 1987364904374352466086t^{14} \\ & - 1143489371242947035052t^{13} \\ & + 534571453903095183576t^{12} \\ & - 187222081778006813708t^{11} \\ & + 38190501318649624878t^{10} \end{aligned}$$

$$\begin{aligned} & + 3640463051927237920t^9 \\ & - 6612210530622475839t^8 \\ & + 2873353737226588120t^7 \\ & - 748108460242930642t^6 \\ & + 127972510817241756t^5 \\ & - 14371540294374703t^4 + 1013310571582176t^3 \\ & - 40376902667904t^2 + 671143753728t \\ & + 2176782336), \end{aligned}$$

$$\begin{aligned} B(t) = & 256t^2(t-1)^4(t+1)^4(t-3)^2(t-5)^2(3t-1)^2 \\ & \times (t-2)^4(t^2+2)^2(t^2-2t+3)^2(t^2+6t-1)^2 \\ & \times (7t^2-18t+23)^2(3t^2-2t+7)^2 \\ & \times (t^3+4t^2-5t+16)^2 \\ & \times (3t^4-17t^3+27t^2-43t+6)^2 \\ & \times (2t^4-7t^3+9t^2-11t-5)^2 \\ & \times (2t^4-17t^3+27t^2-25t+1)^2 \\ & \times (t^4-4t^3+6t^2-12t+1)^2 \\ & \times (5t^4-17t^3+27t^2-79t+16)^2 \\ & \times (t^4-28t^3+54t^2-92t+41)^2 \\ & \times (t^4-9t^3+15t^2-19t+4)^2. \end{aligned}$$

Here we give also the factorization of $A(t)^2 - 4B(t)$ because it is essential for the proof of Theorem 2.1.

$$\begin{aligned} A(t)^2 - 4B(t) = & 16(6561t^{52} - 393876t^{51} \\ & + 12044286t^{50} - 233179992t^{49} + 3037888017t^{48} \\ & - 27010254024t^{47} + 156557186174t^{46} \\ & - 431368937388t^{45} - 1778897440520t^{44} \\ & + 29118669267908t^{43} - 185554409423562t^{42} \\ & + 692712486737480t^{41} - 814770507947971t^{40} \\ & - 9637269870939544t^{39} + 91721999355372182t^{38} \\ & - 511726240316396532t^{37} \\ & + 2209210564256660999t^{36} \\ & - 7976932199520997736t^{35} \\ & + 24934025825524375740t^{34} \\ & - 68735161090418984560t^{33} \\ & + 168954992642195397618t^{32} \\ & - 372775248087820281744t^{31} \\ & + 740995668471372699516t^{30} \\ & - 1328945941188034678776t^{29} \end{aligned}$$

$$\begin{aligned}
& + 2149268803793268828000t^{28} \\
& - 3126651933116879854968t^{27} \\
& + 4072542345072750657372t^{26} \\
& - 4716734217685402094832t^{25} \\
& + 4814383096857805387890t^{24} \\
& - 4296609977077762663152t^{23} \\
& + 3376161521034367049052t^{22} \\
& - 2507937390234783739560t^{21} \\
& + 2175389746176683769207t^{20} \\
& - 2602425451380388744228t^{19} \\
& + 3581059482295238331078t^{18} \\
& - 4560635054158334637368t^{17} \\
& + 4966911843389715455741t^{16} \\
& - 4539431424438631575336t^{15} \\
& + 3452812241784521490182t^{14} \\
& - 2155004986056068938396t^{13} \\
& + 1071003286706676463160t^{12} \\
& - 395973787910697589516t^{11} \\
& + 87894481825369263726t^{10} \\
& + 4136860927499429288t^9 \\
& - 12949727214730449839t^8 \\
& + 5873221034815986696t^7 \\
& - 1551776386124274418t^6 \\
& + 266291310738984156t^5 - 29718680649281967t^4 \\
& + 2058088618943712t^3 - 79201521880704t^2 \\
& + 1255157987328t + 2176782336) \\
& \times (81t^{26} - 2058t^{25} + 22205t^{24} - 136914t^{23} \\
& + 569600t^{22} - 1941994t^{21} + 7144777t^{20} \\
& - 31865642t^{19} + 143465455t^{18} - 557913380t^{17} \\
& + 1796620282t^{16} - 4792045284t^{15} \\
& + 10672893440t^{14} - 19973820452t^{13} \\
& + 31471575770t^{12} - 41625786276t^{11} \\
& + 45790269127t^{10} - 41147326466t^9 \\
& + 29240715721t^8 - 15417678410t^7 \\
& + 5182080208t^6 - 459229234t^5 - 434078947t^4 \\
& + 182700750t^3 - 25979095t^2 + 933744t + 46656)^2.
\end{aligned}$$

Kihara in [9] found five independent points P_1, \dots, P_5 on this curve, corresponding to the five points on H mentioned above, showing that the rank

of E over $\mathbf{Q}(t)$ is ≥ 5 . The torsion subgroup is $\mathbf{Z}/4\mathbf{Z}$. Indeed, the point $T_1 = (a^2 - b, 2a(a^2 - b))$ on (2.2) is of order 4 since $2T_1 = (0, 0)$ and $4T_1 = \mathcal{O}$. Furthermore, from the factorizations of $B(t)$ and $A(t)^2 - 4B(t)$ we see that $T_1 \notin 2E(\mathbf{Q}(t))$ and that $E(\mathbf{Q}(t))$ has exactly one point of order 2. An alternative proof of this fact will be given in Section 3.

Our goal is to prove that the rank of E over $\mathbf{Q}(t)$ is exactly equal to 5 and to find the generators of $E(\mathbf{Q}(t))$. Computations with several specializations indicate that P_1, P_2, P_3, P_4, P_5 are not generators of $E(\mathbf{Q}(t))$. Indeed, from our results it will follow that they generate a subgroup of index 64 in $E(\mathbf{Q}(t))$.

In fact, it holds that $P_1 + P_i \in 2E(\mathbf{Q}(t))$ for $i = 2, 3, 4, 5$, i.e. there exist points W_2, W_3, W_4, W_5 of $E(\mathbf{Q}(t))$ such that $P_1 + P_i = 2W_i$, $i = 2, 3, 4, 5$. Since the torsion subgroup is $\mathbf{Z}/4\mathbf{Z}$, there are two choices for each W_i . We choose one of them. The x -coordinates of these points are

$$\begin{aligned}
x(W_2) &= 4(t-3)(3t-1)(t^2+2) \\
&\times (t^4 - 4t^3 + 6t^2 - 12t + 1) \\
&\times (3t^4 - 17t^3 + 27t^2 - 43t + 6) \\
&\times (t^4 - 9t^3 + 15t^2 - 19t + 4)(7t^2 - 18t + 23) \\
&\times (t^4 - 28t^3 + 54t^2 - 92t + 41)(t-2)^2 \\
&\times (9t^{12} - 110t^{11} + 576t^{10} - 2333t^9 + 7802t^8 \\
&\quad - 19832t^7 + 39488t^6 - 57374t^5 + 61421t^4 \\
&\quad - 42914t^3 + 16488t^2 - 701t - 216)^2(t+1)^4,
\end{aligned}$$

$$\begin{aligned}
x(W_3) &= -16(t-1)^3(t+1)^3(t-3)(t^2+2) \\
&\times (t^2+6t-1)(t^3+4t^2-5t+16) \\
&\times (t^4-9t^3+15t^2-19t+4)(2t^4-17t^3+27t^2 \\
&\quad - 25t+1)(3t^4-17t^3+27t^2-43t+6) \\
&\times (3t^2-2t+7)(t-2)^2t^2(3t-1)^2(t-5)^2 \\
&\times (t^4-28t^3+54t^2-92t+41)^2 \\
&\times (2t^4-7t^3+9t^2-11t-5)^2,
\end{aligned}$$

$$\begin{aligned}
x(W_4) &= 16t(t-1)^3(t+1)(3t-1)(t^2+2) \\
&\times (t^2+6t-1)(t^3+4t^2-5t+16) \\
&\times (3t^4-17t^3+27t^2-43t+6) \\
&\times (t^4-9t^3+15t^2-19t+4) \\
&\times (2t^4-17t^3+27t^2-25t+1) \\
&\times (3t^2-2t+7)(t-2)^2(t-5)^2(7t^2-18t+23)^2 \\
&\times (t^4-4t^3+6t^2-12t+1)^2
\end{aligned}$$

$$\begin{aligned} & \times (2t^4 - 7t^3 + 9t^2 - 11t - 5)^2(3t^7 - 29t^6 + 51t^5 \\ & - 136t^4 + 175t^3 - 267t^2 + 179t - 24)^2 / \\ & (8 - 55t - 29t^2 + 103t^3 - 120t^4 + 69t^5 \\ & - 27t^6 + 3t^7)^2, \\ x(W_5) &= 16(t - 1)^3(t - 2)^3(t - 3)^2(t^2 + 2)^2 \\ & \times (3t^2 - 2t + 7)^2(t^3 + 4t^2 - 5t + 16)^2 \\ & \times (t^4 - 28t^3 + 54t^2 - 92t + 41) \\ & \times (t^4 - 4t^3 + 6t^2 - 12t + 1) \\ & \times (3t^4 - 17t^3 + 27t^2 - 43t + 6) \\ & \times (t^4 - 9t^3 + 15t^2 - 19t + 4) \\ & \times (5t^4 - 17t^3 + 27t^2 - 79t + 16) \\ & \times (2t^4 - 17t^3 + 27t^2 - 25t + 1) \\ & \times (t^2 + 6t - 1)(t - 5)(3t - 1)(t + 1)t. \end{aligned}$$

We also give the x -coordinate of P_1 :

$$\begin{aligned} x(P_1) &= 16t(t - 2)^2(t - 3)(t - 5)(3t - 1)(t^2 + 2) \\ & \times (3t^2 - 2t + 7)(7t^2 - 18t + 23) \\ & \times (t^2 + 6t - 1)(t^3 + 4t^2 - 5t + 16) \\ & \times (2t^4 - 17t^3 + 27t^2 - 25t + 1) \\ & \times (2t^4 - 7t^3 + 9t^2 - 11t - 5) \\ & \times (3t^4 - 17t^3 + 27t^2 - 43t + 6) \\ & \times (t^4 - 4t^3 + 6t^2 - 12t + 1) \\ & \times (t^2 - 2t + 3)(5t^4 - 17t^3 + 27t^2 - 79t + 16) \\ & \times (t^4 - 9t^3 + 15t^2 - 19t + 4) \\ & \times (t^4 - 28t^3 + 54t^2 - 92t + 41) \\ & \times (3t^{13} - 128t^{12} + 1185t^{11} - 5018t^{10} + 13628t^9 \\ & - 27704t^8 + 44162t^7 - 63956t^6 + 84827t^5 \\ & - 100976t^4 + 92061t^3 - 52802t^2 + 10662t - 552)^2 / \\ & (12t^{11} - 219t^{10} + 1699t^9 - 7248t^8 + 21004t^7 \\ & - 45434t^6 + 72862t^5 - 90128t^4 + 77496t^3 \\ & - 46283t^2 + 10095t - 768)^2. \end{aligned}$$

The points P_1, W_2, \dots, W_5 are a natural guess for the generators, and we will show that this is indeed true by proving the following theorem in the next section.

Theorem 2.1. *The elliptic curve E over $\mathbf{Q}(t)$ has rank equal to 5 with free generators the points P_1, W_2, W_3, W_4, W_5 and the torsion group is $\mathbf{Z}/4\mathbf{Z}$.*

3. An injective specialization. As described in the introduction, we use [7, Theorem 1.3] to find rational numbers t_0 for which the special-

ization map at t_0 is injective. The condition is that for each nonconstant square-free divisor h of $B(t)$ or $A(t)^2 - 4B(t)$ in $\mathbf{Z}[t]$ the rational number $h(t_0)$ is not a square in \mathbf{Q} . The condition is easy to check, and we can find many rationals t_0 satisfying it. However, the coefficients of the curve E are polynomials with large degrees and coefficients. Thus, for the success of our approach, it is crucial to find suitable specialization t_0 of reasonably small height. Furthermore, we need a specialization for which the rank of E_{t_0} over \mathbf{Q} is equal to 5, so it is reasonable to consider only specializations for which the the root number of E_{t_0} is -1 (conjecturally implying that the rank is odd).

We find that the specialization at $t_0 = -\frac{11}{4}$ satisfies all requirements, see preceding section for the factoriozation of $B(t)$ and $A(t)^2 - 4B(t)$. It remains to compute the rank and generators of $E_{-11/4}$. For that purpose, we use the excellent program [2] of Cremona, which is included in the program package Sage [13]. By extending significantly the default precision (we use options `-p 800 -b 11`), we get the elliptic curve $E_{-11/4}$ over \mathbf{Q} , given by the equation

$$\begin{aligned} y^2 &= x^3 + 4843712051739169544755051773863036- \\ & 55600428018856419825361x^2 \\ & + 1991079455035325445414429226070637115- \\ & 56495848199368164800^2x, \end{aligned}$$

which is of rank 5 with five free generators G_1, \dots, G_5 and the generator of the torsion group T_0 , given by their x -coordinates ,

$$\begin{aligned} x(T_0) &= -199107945503532544541442922607063711556495848199368164800, \\ x(G_1) &= \frac{35128929795293330966584382924686967322464932483259844000000}{49}, \\ x(G_2) &= \frac{13315895444389669790150691801504854333267656768880200356488633890879100326435093056000}{76159758997263590677307690401}, \\ x(G_3) &= \frac{536936685248255028234168808334606137477987482678373004839681994240}{1416844881}, \\ x(G_4) &= \frac{477616878094060797794543416082366648044181864476558959811339468000}{9223369}, \\ x(G_5) &= \frac{727946070485627419024954469098052196447100359265007972065657045989202204731022849600}{13386309077372650899951050449}. \end{aligned}$$

The rank 5 and generators are also confirmed in the most recent version V2.20-10 of Magma [1] (by the function `MordellWeilShaInformation` with option `SetClassGroupBounds("GRH")`). Now denote by $P_1^*, W_2^*, \dots, W_5^*$ the points obtained from P_1, W_2, \dots, W_5 after specialization $t \mapsto -\frac{11}{4}$. We easily get that

$$\begin{aligned} P_1^* &= T_0 + G_5, \\ W_2^* &= T_0 - G_3 + G_4, \\ W_3^* &= T_0 + G_1 + G_2 - G_3 + G_4 - G_5, \end{aligned}$$

$$\begin{aligned} W_4^* &= T_0 - G_2 + G_3 - G_4 + G_5, \\ W_5^* &= T_0 + G_4. \end{aligned}$$

Here all points are chosen up to sign of y -coordinates. It is easy to check that the matrix of this base change (modulo torsion) is of determinant ± 1 (the signs of the determinant depends on the choice of the signs of y -coordinates), so we see that $P_1^*, W_2^*, \dots, W_5^*$ are free generators of the elliptic curve $E_{-11/4}$ over \mathbf{Q} . From the comments at the end of the introduction, we see that [7, Theorem 1.3] now implies that E has rank 5 over $\mathbf{Q}(t)$ and that P_1, W_2, W_3, W_4, W_5 are its free generators. Since E has a point of fourth order and the torsion group of $E_{-11/4}(\mathbf{Q})$ is $\mathbf{Z}/4\mathbf{Z}$, we conclude that the torsion group of $E(\mathbf{Q}(t))$ is also $\mathbf{Z}/4\mathbf{Z}$. \square

Remark 3.1. Here we prove that the elliptic curves with torsion $\mathbf{Z}/10\mathbf{Z}$, $\mathbf{Z}/12\mathbf{Z}$, $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/8\mathbf{Z}$ and with rank ≥ 0 listed in Kubert's paper [10] have rank equal to 0 over $\mathbf{Q}(t)$.

For torsion group $\mathbf{Z}/10\mathbf{Z}$, the curve is given by the equation

$$\begin{aligned} y^2 &= x(x^2 - (2t^2 - 2t + 1)(4t^4 - 12t^3 + 6t^2 + 2t - 1) \\ &\quad \times x + 16t^5(t - 1)^5(t^2 - 3t + 1)). \end{aligned}$$

The specialization for $t_0 = 6$ satisfies the condition of [7, Theorem 1.3], and the specialized elliptic curve has rank 0, which proves our claim for that torsion group.

For torsion group $\mathbf{Z}/12\mathbf{Z}$, the curve is

$$\begin{aligned} y^2 &= x(x^2 + (t^8 - 12t^6 - 48t^5 - 162t^4 - 480t^3 - 540t^2 \\ &\quad - 624t - 183)x + 1024(t^2 + 3)^2(t + 1)^6), \end{aligned}$$

and the specialization which satisfies the condition of [7, Theorem 1.3] and has rank 0 is $t_0 = 11$.

Finally, for torsion group $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/8\mathbf{Z}$, we have the curve

$$y^2 = x(x + (t^2 - 1)^4)(x + 16t^4).$$

The specialization for $t_0 = 6$ satisfies the condition of [7, Theorem 1.1], and the specialized elliptic curve has rank 0, which proves our claim in that case.

Remark 3.2. An alternative method for computing the rank of an elliptic curves over $\mathbf{Q}(t)$ (and over $\mathbf{C}(t)$) is to use theory of Mordell-Weil

lattices, in particular the Shioda-Tate formula [12]. This method does not require that the curve has nontrivial 2-torsion and it is very efficient if the corresponding surface is rational or $K3$ surface (see e.g. [4,11,14]).

Acknowledgement. The authors were supported by the Croatian Science Foundation under the project no. 6422.

References

- [1] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.* **24** (1997), no. 3-4, 235-265.
- [2] J. E. Cremona, *Algorithms for modular elliptic curves*, 2nd ed., Cambridge Univ. Press, Cambridge, 1997.
- [3] A. Dujella, Infinite families of elliptic curves with high rank and prescribed torsion, <http://web.math.pmf.unizg.hr/~duje/tors/generic.html>.
- [4] A. Dujella and J. C. Peral, Elliptic curves and triangles with three rational medians, *J. Number Theory* **133** (2013), no. 6, 2083-2091.
- [5] A. Dujella and J. C. Peral, High-rank elliptic curves with torsion $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/4\mathbf{Z}$ induced by Diophantine triples, *LMS J. Comput. Math.* **17** (2014), no. 1, 282-288.
- [6] I. Gusić and P. Tadić, A remark on the injectivity of the specialization homomorphism, *Glas. Mat. Ser. III* **47(67)** (2012), no. 2, 265-275.
- [7] I. Gusić and P. Tadić, Injectivity of the specialization homomorphism of elliptic curves, *J. Number Theory* **148** (2015), 137-152.
- [8] S. Kihara, On the rank of the elliptic curves with a rational point of order 4, *Proc. Japan Acad. Ser. A Math. Sci.* **80** (2004), no. 4, 26-27.
- [9] S. Kihara, On the rank of the elliptic curves with a rational point of order 4. II, *Proc. Japan Acad. Ser. A Math. Sci.* **80** (2004), no. 8, 158-159.
- [10] D. S. Kubert, Universal bounds on the torsion of elliptic curves, *Proc. London Math. Soc. (3)* **33** (1976), no. 2, 193-237.
- [11] M. Kuwata and J. Top, An elliptic surface related to sums of consecutive squares, *Exposition. Math.* **12** (1994), no. 2, 181-192.
- [12] T. Shioda, On the Mordell-Weil lattices, *Comment. Math. Univ. St. Paul.* **39** (1990), no. 2, 211-240.
- [13] W. A. Stein *et al.*, Sage Mathematics Software (Version 6.0), The Sage Development Team, 2013, <http://www.sagemath.org>.
- [14] J. Top and F. De Zeeuw, Explicit elliptic $K3$ surfaces with rank 15, *Rocky Mountain J. Math.* **39** (2009), no. 5, 1689-1697.