

On the Geometry of the Tangent Bundle with the Cheeger-Gromoll Metric

Sigmundur GUDMUNDSSON and Elias KAPPOS

Lund University

(Communicated by R. Miyaoka)

Abstract. Let (M, g) be a Riemannian manifold of constant sectional curvature κ and (TM, \tilde{g}) be the tangent bundle of M equipped with the Cheeger-Gromoll metric induced by g . We give necessary and sufficient conditions for TM having positive scalar curvature. This gives counterexamples to a stated theorem of Sekizawa.

1. Introduction.

A Riemannian metric g on a smooth manifold M gives rise to several natural Riemannian metrics on the tangent bundle TM of M . The best known example is the Sasaki metric \hat{g} introduced in [6], see also [2]. In the present paper we study tangent bundles equipped with the so called Cheeger-Gromoll metric. Its construction was suggested in [1] but the first explicit description was given by Musso and Tricerri in [5].

In [7] Sekizawa calculates the Levi-Civita connection $\tilde{\nabla}$, the curvature tensor \tilde{R} , the sectional curvatures \tilde{K} and the scalar curvature \tilde{S} of the Cheeger-Gromoll metric. He then states in his Theorem 6.3 that if (M, g) is an m -dimensional manifold of constant sectional curvature $\kappa \geq -3(m-2)/m$, then (TM, \tilde{g}) has non-negative scalar curvature.

In this paper we prove the following results giving counterexamples to Sekizawa's statement.

THEOREM 1.1. *Let (M, g) be a surface of constant sectional curvature κ . Then there exists a real number $C_2 \geq 40$ such that the tangent bundle (TM, \tilde{g})*

- i. *has positive scalar curvature if and only if $\kappa \in [0, C_2)$,*
- ii. *has non-negative scalar curvature if and only if $\kappa \in [0, C_2]$.*

THEOREM 1.2. *Let (M, g) be a Riemannian manifold of dimension $m > 2$ and of constant sectional curvature κ . Then there exist real numbers $c_m < 0$ and $C_m > 60$ such that the tangent bundle (TM, \tilde{g})*

- i. *has positive scalar curvature if and only if $\kappa \in (c_m, C_m)$,*
- ii. *has non-negative scalar curvature if and only if $\kappa \in [c_m, C_m]$.*

Received June 5, 2001

1991 *Mathematics Subject Classification.* 53C25.

Key words and phrases. tangent bundles, natural metrics.

For further details on the geometry of tangent bundles equipped with natural metrics, such as those of Sasaki and Cheeger-Gromoll we refer the reader to the recent survey given in [4].

2. The Cheeger-Gromoll metric.

Let (M, g) be a Riemannian manifold and let (p, u) be a point on the tangent bundle TM of M . Then the Levi-Civita connection ∇ on M induces a natural splitting of the tangent space $T_{(p,u)}TM$ into its vertical and horizontal subspaces

$$T_{(p,u)} = \mathcal{V}_{(p,u)} \oplus \mathcal{H}_{(p,u)}.$$

This gives rise to the vertical and horizontal lifts X^v, X^h of a vector field X on M . The vertical subspace is the kernel of the bundle map $\pi : TTM \rightarrow TM$ at the point (p, u) . A horizontal curve in TM corresponds to a vector field on M which is parallel with respect to the Levi-Civita connection ∇ on (M, g) . The vector field U on TM given by $U_{(p,u)} = (u_p)^v$ is called *the canonical vector field*.

DEFINITION 2.1. Let (M, g) be a Riemannian manifold. Then the *Cheeger-Gromoll metric* \tilde{g} on TM is the Riemannian metric on the tangent bundle TM given by

- i. $\tilde{g}_{(p,u)}(X^h, Y^h) = g_p(X, Y)$,
- ii. $\tilde{g}_{(p,u)}(X^h, Y^v) = 0$,
- iii. $\tilde{g}_{(p,u)}(X^v, Y^v) = \frac{1}{\alpha}(g_p(X, Y) + g_p(X, u)g_p(Y, u))$

for all vectors $X, Y \in T_pM$. Here $\alpha = 1 + g(u, u)$.

In [7] Sekizawa calculates the Levi-Civita connection $\tilde{\nabla}$ and the curvature tensor \tilde{R} of the Cheeger-Gromoll metric on TM . His results are partially contained in Lemmata 2.2 and 2.4.

LEMMA 2.2. Let (M, g) be a Riemannian manifold and TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} . Then the Levi-Civita connection $\tilde{\nabla}$ of (TM, \tilde{g}) satisfies the following:

$$\begin{aligned} \tilde{\nabla}_{X^v} Y^v &= \frac{1}{\alpha}((1 + \alpha)\tilde{g}(X^v, Y^v)U - \tilde{g}(X^v, U)\tilde{g}(Y^v, U)U \\ &\quad - \tilde{g}(X^v, U)Y^v - \tilde{g}(Y^v, U)X^v). \end{aligned}$$

COROLLARY 2.3. Let (M, g) be a Riemannian manifold and TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} . Then the projection map $\pi : TM \rightarrow M$ is a harmonic morphism.

For further information on harmonic morphisms between Riemannian manifolds see the regularly updated bibliography on [3].

PROOF. It follows from Definition 2.1 and Lemma 2.2 that $\pi : TM \rightarrow M$ is a Riemannian submersion with totally geodesic fibres. Hence it is a harmonic morphism. \square

LEMMA 2.4. *Let (M, g) be a Riemannian manifold and TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} . Then the curvature tensor \tilde{R} of (TM, \tilde{g}) satisfies the following:*

$$\begin{aligned}\tilde{R}(X^v, Y^v)Z^v &= \frac{\alpha + 2}{\alpha^2}(\tilde{g}(X^v, Z^v)g(Y, u)U - \tilde{g}(Y^v, Z^v)g(X, u)U) \\ &\quad + \frac{1 + \alpha + \alpha^2}{\alpha^2}(\tilde{g}(Y^v, Z^v)X^v - \tilde{g}(X^v, Z^v)Y^v) \\ &\quad + \frac{\alpha + 2}{\alpha^2}(g(X, u)g(Z, u)Y^v - g(Y, u)g(Z, u)X^v).\end{aligned}$$

In his paper Sekizawa attempts to calculate the sectional curvatures \tilde{K} and the scalar curvature \tilde{S} for (TM, \tilde{g}) but unfortunately his calculations are wrong. In the rest of this section we shall correct the error and obtain valid expressions for \tilde{K} and \tilde{S} .

Let $\|\cdot\|$ denote the norm with respect to the Cheeger-Gromoll metric \tilde{g} and let $\tilde{Q}(V, W)$ be the square of the area of the parallelogram with sides $V, W \in T_{(p,u)}TM$ given by

$$\tilde{Q}(V, W) = \|V\|^2\|W\|^2 - \tilde{g}(V, W)^2.$$

Furthermore let \tilde{G} be the $(2, 0)$ -tensor on the tangent bundle TM given by

$$\tilde{G} : (V, W) \mapsto \tilde{g}(\tilde{R}(V, W)W, V).$$

LEMMA 2.5. [7] *Let $X, Y \in T_pM$ be two orthonormal vectors in the tangent spaces T_pM of M at p . Then*

- i. $\tilde{Q}(X^h, Y^h) = 1$,
- ii. $\tilde{Q}(X^h, Y^v) = \frac{1}{\alpha}(1 + g(Y, u)^2)$,
- iii. $\tilde{Q}(X^v, Y^v) = \frac{1}{\alpha^2}(1 + g(Y, u)^2 + g(X, u)^2)$.

Sekizawa's mistake is contained in the proof of the last part of the following Lemma.

LEMMA 2.6. *Let X, Y be two orthonormal vectors in the tangent space T_pM of M at p . Then*

- i. $\tilde{G}(X^h, Y^h) = K(X, Y) - \frac{3}{4\alpha}|R(X, Y)u|^2$,
- ii. $\tilde{G}(X^h, Y^v) = \frac{1}{4\alpha^2}|R(u, Y)X|^2$,
- iii. $\tilde{G}(X^v, Y^v) = \frac{(1+\alpha+\alpha^2)}{\alpha^2}\tilde{Q}(X^v, Y^v) - \frac{(\alpha+2)}{\alpha^3}(g(X, u)^2 + g(Y, u)^2)$.

PROOF. Here we only prove iii. The rest can be found in [7].

$$\begin{aligned}\tilde{G}(X^v, Y^v) &= \tilde{g}(\tilde{R}(X^v, Y^v)Y^v, X^v) \\ &\quad + \frac{(\alpha + 2)}{\alpha^2}(\tilde{g}(X^v, Y^v)g(Y, u)g(X, u) - \tilde{g}(Y^v, Y^v)g(X, u)^2) \\ &\quad + \frac{(1 + \alpha + \alpha^2)}{\alpha^2}(\tilde{g}(Y^v, Y^v)\tilde{g}(X^v, X^v) - \tilde{g}(X^v, Y^v)^2) \\ &\quad + \frac{(\alpha + 2)}{\alpha^2}(g(X, u)g(Y, u)\tilde{g}(X^v, Y^v) - g(Y, u)^2\tilde{g}(X^v, X^v))\end{aligned}$$

$$= \frac{(1 + \alpha + \alpha^2)}{\alpha^2} \tilde{Q}(X^v, Y^v) - \frac{(\alpha + 2)}{\alpha^3} (g(X, u)^2 + g(Y, u)^2).$$

□

PROPOSITION 2.7. *Let (M, g) be a Riemannian manifold and TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} . Then the sectional curvatures \tilde{K} of (TM, \tilde{g}) satisfy the following:*

- i. $\tilde{K}(X^h, Y^h) = K(X, Y) - \frac{3}{4\alpha} |R(X, Y)u|^2$,
- ii. $\tilde{K}(X^h, Y^v) = \frac{1}{4\alpha} \frac{|R(u, Y)X|^2}{(1+g(Y, u)^2)}$,
- iii. $\tilde{K}(X^v, Y^v) = \frac{1-\alpha}{\alpha^2} + \frac{\alpha+2}{\alpha} \frac{1}{(1+g(X, u)^2+g(Y, u)^2)}$.

PROOF. The statements follow directly from the division of $\tilde{G}(X^i, Y^j)$ by $\tilde{Q}(X^i, Y^j)$ for $i, j \in \{h, v\}$. □

For a given point $(p, u) \in TM$ with $u \neq 0$ let $\{e_1, \dots, e_m\}$ be an orthonormal basis for the tangent space $T_p M$ of M at p such that $e_1 = u/|u|$, where $|u|$ is the norm of u with respect to the metric g on M . Then for $i \in \{1, \dots, m\}$ and $k \in \{2, \dots, m\}$ define the horizontal and vertical lifts by $f_i = e_i^h$, $f_{m+1} = e_1^v$ and $f_{m+k} = \sqrt{\alpha} e_k^v$. Then $\{f_1, \dots, f_{2m}\}$ is an orthonormal basis for the tangent space $T_{(p,u)} M$ with respect to the Cheeger-Gromoll metric.

LEMMA 2.8. *Let (p, u) be a point on TM and $\{f_1, \dots, f_{2m}\}$ be an orthonormal basis for the tangent space $T_{(p,u)} TM$ as above. Then the sectional curvatures \tilde{K} satisfy the following equations.*

$$\begin{aligned} \tilde{K}(f_i, f_j) &= K(e_i, e_j) - \frac{3}{4\alpha} |R(e_i, e_j)u|^2, \\ \tilde{K}(f_i, f_{m+1}) &= 0, \\ \tilde{K}(f_i, f_{m+k}) &= \frac{1}{4} |R(u, e_k)e_i|^2, \\ \tilde{K}(f_{m+1}, f_{m+k}) &= \frac{3}{\alpha^2}, \\ \tilde{K}(f_{m+k}, f_{m+l}) &= \frac{\alpha^2 + \alpha + 1}{\alpha^2}, \end{aligned}$$

for $i, j \in \{1, \dots, m\}$ and $k, l \in \{2, \dots, m\}$.

PROOF. The result is a direct consequence of Proposition 2.7. □

PROPOSITION 2.9. *Let (M, g) be a Riemannian manifold with scalar curvature S . Let TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} and (p, u) be a point on TM . Then the scalar curvature \tilde{S} of (TM, \tilde{g}) satisfies the following:*

$$\tilde{S}_{(p,u)} = S_p + \frac{(2\alpha - 3)}{4\alpha} \sum_{i,j=1}^m |R(e_i, e_j)u|^2$$

$$+ \frac{(m-1)}{\alpha^2} (6 + (m-2)(\alpha^2 + \alpha + 1)).$$

PROOF. Let $\{f_1, \dots, f_{2m}\}$ be an orthonormal basis for the tangent space $T_{(p,u)}TM$ as above. By the definition of the scalar curvature we know that

$$\begin{aligned} \tilde{S} &= \sum_{i \neq j} \tilde{K}(f_i, f_j) \\ &= 2 \sum_{\substack{i,j=1 \\ i < j}}^m \tilde{K}(f_i, f_j) + 2 \sum_{i,j=1}^m \tilde{K}(f_i, f_{m+j}) + 2 \sum_{\substack{i,j=1 \\ i < j}}^m \tilde{K}(f_{m+i}, f_{m+j}) \\ &= \sum_{i \neq j}^m K(e_i, e_j) - \frac{3}{4\alpha} \sum_{i,j=1}^m |R(e_i, e_j)u|^2 \\ &\quad + \frac{1}{2} \sum_{i,j=1}^m |R(u, e_j)e_i|^2 + 2 \sum_{i=2}^m \frac{3}{\alpha^2} + \sum_{\substack{i,j=2 \\ i \neq j}}^m \frac{(1 + \alpha + \alpha^2)}{\alpha^2} \\ &= S + \frac{(2\alpha - 3)}{4\alpha} \sum_{i,j=1}^m |R(e_i, e_j)u|^2 + \frac{(m-1)}{\alpha^2} (6 + (m-2)(1 + \alpha + \alpha^2)). \end{aligned}$$

For the above calculations we have used Lemma 2.10. \square

LEMMA 2.10. Let (M, g) be a Riemannian manifold with curvature tensor R . If $p \in M$ and $\{e_1, \dots, e_m\}$ is an orthonormal basis for the tangent space T_pM , then

$$\sum_{i,j=1}^m |R(e_j, u)e_i|^2 = \sum_{i,j=1}^m |R(e_j, e_i)u|^2.$$

PROOF. With $u = \sum_{i=1}^m u_i X_i$ we get

$$\begin{aligned} \sum_{i,j=1}^m |R(e_j, u)e_i|^2 &= \sum_{i,j,k,l=1}^m u_k u_l g(R(e_j, e_k)e_i, R(e_j, e_l)e_i) \\ &= \sum_{i,j,k,l,s=1}^m u_k u_l g(R(e_s, e_i)e_k, e_j) g(R(e_s, e_i)e_l, e_j) \\ &= \sum_{i,j,k,l=1}^m u_k u_l g(R(e_j, e_i)e_k, R(e_j, e_i)e_l) \\ &= \sum_{i,j=1}^m |R(e_j, e_i)u|^2. \end{aligned}$$

\square

3. The constant curvature case.

In this section we shall assume that (M, g) is an m -dimensional Riemannian manifold of constant sectional curvature κ . This implies that its curvature tensor R takes the special form

$$R(X, Y)Z = \kappa(g(Y, Z)X - g(X, Z)Y)$$

and the scalar curvature S is given by $S = \kappa m(m - 1)$.

PROPOSITION 3.1. *Let (M, g) be a Riemannian manifold of constant sectional curvature κ . Let TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} . Then the sectional curvatures \tilde{K} of (TM, \tilde{g}) satisfy the following:*

- i. $\tilde{K}(X^h, Y^h) = \kappa - \frac{3\kappa^2}{4\alpha}(g(u, X)^2 + g(u, Y)^2)$,
- ii. $\tilde{K}(X^h, Y^v) = \frac{\kappa^2 g(X, u)^2}{4\alpha(1+g(Y, u)^2)}$,
- iii. $\tilde{K}(X^v, Y^v) = \frac{1-\alpha}{\alpha^2} + \frac{\alpha+2}{\alpha} \frac{1}{1+g(X, u)^2+g(Y, u)^2}$,

for any orthonormal vectors $X, Y \in T_p M$.

PROOF. This is a simple calculation using the special form of the curvature tensor. \square

PROPOSITION 3.2. *Let (M, g) be a Riemannian manifold of constant sectional curvature κ . Let TM be its tangent bundle equipped with the Cheeger-Gromoll metric \tilde{g} and \tilde{K} be the sectional curvatures of (TM, \tilde{g}) . Then*

- i. $\tilde{K}(X^h, Y^h)$ is non-negative if $0 \leq \kappa \leq \frac{4}{3}$,
- ii. $\tilde{K}(X^h, Y^v)$ is non-negative,
- iii. $\tilde{K}(X^v, Y^v)$ is positive.

PROOF. If $X, Y \in T_p M$ are orthonormal, then obviously

$$g(Y, u)^2 + g(X, u)^2 \leq |u|^2 < \alpha.$$

With this in hand the result follows directly by Proposition 3.1. \square

PROPOSITION 3.3. *Let (M, g) be a Riemann manifold of constant sectional curvature κ . Let (TM, \tilde{g}) be the tangent bundle equipped with the Cheeger-Gromoll metric. Then the scalar curvature \tilde{S} of TM is given by*

$$\begin{aligned} \tilde{S} &= \frac{(m-1)}{2\alpha^2} (\alpha(\alpha-1)(2\alpha-3)\kappa^2 + 2m\alpha^2\kappa \\ &\quad + 2(6+(m-2)(1+\alpha+\alpha^2))). \end{aligned}$$

PROOF. The result follows from Proposition 2.9 and the following calculations

$$\begin{aligned} \sum_{i,j=1}^m |R(e_i, e_j)u|^2 &= \sum_{i,j=1}^m (\alpha-1)\kappa^2 |\delta_{1j}e_i - \delta_{i1}e_j|^2 \\ &= 2(m-1)(\alpha-1)\kappa^2. \end{aligned}$$

\square




FIGURE 1.

For a given $m \geq 2$, we are now interested in determining the sign of the scalar curvature $\tilde{S}_m(\alpha, \kappa)$ as a function of $(\alpha, \kappa) \in D = [1, \infty) \times \mathbf{R}$. The contour $D_0 = \{(\alpha, \kappa) \in D \mid \tilde{S}_m(\alpha, \kappa) = 0\}$ in the (α, κ) -plane is determined by the second order polynomial equation

$$(1) \quad \alpha(\alpha - 1)(2\alpha - 3)\kappa^2 + 2m\alpha^2\kappa + 2(6 + (m - 2)(1 + \alpha + \alpha^2)) = 0$$

in κ . If $\alpha \neq 1$, $\alpha \neq 3/2$ and the discriminant of equation (1) is non-negative we get the solutions

$$\kappa_{\pm} = \frac{-m\alpha^2 \pm \sqrt{m^2\alpha^4 - \alpha(\alpha - 1)(2\alpha - 3)2(6 + (m - 2)(1 + \alpha + \alpha^2))}}{\alpha(\alpha - 1)(2\alpha - 3)}.$$

In Figure 1 we have plotted the contour D_0 in the (α, κ) -plane for the case when $m = 3$. When removing D_0 from D the rest falls into three connected components. The scalar curvature is positive in the component D_+ containing the point $(2, 40)$ and negative in the other two.

We are interested in determining those $\kappa \in \mathbf{R}$ such that $S_3(\alpha, \kappa)$ is positive (non-negative) for all $\alpha \in [1, \infty)$ i.e. which are the horizontal half-lines completely contained in the component D_+ . The connected component of D_0 contained in the upper halfspace ($\kappa > 0$) is the graph of the solution κ_- for $\alpha \in (1, 3/2)$. It has exactly one minimum $C_3 > 0$.




FIGURE 2.

The graph of the other solution κ_+ , where defined, has exactly one maximum $c_3 < 0$. The family of horizontal lines that we are looking for are then parametrized by $\kappa \in (c_3, C_3)$.

It is easy to see that for $m > 3$ we get exactly the same qualitative behaviour of the two solutions κ_- and κ_+ as for $m = 3$. This provides us with the following result:

THEOREM 3.4. *Let (M, g) be a Riemannian manifold of dimension $m > 2$ and of constant sectional curvature κ . Then there exist real numbers $c_m < 0$ and $C_m > 60$ such that the tangent bundle (TM, \tilde{g})*

- i. *has positive scalar curvature if and only if $\kappa \in (c_m, C_m)$,*
- ii. *has non-negative scalar curvature if and only if $\kappa \in [c_m, C_m]$.*

Notice that for $\alpha \in (1, 3/2)$ and $m \geq 3$ the function κ is increasing in m , so if $m < \bar{m}$ then $C_m < C_{\bar{m}}$.

When $m = 2$ the discriminant of equation (1) is positive everywhere, and the solution κ_- and κ_+ are given by

$$\kappa_{\pm} = \frac{-2\alpha^2 \pm 2\sqrt{\alpha^4 - 3\alpha(\alpha - 1)(2\alpha - 3)}}{\alpha(\alpha - 1)(2\alpha - 3)}.$$

In Figure 2 we have plotted the contour D_0 in the (α, κ) for $m = 2$. When this is removed from D the rest falls into four connected components. The scalar curvature is positive in the two components containing the points $(2, \pm 40)$ and negative in the others. When $\alpha > 3/2$ both the solutions κ_{\pm} are negative and approaching 0 in the limit $\alpha \rightarrow \infty$.

This leads to the following result

THEOREM 3.5. *Let (M, g) be a surface of constant sectional curvature κ . Then there exists a real number $C_2 \geq 40$ such that the tangent bundle (TM, \tilde{g})*

- i. *has positive scalar curvature if and only if $\kappa \in [0, C_2]$,*
- ii. *has non-negative scalar curvature if and only if $\kappa \in [0, C_2]$.*

ACKNOWLEDGMENTS. This paper has benefitted from our discussions with Prof. Masami Sekizawa and Martin Svensson.

References

- [1] J. CHEEGER and D. GROMOLL, On the structure of complete manifolds of nonnegative curvature, *Ann. of Math.* **96** (1972), 413–443.
- [2] P. DOMBROWSKI, On the Geometry of the Tangent Bundle, *J. Reine Angew. Math.* **210** (1962), 73–88.
- [3] S. GUDMUNDSSON, *The Bibliography of Harmonic Morphisms*, [<http://www.maths.lth.se/matematiklu/personal/sigma/harmonic/bibliography>].
- [4] E. KAPPOS, Natural Metrics on Tangent Bundles, Master's Dissertation, Lund University (2001), [<http://www.maths.lth.se/matematiklu/personal/sigma/students/Kappos.ps>].
- [5] E. MUSSO and F. TRICERRI, Riemannian Metrics on Tangent Bundles, *Ann. Mat. Pura Appl.* (4) **150** (1988), 1–19.
- [6] S. SASAKI, On the differential geometry of tangent bundles of Riemannian manifolds, *Tôhoku Math. J.* **10** (1958), 338–354.
- [7] M. SEKIZAWA, Curvatures of tangent bundles with Cheeger-Gromoll metric, *Tokyo J. Math.* **14** (1991), 407–417.

Present Addresses:

SIGMUNDUR GUDMUNDSSON
CENTRE FOR MATHEMATICAL SCIENCES, UNIVERSITY OF LUND,
BOX 118, S-221 00 LUND, SWEDEN.
e-mail: Sigmundur.Gudmundsson@math.lu.se

ELIAS KAPPOS
INSTITUT FÜR MATHEMATIK, GEORG-AUGUST-UNIVERSITÄT ZU GÖTTINGEN,
D-37073 GÖTTINGEN, GERMANY.
e-mail: ekappos@netscape.net