

ON THE VOLUME OF MANIFOLDS ALL OF WHOSE GEODESICS ARE CLOSED

ALAN WEINSTEIN

1. C_L -manifolds

A riemannian manifold (M, g) will be called a C_L -manifold if all the geodesics on M are closed and have length $2\pi L$, i.e., if all the orbits of the geodesic flow on the unit tangent bundle $U(M, g)$ are periodic with least period $2\pi L$. It is a problem of some interest to characterize these manifolds, which are the "simple harmonic oscillators" of riemannian geometry.

The best known examples of C_L -manifolds are the symmetric spaces of rank one, or SC -manifolds, as they are called by Berger [1, III. 4]. These are the spheres (S^n, can) , projective spaces $(P^n(K), \text{can})$ for $k = \mathbf{R}, \mathbf{C}$, or \mathbf{H} , and the Cayley projective plane $(P^2(I), \text{can})$, with their canonical metrics. The spheres are C_1 -manifolds, and the projective spaces are, with Berger's normalization, $C_{1/2}$ -manifolds.

Zoll (see [1, IV. 8]) in 1903 constructed examples of non-standard C_L -metrics on S^2 (surfaces of revolution), and Blaschke [3, p. 233] gives an example, due to Thomsen, of a C_L -metric on S^2 with no nontrivial isometries. These constructions can be carried out on higher dimensional spheres as well. If one strengthens the C_L condition to require that the geodesics be simple closed curves on M , then a theorem of Green (see [1, VIII. 9]) states that any such simple C_L -metric on $P^2(\mathbf{R})$ has constant curvature. Furthermore, it is a theorem of Bott (see [1, IV. 6]) that every simple C_L -manifold has the same integer cohomology ring as some SC -manifold. In fact, this result requires only that all the geodesics through a single point of M be simply closed with the same length. The earliest topological study of C_L -manifolds seems to be that of Reeb [7], who proved, among other things, that the product of two spheres of different odd dimensions cannot carry a C_L -metric.

The aim of the present paper is to demonstrate the following geometric result.

Theorem A. *If (M, g) is an n -dimensional C_L -manifold, then the ratio*

$$i(M, g) = \frac{\text{vol}(M, g)}{L^n \text{vol}(S^n, \text{can})}$$

is an integer.

We will actually prove the following theorem, of which Theorem A is an immediate consequence.

Theorem B. *If (M, g) is an n -dimensional C_L -manifold, then the real number $j(M, g)$ defined by the equation*

$$(1) \quad \text{vol}(M, g) = \frac{(2\pi L)^n \cdot j(M, g)}{(n - 1)! \text{vol}(S^{n-1}, \text{can})}$$

is an even integer.

To prove Theorem A from Theorem B, one has merely to check, using the values of $\text{vol}(S^{n-1}, \text{can})$ [1, VI. 7], that $j(S^n, \text{can}) = 2$; then set $i(M, g) = \frac{1}{2}j(M, g)$.

Remarks

1. The proof of Theorem B, contained in the following two sections of this paper, identifies the integer $j(M, g)$ as a topological invariant of the fibration of $U(M, g)$ by the orbits of the geodesic flow.

2. Using Gysin sequences one can prove that $j(M, g) = 2$ and $i(M, g) = 1$ if M is an even-dimensional sphere. It would be interesting to prove that $i(M, g)$ is independent of g when M is any SC -manifold. This may be a step in the direction of generalizing the theorem of Green mentioned above.

3. In the succeeding paper in this journal [2], Marcel Berger proves the following application of Theorem A. Let g be a Kählerian metric on $P^n(C)$, compatible with the standard complex structure. Suppose that the distance to the first conjugate point in each direction from each point on $P^n(C)$ is $\frac{1}{2}\pi$. Then, at least if g is sufficiently near the canonical metric in the C^0 topology, $(P^n(C), g)$ is isometric to $(P^n(C), \text{can})$.

4. Funk [4, p. 283] remarks that the area of a C_1 -surface of revolution must be 4π . Otherwise, our result seems to be new even for $M = S^2$.

5. An amusing consequence of Theorem A is that one cannot apply a slight perturbation to (S^n, can) to make the geodesics close only after $k > 1$ "revolutions", for then the volume of the manifold would have to be multiplied by k^n .

6. For reference, we present the following formulas, obtained from the calculations in [1, VI. 7]:

$$i(S^n, \text{can}) = 1, \quad i(P^n R, \text{can}) = 2^{n-1},$$

$$i(P^n C, \text{can}) = \binom{2n - 1}{n - 1}, \quad i(P^n H, \text{can}) = \frac{1}{2n + 1} \binom{4n - 1}{2n - 1},$$

$$i(P^2 \Gamma, \text{can}) = 39.$$

7. The work described here was carried out at the Institute for Advanced

Study, Princeton and Institut des Hautes Études Scientifiques, Bures-sur-yvette with support from a Sloan fellowship and was completed in Berkeley, with support from NSF grant GP-34785X. The author would like to thank M. Berger and F. Hirzebruch for their helpful comments.

2. Proof of Theorem B

The unit tangent bundle $U(M, g)$ of a riemannian manifold carries the following geometric objects:

- the geodesic spray G , [1, IV. 2],
- the canonical one-form α , [1, III. 6],
- the canonical two-form $d\alpha$, [1, III. 6],
- the riemannian metric \bar{g} , [1, V. 2.4],
- the volume element $\bar{\theta}$, [1, V. 2.4].

These objects satisfy the following relations:

- i. $\left| \frac{\alpha \wedge (d\alpha)^{n-1}}{(n-1)!} \right| = \bar{\theta}$, [1, V. 2.5],
- ii. $\text{vol}(U(M, g), \bar{g}) = \text{vol}(M, g) \cdot \text{vol}(S^{n-1}, \text{can})$, [1, V. 2.13],
- iii. the flow of G leaves α invariant [1, IV. 3.10],
- iv. $\alpha(G) \equiv 1$, [1, p. 125],
- v. the null space (characteristic distribution) of $d\alpha$ is generated by G , [5, Thm. 5.9].

Since the orbits of G are all periodic with period $2\pi L$, the vector field $2\pi LG$ generates a free action of $S^1 = \mathbf{R}/\mathbf{Z}$ on $U(M, g)$, with quotient a manifold $C(M, g)$. The projection $U(M, g) \xrightarrow{p} C(M, g)$ is a principal bundle with structure group S^1 . Relations iii and iv above mean that $\alpha/(2\pi L)$ is a connection form on this bundle, and $d\alpha/(2\pi L)$ is the curvature form. There is then a uniquely determined form Ω on $C(M, g)$ such that $p^*\Omega = d\alpha/(2\pi L)$; the de Rham cohomology class $[\Omega] \in H^2(C(M, g); \mathbf{R})$ is the image under the coefficient homomorphism $\rho_2: H^2(C(M, g); \mathbf{Z}) \rightarrow H^2(C(M, g); \mathbf{R})$ of the Euler class $e(p)$ of the bundle p . (We identify the group S^1 with $SO(2)$.) Then $[\Omega^{n-1}] = [\Omega]^{n-1}$ is the image of $[e(p)]^{n-1}$ under the coefficient homomorphism ρ_{2n-2} .

By relation v, the form Ω is nonsingular on $C(M, g)$, which is oriented by Ω^{n-1} . Denoting by $[C(M, g)]$ the fundamental $(2n-2)$ -cycle, we have

$$(2) \quad \int_{C(M, g)} \Omega^{n-1} = \langle [e(p)]^{n-1}, [C(M, g)] \rangle .$$

Let $j(M, g)$ be the quantity on either side of (2). The left hand side of the equation is positive, and the right hand side is an integer, so $j(M, g)$ is a positive integer.

The argument up to here is essentially contained in [7]. At this point, we use the Fubini theorem for fibrations [1, 0.3.17] to calculate

$$\begin{aligned} \text{vol}(U(M, g); \bar{g}) &= \int_{U(M, g)} \frac{\alpha \wedge (d\alpha)^{n-1}}{(n-1)!} \\ &= \frac{1}{(n-1)!} \int_{U(M, g)} \alpha \wedge p^*(2\pi L \Omega)^{n-1} \\ &= \frac{(2\pi L)^{n-1}}{(n-1)!} \int_{x \in C(M, g)} \left[\int_{p^{-1}(x)} \alpha \right] \Omega^{n-1}. \end{aligned}$$

By relation iv, $\int_{p^{-1}(x)} \alpha = 2\pi L$ for each x , so the above expression becomes

$$\frac{(2\pi L)^n}{(n-1)!} \int_{C(M, g)} \Omega^{n-1} = \frac{2\pi L}{(n-1)!} j(M, g).$$

Combining this with relation ii shows that $j(M, g)$ satisfies (1). To complete the proof of Theorem B, it remains only to show that $j(M, g)$ is even. This is done in the next section.

3. Involutions and evenness

Let $\xi: P \rightarrow B$ be a principal bundle with structure group $SO(2)$. By means of the embedding $SO(2) \rightarrow O(2)$, we can consider ξ as a bundle with fibre $SO(2)$ and structure group $O(2)$. Let $\beta: \xi \rightarrow \xi$ be a mapping of $O(2)$ bundles (see [8, 2.5] for a definition) with $\beta^2 = \text{identity}$ and such that the induced mapping $\gamma: B \rightarrow B$ has no fixed points. Suppose further that B is an orientable manifold of dimension $2n$.

Proposition. *The class $[e(\xi)]^n$ is an even multiple of the generator of $H^{2n}(B; \mathbb{Z}) \approx \mathbb{Z}$.*

Proof. By [6, 4.11.2 III], $[e(\xi)]^n = e(n\xi)$ where $n\xi$ is the $SO(2n)$ bundle obtained by taking the n -fold Whitney sum of ξ with itself. The involution β induces an involution $n\beta: n\xi \rightarrow n\xi$ of $O(2n)$ bundles; therefore there is an $O(2n)$ bundle $\overline{n\xi}$ over the quotient manifold $\overline{B} = B/\gamma$ such that $n\xi = \pi^* \overline{n\xi}$, $\pi: B \rightarrow \overline{B}$ being the projection. The Whitney classes of $n\xi$ and $\overline{n\xi}$ satisfy the relation $w_{2n}(n\xi) = \pi^* w_{2n}(\overline{n\xi})$ [6, p. 73]. Since π is a double covering, it induces the zero map from $H^{2n}(\overline{B}; \mathbb{Z}_2)$ to $H^{2n}(B; \mathbb{Z}_2)$, so $w_{2n}(n\xi) = 0$. But $w_{2n}(n\xi)$ is the mod 2 reduction [6, p. 73] of $e(n\xi) = e(\xi)^n$, so $[e(\xi)]^n$ is even. q.e.d.

To apply this Proposition to Theorem 2, we use the involution $h_{-1}: U(M, g) \rightarrow U(M, g)$ defined by multiplying each tangent vector by -1 . Since G is a spray, G is h_{-1} -related to $-G$, so h_{-1} is an $O(2)$ bundle mapping. Finally, the induced map on $C(M, g)$ has no fixed points, because a geodesic cannot double back upon itself in the reverse direction.

Hence the class $[e(p)]^{n-1}$ is an even multiple of the generator of $H^{2n-2}(C(M, g); \mathbb{Z})$, and $j(M, g) = \langle [e(p)]^{n-1}, [C(M, g)] \rangle$ is an even integer.

References

- [1] M. Berger, *Lectures on geodesics in riemannian geometry*, Tata Institute of Fundamental Research, Bombay, 1965.
- [2] —, *Sur certaines variétés kähleriennes à géodésiques toutes fermées*, J. Differential Geometry **9** (1974) 519–520.
- [3] W. Blaschke, *Vorlesungen über Differentialgeometrie*, 3rd. edition, Springer, Berlin, 1930.
- [4] P. Funk, *Über Flächen mit lauter geschlossenen geodätischen Linien*, Math. Ann. **74** (1913) 278–300.
- [5] C. Godbillon, *Géométrie différentielle et mécanique analytique*, Hermann, Paris, 1969.
- [6] F. Hirzebruch, *Topological methods in algebraic geometry*, Third enlarged edition (translated by R. L. E. Schwarzenberger), Springer, New York, 1966.
- [7] G. Reeb, *Variétés de Riemann dont toutes les géodésiques sont fermées*, Acad. Roy. Belg. Bull. Cl. Sci. (5) **36** (1950) 324–329.
- [8] N. Steenrod, *The topology of fibre bundles*, Princeton University Press, Princeton, 1951.

UNIVERSITY OF CALIFORNIA, BERKELEY

