

versely if the first row is multiplied by the inverse of $c \pmod{p^k}$. This inverse exists, and the correspondence is one-to-one, because c is prime to p . This proves (3).

The sum of the probabilities $P_n(ap^\alpha, p^k)$, where a runs through the values $1, 2, \dots, p^{k-\alpha}$, is clearly the probability that a determinant be divisible by p^α . The terms of this sum can be simplified and collected by use of (3), and we have

$$(11) \quad P_n(0, p^\alpha) = \sum_{r=0}^{k-\alpha} \phi(p^{k-\alpha-r}) P_n(p^{\alpha+r}, p^k).$$

Replacing α by $\alpha+1$, and subtracting the resulting equation from (11), we arrive at (4).

PURDUE UNIVERSITY

ON THE NOTION OF THE RING OF QUOTIENTS OF A PRIME IDEAL

CLAUDE CHEVALLEY

Let \mathfrak{o} be a domain of integrity (that is, a ring with unit element and with no zero divisor not equal to 0), and let \mathfrak{u} be a prime ideal in \mathfrak{o} . We can construct two auxiliary rings associated with \mathfrak{u} : the factor ring $\mathfrak{o}/\mathfrak{u}$, composed of the residue classes of elements of \mathfrak{o} modulo \mathfrak{u} , and the ring of quotients $\mathfrak{o}_{\mathfrak{u}}$, composed of the fractions whose numerator and denominator belong to \mathfrak{o} , but whose denominators do not belong to \mathfrak{u} . These constructions are of paramount importance in algebraic geometry; if \mathfrak{o} is the ring of a variety V , there corresponds to \mathfrak{u} a subvariety U of V ; $\mathfrak{o}/\mathfrak{u}$ is the ring of U , whereas the ring $\mathfrak{o}_{\mathfrak{u}}$ is the proper algebraic tool to investigate the neighborhood of U with respect to V .

Now, the local theory of algebraic varieties involves the consideration of rings which are not domains of integrity (this, because the completion of a local ring may introduce zero divisors). Let then \mathfrak{o} be any commutative ring with unit element, and let again \mathfrak{u} be a prime ideal in \mathfrak{o} . We may define the factor ring $\mathfrak{o}/\mathfrak{u}$ exactly in the same way as above, but we cannot so easily generalize the notion of the ring of quotients $\mathfrak{o}_{\mathfrak{u}}$. If there exist zero divisors outside \mathfrak{u} , these zero divisors cannot be used as denominators of fractions, which shows that the definition of $\mathfrak{o}_{\mathfrak{u}}$ cannot be extended verbatim. If we

Received by the editors September 4, 1943.

consider only fractions whose denominators are not zero divisors and do not belong to \mathfrak{u} , we obtain a ring \mathfrak{o}' ; but \mathfrak{o}' fails in general to have the essential property of a ring of quotients, namely, of being a local ring in the sense of Krull (that is, the non-units in \mathfrak{o}' will not form an ideal). The object of this note is to construct a ring for which the essential properties of rings of quotients are preserved.

Throughout this paper we shall denote by \mathfrak{o} a Noetherian ring (that is, a ring in which the maximal condition for ideals is satisfied) with a unit element. Generalizing the problem of defining the ring of quotients of a prime ideal, we take any multiplicatively closed¹ subset S of \mathfrak{o} which does not contain 0 (a set is said to be multiplicatively closed if the product of any two elements of the set belongs to the set; if we are concerned with a prime ideal \mathfrak{u} in \mathfrak{o} , we take S to be the complement of \mathfrak{u} in \mathfrak{o}). There exists at least one primary ideal which does not meet S (otherwise, 0 would belong to S as we see at once by representing the zero ideal as an intersection of primary ideals). We shall denote by \mathfrak{s} the intersection of all primary ideals in \mathfrak{o} which do not meet S .

PROPOSITION 1. *Let $\{0\} = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_h$ be an irredundant representation of the zero ideal in \mathfrak{o} as an intersection of primary ideals, and let $\mathfrak{p}_i (1 \leq i \leq h)$ be the associated prime ideal of \mathfrak{q}_i . Assume that $\mathfrak{p}_i \cap S = \emptyset$ for $i \leq g$, but not for $i > g$. Then $\mathfrak{s} = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_g$.*

It is clear that $\mathfrak{s} \subset \mathfrak{q}' = \mathfrak{q}_1 \cap \cdots \cap \mathfrak{q}_g$. Let \mathfrak{v} be any primary ideal which does not meet S ; we shall prove that $\mathfrak{q}' \subset \mathfrak{v}$. Let \mathfrak{q}'' be the ideal $\mathfrak{q}_{g+1} \cap \cdots \cap \mathfrak{q}_h$. We have $\{0\} = \mathfrak{q}' \cap \mathfrak{q}'' = \mathfrak{q}' \mathfrak{q}'' \subset \mathfrak{v}$, whence $\mathfrak{q}' \subset \mathfrak{v} : \mathfrak{q}''$. Let \mathfrak{u} be the associated prime ideal of \mathfrak{v} ; since \mathfrak{v} contains some power of \mathfrak{u} , it follows from the multiplicatively closed character of S that \mathfrak{u} does not meet S . If $i > g$, the ideal \mathfrak{p}_i meets S and is therefore not contained in \mathfrak{u} . It follows² that $\mathfrak{v} : \mathfrak{q}'' = \mathfrak{v}$, whence $\mathfrak{q}' \subset \mathfrak{v}$. Proposition 1 is thereby proved.

LEMMA 1. *Let \mathfrak{p} be a prime ideal in \mathfrak{o} , and let \mathfrak{a} be an ideal contained in \mathfrak{p} . If \mathfrak{q} is an ideal containing \mathfrak{a} , the statements " \mathfrak{q} is primary for \mathfrak{p} " and " $\mathfrak{q}/\mathfrak{a}$ is primary for $\mathfrak{p}/\mathfrak{a}$ " are equivalent.*

¹ It was H. Grell who observed for the first time that, S being any multiplicatively closed set of nonzero divisors in a ring, it is possible to associate with S a ring of quotients, whose elements are the fractions whose denominators belong to S (Cf. H. Grell, *Beziehungen zwischen Ideale verschiedener Ringe*, Math. Ann. vol. 97 (1926) p. 510). For the properties of these rings of quotients, cf. Krull, *Idealtheorie* (Ergebnisse der Mathematik) or my paper *On the theory of local rings*, Ann. of Math. vol. 44 (1943) p. 690.

² Cf. van der Waerden, *Moderne Algebra*, vol. 2, chap. 12, p. 36.

Lemma 1 follows trivially from the definitions.

The zero ideal in $\mathfrak{o}/\mathfrak{s}$ is the intersection of the primary ideals $\mathfrak{q}_1/\mathfrak{s}, \dots, \mathfrak{q}_g/\mathfrak{s}$ whose associated prime ideals are $\mathfrak{p}_1/\mathfrak{s}, \dots, \mathfrak{p}_g/\mathfrak{s}$. Let S^* be the set of the residue classes modulo \mathfrak{s} of the elements of S ; then $S^* \cap (\mathfrak{p}_i/\mathfrak{s}) = \emptyset$ ($1 \leq i \leq g$), which means that no element of S^* is a zero divisor in $\mathfrak{o}/\mathfrak{s}$.³ We may therefore construct the ring of quotients $(\mathfrak{o}/\mathfrak{s})_{S^*}$ of S^* with respect to the ring $\mathfrak{o}/\mathfrak{s}$.

DEFINITION 1. *The ring $(\mathfrak{o}/\mathfrak{s})_{S^*}$ will be called the ring of quotients of S with respect to \mathfrak{o} . This ring will be denoted by \mathfrak{o}_S .*

This definition coincides with the usual one in the case where S does not contain any zero divisor. We shall now prove that the essential properties of rings of quotients in the usual sense still hold in our case.

If \mathfrak{a} is an ideal in \mathfrak{o} , $(\mathfrak{a} + \mathfrak{s})/\mathfrak{s}$ is an ideal in \mathfrak{o}_S which we shall denote symbolically by $\mathfrak{a}\mathfrak{o}_S$ (in spite of the fact that \mathfrak{o} is not in general a subring of \mathfrak{o}_S , so that we cannot multiply elements of \mathfrak{o} by elements of \mathfrak{o}_S). If \mathfrak{b} is any ideal in \mathfrak{o}_S , the set $\mathfrak{b} \cap (\mathfrak{o}/\mathfrak{s})$ may be written in the form $\mathfrak{a}/\mathfrak{s}$, where \mathfrak{a} is an ideal in \mathfrak{o} which contains \mathfrak{s} . We shall denote \mathfrak{a} symbolically by $\mathfrak{b} \cap \mathfrak{o}$ (although $\mathfrak{b} \cap \mathfrak{o}$ is not a set theoretic intersection).

PROPOSITION 2. *If \mathfrak{b} is any ideal in \mathfrak{o}_S , we have $\mathfrak{b} = (\mathfrak{b} \cap \mathfrak{o})\mathfrak{o}_S$.*

Since \mathfrak{o}_S is a ring of quotients of $\mathfrak{o}/\mathfrak{s}$, we have $\mathfrak{b} = (\mathfrak{b} \cap (\mathfrak{o}/\mathfrak{s}))\mathfrak{o}_S$.¹ Proposition 2 follows immediately from this formula.

PROPOSITION 3. *Let \mathfrak{p} be a prime ideal in \mathfrak{o} , and let \mathfrak{q} be primary for \mathfrak{p} . If \mathfrak{q} meets S , we have $\mathfrak{p}\mathfrak{o}_S = \mathfrak{q}\mathfrak{o}_S = \mathfrak{o}_S$. If \mathfrak{p} does not meet S , $\mathfrak{p}\mathfrak{o}_S$ is prime, $\mathfrak{q}\mathfrak{o}_S$ is primary for $\mathfrak{p}\mathfrak{o}_S$ and $\mathfrak{p}\mathfrak{o}_S \cap \mathfrak{o} = \mathfrak{p}$, $\mathfrak{q}\mathfrak{o}_S \cap \mathfrak{o} = \mathfrak{q}$.*

If \mathfrak{q} meets S , $\mathfrak{q} + \mathfrak{s}/\mathfrak{s}$ meets S^* , whence $\mathfrak{p}\mathfrak{o}_S = \mathfrak{q}\mathfrak{o}_S = \mathfrak{o}_S$. If \mathfrak{p} does not meet S , the same holds for \mathfrak{q} , whence $\mathfrak{s} \subset \mathfrak{q} \subset \mathfrak{p}$. By Lemma 1, $\mathfrak{p}/\mathfrak{s}$ is prime and $\mathfrak{q}/\mathfrak{s}$ is primary for $\mathfrak{p}/\mathfrak{s}$. Furthermore, $\mathfrak{p}/\mathfrak{s}$ does not meet S^* . Proposition 3 follows therefore from the corresponding proposition which is known to hold for ordinary rings of quotients.¹ We see also that, if \mathfrak{p} does not meet \mathfrak{s} , the formula $\mathfrak{q} \leftrightarrow \mathfrak{q}\mathfrak{o}_S$ establishes a one-to-one inclusion preserving correspondence between the primary ideals for \mathfrak{p} in \mathfrak{o} and the primary ideals for $\mathfrak{p}\mathfrak{o}_S$ in \mathfrak{o}_S .

PROPOSITION 4. *Let $\mathfrak{a} = \mathfrak{v}_1 \cap \dots \cap \mathfrak{v}_r$ be an irredundant representation of an ideal \mathfrak{a} in \mathfrak{o} as an intersection of primary ideals. Let \mathfrak{u}_i be the associated prime ideal of \mathfrak{v}_i , and assume that \mathfrak{u}_i meets S for $i > s$ but not for $i \leq s$. Then we have $\mathfrak{a}\mathfrak{o}_S = \mathfrak{v}_1\mathfrak{o}_S \cap \dots \cap \mathfrak{v}_r\mathfrak{o}_S$, and this is an irredundant representation of $\mathfrak{a}\mathfrak{o}_S$ as intersection of primary ideals in \mathfrak{o}_S .*

³ If $s^* \in S^*$, we have $\{0\} :_{s^*} (\mathfrak{o}/\mathfrak{s}) = \{0\}$ by the result quoted in footnote 2.

It is obvious that $a\mathfrak{o}_S \subset v_1\mathfrak{o}_S \cap \dots \cap v_s\mathfrak{o}_S$. Let conversely a be any element of $v_1\mathfrak{o}_S \cap \dots \cap v_s\mathfrak{o}_S$. We know³ that $v_1\mathfrak{o}_S \cap \dots \cap v_s\mathfrak{o}_S$ is equal to $(v_1/\mathfrak{s} \cap \dots \cap v_s/\mathfrak{s})\mathfrak{o}_S$; it follows that a may be written in the form b^*/c^* , where $b^* \in v_1/\mathfrak{s} \cap \dots \cap v_s/\mathfrak{s}$ and $c^* \in S^*$. If $i > s$, the ideal u_i has an element u_i in common with S ; if m is large enough, we have $u = (\prod_{i=s+1}^r u_i)^m \in v_{s+1} \cap \dots \cap v_r$. Since $u_i \in S$ ($s+1 \leq i \leq r$), we have $u \in S$, whence $u^* \in S^*$, if u^* is the residue class of u modulo \mathfrak{s} . We may write $a = (u^*b^*)/(u^*c^*)$, $u^*c^* \in S^*$. Let b be any element of the residue class b^* modulo \mathfrak{s} ; since $b^* \in v_i/\mathfrak{s}$ ($1 \leq i \leq s$), we have $b \in v_1 \cap \dots \cap v_s$, whence $ub \in a$ and $u^*b^* \in a + \mathfrak{s}/\mathfrak{s}$, $a \in a\mathfrak{o}_S$. It is clear that none of the ideals v_i/\mathfrak{s} contains the intersection of the others; making use of a known result¹ for ordinary rings of quotients, it follows that the representation $a\mathfrak{o}_S = v_1\mathfrak{o}_S \cap \dots \cap v_s\mathfrak{o}_S$ is irredundant.

We shall now consider more specifically the case where S is the complement of a prime ideal u . The ring \mathfrak{o}_S will then also be denoted by \mathfrak{o}_u . In that case, the ideal \mathfrak{s} coincides with the intersection of all primary ideals for u . In fact, the set S^* is clearly the complement of u/\mathfrak{s} with respect to $\mathfrak{o}/\mathfrak{s}$; the ring \mathfrak{o}_S is the ring of quotients (in the ordinary sense) of the prime ideal u/\mathfrak{s} with respect to $\mathfrak{o}/\mathfrak{s}$. It follows that $u\mathfrak{o}_S$ is the ideal of non-units in \mathfrak{o}_S , whence $\bigcap_{n=1}^\infty (u/\mathfrak{s})^n \mathfrak{o}_S = \{0\}$.⁴ For every n , the ideal $(u/\mathfrak{s})^n \mathfrak{o}_S \cap \mathfrak{o}/\mathfrak{s}$ is a primary ideal for u/\mathfrak{s} in $\mathfrak{o}/\mathfrak{s}$; it follows that the intersection of all primary ideals for u/\mathfrak{s} is the zero ideal in $\mathfrak{o}/\mathfrak{s}$. Our assertion then follows from Lemma 1.⁵ At the same time, we see that \mathfrak{o}_u is a local ring in the sense of Krull.

Assume now that \mathfrak{o} is a semi-local ring⁶ and that u is one of the maximal prime ideals in \mathfrak{o} . Let $\bar{\mathfrak{o}}$ be the completion of \mathfrak{o} ; there corresponds to u an idempotent ϵ in $\bar{\mathfrak{o}}$. We shall prove the following results:

PROPOSITION 5. *The ring $\mathfrak{o}/\mathfrak{s}$ is isomorphic with the subring $\mathfrak{o}\epsilon$ of the ring $\bar{\mathfrak{o}}\epsilon$. This isomorphism may be extended to an isomorphism of the completion of \mathfrak{o}_u with $\bar{\mathfrak{o}}\epsilon$.*

The first statement will be proved if we show that \mathfrak{s} coincides with the set of elements $x \in \mathfrak{o}$ which satisfy the condition $x\epsilon = 0$. If x is any

⁴ Cf. Krull, *Dimensionstheorie in Stellenringen*, J. Reine Angew. Math. vol. 179 (1938) p. 204 or my paper quoted in footnote 1.

⁵ This result, together with Proposition 1 above, yields a proof of a theorem of Krull; cf. Krull, *Primidealketten in allgemeinen Ringbereichen*, Sitzungsberichte der Heidelberger Akademie, 1928, p. 7.

⁶ A semi-local ring is a Noetherian ring (that is, the maximal chain condition holds in the ring) in which there exist only a finite number of maximal prime ideals. For the proofs of the results on semi-local rings which are used in this paper, cf. my paper quoted in footnote 1.

element of \mathfrak{o} , we may write $x = x\epsilon + x(1 - \epsilon)$, and we know that $1 - \epsilon \in u^n \bar{\mathfrak{d}}$ for every n . If $x\epsilon = 0$, we have $x \in u^n \bar{\mathfrak{d}} \cap \mathfrak{o} = u^n$ for every n ; in particular, x belongs to every primary ideal for u , whence $x \in \mathfrak{s}$. If $x \in \mathfrak{s}$, we have $x \in u^n$ for every n (u^n is primary because u is a *maximal* prime ideal), whence $x\epsilon \in u^n \bar{\mathfrak{d}}\epsilon$. Since $u\bar{\mathfrak{d}}\epsilon$ is the ideal of non-units in $\bar{\mathfrak{d}}\epsilon$, it follows that $x\epsilon = 0$.

If we identify $\mathfrak{o}/\mathfrak{s}$ with $\mathfrak{o}\epsilon$, every element of S^* is a unit in $\bar{\mathfrak{d}}\epsilon$. In fact, if $y \in S$, we have $y = y\epsilon + y(1 - \epsilon)$, $y(1 - \epsilon) \in u\bar{\mathfrak{d}}$. If we had $y \in u\bar{\mathfrak{d}}$, we would have $y \in u\bar{\mathfrak{d}} \cap \mathfrak{o} = u$, which is not the case. It follows that $\bar{\mathfrak{d}}\epsilon$ contains the ring \mathfrak{o}_u . The ring $\bar{\mathfrak{d}}\epsilon$ is a complete ring with $u\bar{\mathfrak{d}}\epsilon$ as unique maximal prime ideal; it is clear that $\mathfrak{o}\epsilon$ (and, a fortiori, \mathfrak{o}_u) is dense in $\bar{\mathfrak{d}}\epsilon$. In order to prove that $\bar{\mathfrak{d}}\epsilon$ is the completion of \mathfrak{o}_u , it is sufficient to prove that \mathfrak{o}_u is topologically a subspace of $\bar{\mathfrak{d}}\epsilon$. We show first that $u^n \bar{\mathfrak{d}}\epsilon \cap \mathfrak{o}\epsilon = u^n \mathfrak{o}\epsilon$ for every n . Let $x\epsilon$ ($x \in \mathfrak{o}$) be an element of $u^n \bar{\mathfrak{d}}\epsilon \cap \mathfrak{o}\epsilon$; we have $x = x\epsilon + x(1 - \epsilon) \in u^n \bar{\mathfrak{d}}$, whence $x \in u^n \bar{\mathfrak{d}} \cap \mathfrak{o} = u^n$, $x\epsilon \in u^n \mathfrak{o}\epsilon$. The ideal $u^n \bar{\mathfrak{d}} \cap \mathfrak{o}_u$ is equal to $((u^n \bar{\mathfrak{d}} \cap \mathfrak{o}_u) \cap \mathfrak{o}) \mathfrak{o}_u = (u^n \mathfrak{o}\epsilon) \mathfrak{o}_u = u^n \mathfrak{o}_u$; Proposition 5 is thereby completely proved.

PRINCETON UNIVERSITY