

156. J. A. Shohat: *Series expansions for the periodic solution of Van der Pol's equation and its frequency for all values of the parameter.*

If the parameter  $\mu$  in Van der Pol's equation  $d^2u/dt^2 - \mu(1-u^2)du/dt + u = 0$  is small, power series expansions for the periodic solution  $u$  (unique, save for time-translation) and its frequency  $\nu$  can be and have been given, say, by Lidstedt's method. In the present paper the author gives (for the first time, he believes) series expansion for  $u$  and  $\nu$ , valid for all values of  $\mu$ —large and small. Numerical computation agrees quite well with known numerical results. (Received March 13, 1944.)

#### GEOMETRY

157. Reinhold Baer: *The fundamental theorems of elementary geometry. An axiomatic analysis.*

It is the object of this paper to evaluate the logical interdependence of certain fundamental theorems in elementary geometry. The paper deals with the theorems asserting the copunctuality of each of the following triplets of lines: medians, altitudes, perpendicular bisectors, and bisectors of angles; and the theorem stating that the locus of the points of equal distance from two different points is a line. The framework of our discussion is provided by a general affine plane in which we introduce just as many further relations as are needed for stating the investigated theorems. (Received March 22, 1944.)

158. P. O. Bell: *A study of surfaces by means of a system of differential equations of the first order.*

The projective differential geometry of a surface in ordinary space is studied by means of tetrads of surfaces whose corresponding points  $x_i$  ( $i=0, 1, 2, 3$ ) are linearly independent. The general homogeneous coordinates of  $x_i$  satisfy a system of equations  $\partial x_i / \partial u^\alpha = C_{h,i\alpha} x_h$ ,  $\alpha=1, 2$ , summed for  $h=0, 1, 2, 3$ . With the points  $x_i$  as vertices of a local reference tetrahedron an algebraic surface  $a_{ij} \dots x^i x^j \dots x^l = 0$  is fixed as  $u^1, u^2$  vary independently, if and only if the coefficients  $a_{ij} \dots$  are proportional to the corresponding components of the covariant derivatives, of the aggregate of these coefficients, with respect to the connection  $C_{h,i\alpha}$ . Such conditions of immovability form the basis for a general theory of envelopes. Tetrads of surfaces are first investigated. The study of a surface  $S_0$  is then undertaken by specializing the general theory. Auxiliary surfaces  $S_1, S_2, S_3$  covariantly determined with respect to  $S_0$  are selected so that the fundamental differential equations are as simple as possible and exhibit desirable properties of symmetry. Some differential invariants are characterized geometrically. When the asymptotic curves are parametric on one of the surfaces one of these invariants becomes the projective linear element and another becomes Fubini's element of projective arc length. (Received April 1, 1944.)

159. S. S. Chern: *Laplace transforms of a class of higher dimensional varieties in a projective space of  $n$  dimensions.*

In a projective space of  $n$  dimensions a class of  $r$ -dimensional varieties is defined, which form a natural generalization of the surfaces sustaining conjugate nets. These varieties are characterized by the property that the asymptotic net is an  $(r-1)$ -parameter linear system of cones whose base cones are linear spaces counted twice. (See E. Cartan, Bull. Soc. Math. France vol. 47 (1919) pp. 125-160.) This geometrical

characterization is in general equivalent to the following: On the variety there exist  $r$  families of curves such that when the tangent  $r$ -plane is displaced along a curve of the family its intersection with a neighboring tangent  $r$ -plane is the tangent  $(r-1)$ -plane formed by the tangents of the other  $r-1$  curves. Basing on this property one can define on each of these  $r$  tangents  $AA_i$ ,  $i=1, \dots, r-1$  points  $A_{ij}$ ,  $j \neq i$ , having the property that when the  $s$ -plane  $AA_{i_1} \dots A_{i_s}$  is displaced along  $AA_j$ ,  $j \neq i_1, \dots, i_s$ , its intersection with a neighboring  $s$ -plane is the  $(s-1)$ -plane  $A_{i_1 j} \dots A_{i_s j}$ . The points  $A_{ij}$  describe varieties of the same type and are naturally defined as the Laplace transforms of the given variety, there being altogether  $r(r-1)$  transforms. Many well known properties of Laplace transforms can be generalized. (Received February 23, 1944.)

160. S. B. Jackson: *Vertices of plane curves.*

A closed curve of class  $C''$ , not a circle, has two vertices by the continuity of the curvature. The present paper seeks to characterize geometrically those curves with exactly two vertices. Let a curve be called normalized if it contains no complete circles, and let a simple closed arc of the curve which is never crossed by the curve be called a simple loop. The following facts are established for any normalized curve  $C$  with two vertices: (a)  $C$  may be divided into two simple arcs; (b) all double points are simple; (c)  $C$  contains exactly two simple loops, one containing each vertex; (d) none of the plane regions bounded by  $C$  are bounded always in the same sense except those regions bounded by the loops; (e) at any points of tangency the directed tangents coincide. For a curve which is not normalized these results are modified slightly. Two familiar theorems regarding the number of vertices on an oval are generalized to any simple closed curve. The methods employed are entirely elementary, extensive use being made of the invariance of vertices under direct circular transformations. (Received February 4, 1944.)

161. J. E. Wilkins: *The contact of a cubic surface with a ruled surface.*

It is shown that there exist  $\infty^1$  cubic surfaces having contact of order 5 with a non-developable ruled surface. If there is any cubic surface having contact of order 6 with a nondevelopable ruled surface, then the surface is itself a cubic surface. In order to obtain these results, there are first derived power series expansions for a nondevelopable ruled surface to terms of the sixth degree. Similar investigations are made for developable surfaces. (Received April 1, 1944.)

## STATISTICS AND PROBABILITY

162. C. W. Churchman and Benjamin Epstein: *Estimates of error in parallel experiments.*

It is common in many types of tests to have not only a random error from test to test due to a large number of unallocable causes, but it is also possible to have systematic errors present. It is because of this possibility that one tests not only samples of the unknown, but also control samples. The purpose of using control samples is two-fold—(a) to find out whether or not abnormal experimental conditions exist during the test and (b) to establish tentatively a level for the particular test under consideration. It is shown that a statistic can be found which gives the most efficient estimate of the corrections to be applied to the unknown under test for a variety of experimental conditions. It is further shown that this statistic must be a linear func-