

For by Proposition 4, $FA = A$ for every A and $A(A^{-1}F) = EF = F$.

Let S_1, \dots, S_n be n statements. Let A_i be the statement: "All the preceding statements are annulled but S_i is true." It is interesting to note that the statements A_i form an idempotent (I, r) system.

OHIO STATE UNIVERSITY

A CONJECTURE IN ELEMENTARY NUMBER THEORY

LEON ALAOGU AND PAUL ERDÖS

A well known conjecture of Catalan states that *if $f(n)$ is the sum of all divisors of n except n , then the sequence of iterates of $f(n)$ is either eventually periodic or ends at 1*. It not only seems impossible to prove this, but it is also very difficult to verify.¹

Another conjecture of Poulet,² which appears equally difficult to prove, has the doubtful merit that it is easy to verify. Let $\sigma(n)$ be the sum of all divisors of n , and let $\phi(n)$ be Euler's function. Then *for any integer n the sequence*

$$f_0(n) = n, \quad f_{2k+1}(n) = \sigma(f_{2k}(n)), \quad f_{2k}(n) = \phi(f_{2k-1}(n))$$

is eventually periodic.

We have verified this conjecture to $n = 10000$ (extending Poulet's verification) by using Glaisher's tables.³ The checking was facilitated by the following observation: if the conjecture is to be checked for all $n < x$, it is enough to find a member of the sequence other than the first which is less than x .

The longest cycle found was in the sequence $f_k(9216)$. It starts with $f_0(9216)$, and is: 34560, 122640, 27648, 81800, 30976, 67963, 54432, 183456, 48384, 163520, 55296, 163800, 34560. However our method of checking does not show that this is the largest cycle up to 10000, and in fact Poulet found that $f_k(1800)$ has the same length 12.

As a rule $\phi(\sigma(n))$ is less than n . In fact, it can be shown that *for every $\epsilon > 0$, $\phi(\sigma(n)) < \epsilon n$, except for a set of density 0*. The proof follows from the following two observations:

Received by the editors May 18, 1944.

¹ L. E. Dickson, *Theorems and tables on the sum of the divisors of a number*, Quart. J. Math. vol. 44 (1913) pp. 264-296, and P. Poulet, *La Chasse aux Nombres*, vol. 1, pp. 68-72, and vol. 2, p. 188.

² P. Poulet, *Nouvelles suites arithmétiques*, Sphinx vol. 2 (1932) pp. 53-54.

³ J. W. L. Glaisher, *Number-divisor tables*, British Association for the Advancement of Science, Mathematical Tables, vol. 8.

(1) For a given prime p , the set of all n such that $\sigma(n) \equiv 0 \pmod{p}$ is of density 1.

The set of all integers not divisible by any prime q of the form $px - 1$ is of density zero, since $\sum_q 1/q$ diverges. Hence the set of all integers divisible by a prime $q > N$ of this type is of density 1. But the set of all integers divisible by q^2 , $q > N$, is of density less than $\sum_{q > N} 1/q^2 = o(1)$. Therefore, if x is large, the number of n less than x such that $\sigma(n) \equiv 0 \pmod{p}$ exceeds $(1 - \epsilon)x$.

(2) Except for ϵx integers n less than x , $\sigma(n) < c(\epsilon)n$.

This follows from the fact that $\sum_{n < x} \sigma(n) \sim \pi^2 n^2 / 12$.

Choose p so that $\prod_{q \leq p} (1 - 1/q) < \delta/c(\epsilon)$. Then, if x is sufficiently large, all except $\eta x + \epsilon x$ integers n less than x have $\sigma(n) < c(\epsilon)n$, $\sigma(n) \equiv 0 \pmod{q}$ for all $q \leq p$. But, with these exceptions, $\phi[\sigma(n)] < \delta n$, which completes the proof, since η and ϵ are arbitrary.

In much the same way it can be shown that for every $c > 0$, $\sigma[\phi(n)] > cn$ except for a set of density zero.

Actually, much more can be shown. Except for a set of density zero, $e^\gamma \phi[\sigma(n)] \log \log \log n \sim \sigma(n)$, and $e^{-\gamma} \sigma[\phi(n)] / \log \log \log n \sim \phi(n)$, where γ is Euler's constant. The proof is suppressed, but it might be noted that the reason for this result is that, for almost all n , $\phi(n)$ and $\sigma(n)$ are both divisible by all primes less than $(\log \log n)^{1-\epsilon}$, and by relatively few primes greater than $(\log \log n)^{1+\epsilon}$.

There exist numbers for which $\phi(\sigma(n)) = n$. Up to 2500 these numbers are 1, 2, 8, 12, 128, 240, 720; while two further solutions are 2^{15} and 2^{31} . Poulet gives many others; we do not know whether there are infinitely many solutions.

We state two further conjectures:

(a) Form the sequence $\sigma(n), \sigma(\sigma(n)), \phi(\sigma(\sigma(n))), \sigma(\phi(\sigma(\sigma(n))))$ in which the functions are successively applied in the order $\sigma, \sigma, \phi, \sigma, \sigma, \phi, \sigma, \sigma, \phi, \dots$. This sequence seems to tend to infinity if n is large enough.

(b) On the other hand, the sequence $\phi(n), \phi(\phi(n)), \sigma(\phi(\phi(n))), \dots$, in which the order is $\phi, \phi, \sigma, \phi, \phi, \sigma, \phi, \phi, \sigma, \dots$, seems to converge to 1, for all n .

Obviously many more such conjectures can be formulated.