

SOME REMARKS ABOUT ADDITIVE AND MULTIPLICATIVE FUNCTIONS

P. ERDÖS

The present paper contains some results about the classical multiplicative functions $\phi(n)$, $\sigma(n)$ and also about general additive and multiplicative functions.

(1) It is well known that $n/\phi(n)$ and $\sigma(n)/n$ have a distribution function.¹ Denote these functions by $f_1(x)$ and $f_2(x)$. ($f_1(x)$ denotes the density of integers for which $n/\phi(n) \leq x$.) It is known that both $f_1(x)$ and $f_2(x)$ are strictly increasing and purely singular.¹ We propose to investigate $f_1(x)$ and $f_2(x)$; we shall give details only in case of $f_1(x)$. First we prove the following theorem.

THEOREM 1. *We have for every ϵ and sufficiently large x*

$$(1) \quad \exp(-\exp[(1+\epsilon)ax]) < 1 - f_1(x) < \exp(-\exp[(1-\epsilon)ax])$$

where $a = \exp(-\gamma)$, γ Euler's constant.

We shall prove a stronger result. Put $A_r = \prod_{i=1}^r p_i$, p_i consecutive primes. Define A_k by $A_k/\phi(A_k) \geq x > A_{k-1}/\phi(A_{k-1})$. Then we have

$$(2) \quad 1/A_k < 1 - f_1(x) < 1/A_k^{1-\epsilon}.$$

First of all it is easy to see that Theorem 1 follows from (2), since from the prime number theorem we easily obtain that $\log \log A_k = (1+o(1))ax$, which shows that (1) follows from (2).

(2) means that the density of integers with $\phi(n) \leq (1/x)n$ is between $1/A_k$ and $1/A_k^{1-\epsilon}$.

We evidently have for every $n \equiv 0 \pmod{A_k}$, $n/\phi(n) \geq x$, which proves

$$1/A_k \leq 1 - f_1(x).$$

To get rid of the equality sign, it will be sufficient to observe that there exist integers u with $u/\phi(u) \geq x$, $(u, A_k) = 1$, and that the density of the integers $n \equiv 0 \pmod{u}$, $n \not\equiv 0 \pmod{A_k}$ is positive. This proves the first part of (2). The proof of the second part will be much harder. We split the integers satisfying $n/\phi(n) \geq x$ into two classes. In the first class are the integers which have more than $[(1-\epsilon_1)k] = r$ prime factors not greater than Bp_k , where $B = B(\epsilon_1)$ is a large number. In

Received by the editors June 22, 1945, and, in revised form, January 28, 1946.

¹ These results are due to Schönberg and Davenport. For a more general result see P. Erdős, J. London Math. Soc. vol. 13 (1938) pp. 119-127.

the second class are the other integers satisfying $n/\phi(n) \geq x$. It is easy to see that the number of integers of the first class does not exceed

$$(3) \quad 2^{\pi(Bp_k)}/A_r = 2^{o(p_k)}/A_r < 1/A_k^{1-\epsilon}$$

since $\pi(Bp_k) = o(p_k)$ ($\pi(x)$ denotes the number of primes not greater than x), and from the prime number theorem $\log A_r > (1 - \epsilon)p_k$ if ϵ_1 is small.

Let now n be any integer of the second class. A simple argument shows that

$$\prod_{p|n} \left(1 - \frac{1}{p}\right) < \prod_{t=r+1}^{k-1} \left(1 - \frac{1}{p_t}\right) < 1 - \frac{c_1 \epsilon_1}{\log p_k}.$$

The prime indicates that the product is extended over the $p > Bp_k$. The first inequality follows from the definition of A_k , and from the fact that n is of the second class, the second inequality follows from the prime number theorem. Thus we have

$$(4) \quad \sum'_{p|n} \frac{1}{p} > \frac{c_1 \epsilon_1}{\log p_k}.$$

Denote now by J_t the interval $(B^t p_k, B^{t+1} p_k)$, $t = 1, 2, \dots$. It follows from (4) that for every integer of the second class there exists some t such that

$$(5) \quad \sum_{p|n} \frac{1}{p} > c_1 \frac{\epsilon_1}{2^t \log p_k}$$

where in \sum_t the summation is extended over the primes in J_t . Thus for some t , n must divide more than

$$(6) \quad c_1 \epsilon_1 (B^t/2^t) (p_k/\log p_k) = B_t$$

primes in J_t . The density of the integers satisfying (6), that is, the density of the integers of the second class, is less than

$$(7) \quad \sum_{t=1}^{\infty} \left(\sum_{p \text{ in } J_t} \frac{1}{p} \right)^{B_t} / [B_t]! < \frac{1}{[B_t]!} < e^{-2p_k} < \frac{1}{A_k}$$

that is, $\sum_{p \text{ in } J_t} 1/p < 1$ for large enough k (B is independent of k), if $B = B(\epsilon_1)$ is large enough. Theorem 1 now follows from (3) and (7).

From Theorem 1 we easily obtain that

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n=1}^x \exp(\phi(n))$$

exists. In fact we can also prove that for $\alpha < a$

$$\lim_{x \rightarrow \infty} \frac{1}{x} \sum_{n=1}^x \exp(\exp(\phi(n)))$$

exists. For $\alpha > a$ the limit is infinite.

THEOREM 2.

$$1/A_k^{1+\epsilon} < 1 - f_1(x) < 1/A_k^{1-\epsilon}.$$

We omit the proof since it is very similar to that of Theorem 1.

THEOREM 3. Let $\epsilon \rightarrow 0$, then

$$f_1(1 + \epsilon) = (1 + o(1))a/\log \epsilon^{-1}, \quad f_2(1 + \epsilon) = (1 + o(1))a/\log \epsilon^{-1}.$$

We prove only the first statement since the proof of the second is essentially the same. Let n be an integer with $n/\phi(n) \leq 1 + \epsilon$. Clearly n does not divide any prime $p < (1 - (1 + \epsilon)^{-1})^{-1} = \epsilon^{-1} + O(1)$. Thus

$$(8) \quad f_1(1 + \epsilon) < (1 + o(1))a/\log \epsilon^{-1}.$$

Denote by J_t the interval

$$(4^{t-1}(1 - (1 + \epsilon)^{-1})^{-1}, 4^t(1 - (1 + \epsilon)^{-1})^{-1}).$$

If an integer $n \not\equiv 0 \pmod{p_i}$, $p_i < (1 - (1 + \epsilon)^{-1})^{-1}$, does not satisfy $n/\phi(n) \leq 1 + \epsilon$, then a simple computation shows that for some t it must have at least t prime factors in J_t . Thus the number of these integers does not exceed

$$(1 + o(1)) \frac{a}{\log \epsilon^{-1}} \sum_{t=1}^{\infty} \left(\sum_{p \text{ in } J_t} \frac{1}{p} \right)^t / t! = o(a/\log \epsilon^{-1}),$$

which together with (8) proves Theorem 3.

It follows from Theorem 3 that $f_1'(1) = \infty$. It would be easy to show that $f_1'(n/\phi(n)) = \infty$ for every n .

Denote by f_1^α and f_2^α the distribution functions of

$$\prod_{p|n} \left(1 - \frac{1}{p}\right)^{-\alpha} \quad \text{and} \quad \sum_{d|n} \frac{1}{d^\alpha}, \quad \alpha > 0.$$

THEOREM 4.

$$f_1^{(\alpha)}(1 + \epsilon) = (1 + o(1)) \frac{a\alpha}{\log \epsilon^{-1}}, \quad f_2^{(\alpha)} = (1 + o(1)) \frac{a\alpha}{\log \epsilon^{-1}}.$$

We omit the proof since it is very similar to that of Theorem 3.

Let us denote by $F_\alpha(x)$, $\alpha > 0$, the distribution function of $\prod_{p|n} (1 - 1/\log p^\alpha)^{-1}$, $\alpha > 0$.

THEOREM 5.

$$F_1(1 + \epsilon) = (1 + o(1))b\epsilon,$$

that is, $F_1'(1) = b$. Also $F_\alpha'(1) = 0$ for $\alpha < 1$ and $F_1'(1) = \infty$ for $\alpha > 1$.

We do not give the details of the proof since it would be long and similar to that of Theorem 3. We just make the following remarks: If n satisfies

$$\sum_{p|n} \frac{1}{\log p} \leq 1 + \epsilon$$

then n does not divide any prime $p \leq \exp(1/\epsilon)$. Thus $F_1'(1 + \epsilon) \leq (1 + o(1))a\epsilon$. But here (unlike in Theorem 3) we have $F_1(1 + \epsilon) = (1 + o(1))b$, $b < a$. We obtain analogous results if we consider the additive function $\sum_{p|n} 1/\log p$. It is possible that $F_1'(x)$ exists for every $1 \leq x$, but this we can not prove.

(2) The following results are well known:

$$\sum_{m=1}^x \frac{\phi(m)}{m} = (1 + o(1)) \frac{6}{\pi^2} x, \quad \sum_{m=1}^x \frac{\sigma(m)}{m} = (1 + o(1)) \frac{\pi^2}{6} x.$$

The density of integers for which $\sigma(n+1)/(n+1) > \sigma(n)/n$ is $1/2$, also the density of integers for which $\phi(n+1)/(n+1) > \phi(n)/n$ is $1/2$.² Now we prove the following theorem.

THEOREM 6. Let $g(n)/\log \log \log n \rightarrow \infty$. Then we have

$$(i) \quad \sum_{m=n}^{n+g(n)} \frac{\phi(m)}{m} = (1 + o(1)) \frac{6}{\pi^2} g(n).$$

(ii) The number of integers m in $(n, n+g(n))$ which satisfy $\phi(m+1)/(m+1) > \phi(m)/m$ equals $(1 + o(1))g(n)/2$.

(iii) The number of integers m in $(n, n+g(n))$ which satisfy $m/\phi(m) \leq c$ equals $(1 + o(1))g(n)f_1(c)$. In other words the distribution function of $\phi(m)/m$ in $(n, n+g(n))$ is the same as the distribution function of $\phi(m)/m$.

All these results are best possible; they become false if for infinitely many n , $g(n) < c \log \log \log n$.

We prove only (i); the proof of (ii) and (iii) are similar. Let $A = A(n)$ tend to infinity sufficiently slowly. Put

$$\frac{\phi(m)}{m} = D_1(m)D_2(m),$$

² P. Erdős, Proc. Cambridge Philos. Soc. vol. 32 (1936) pp. 530-540.

where

$$D_1(m) = \prod'_{p|m} \left(1 - \frac{1}{p}\right), \quad D_2(m) = \prod''_{p|m} \left(1 - \frac{1}{p}\right).$$

The prime indicates that $p \leq A$, the two primes that $p > A$. We evidently have

$$(9) \quad \sum_{m=n}^{n+g(n)} \frac{\phi(m)}{m} < \sum_{m=n}^{n+g(n)} D_1(m) = \sum_a''' \left(\frac{g(n)}{d}\right) \frac{\mu(d)}{d} \\ = (1 + o(1))g(n) \prod_{p \leq A} \left(1 - \frac{1}{p^2}\right) = (1 + o(1)) \frac{\pi^2}{6} g(n)$$

where the three primes indicate that the prime factors of d are not greater than A , and $(g(n)/d)$ denotes the number of multiples of d in $(n, n+g(n))$. Now we show that for sufficiently large A the number of integers in $(n, n+g(n))$ which satisfy

$$(10) \quad D_2(m) < 1 - \epsilon$$

is $o(g(n))$. It will be sufficient to show that

$$(11) \quad \prod_m D_2(m) > (1 - \eta)^{o(n)}$$

for every $\eta > 0$, the product over m runs in $(n, n+g(n))$. We evidently have

$$\prod_m D_2(m) > \prod_1 \left(1 - \frac{1}{p}\right)^{2g(n)/p-1} \prod_2 \left(1 - \frac{1}{p}\right)$$

where, in \prod_1 , $A < p \leq g(n)$, and in \prod_2 , p runs through the prime factors greater than $g(n)$ of $n(n+1) \cdots (n+g(n))$. Clearly

$$\prod_1 > \prod_{p>A} \left(1 - \frac{c}{p^2}\right)^{o(n)} > (1 - \eta_1)^{o(n)}.$$

From the prime number theorem we have $\prod_{p \leq 2y} p < e^{2y}$. Thus

$$\prod_2 > \prod_{p \leq 2y} \left(1 - \frac{1}{p}\right) > \frac{c_1}{\log y}$$

where $y = \log [n(n+1) \cdots (n+g(n))]$. Hence using $g(n)/\log \log \log n \rightarrow \infty$, we obtain by a simple calculation that

$$\prod_2 > (1 - \eta_2)^{o(n)}$$

which proves (11) and therefore (10). From (9) and (10) we obtain by a simple argument that

$$(12) \quad \sum_{m=n}^{n+o(n)} \frac{\phi(m)}{m} > (1 - o(1)) \sum_{m=n}^{n+o(n)} D_1(m) = (1 + o(1))g(n) \frac{\pi^2}{6}.$$

(i) now follows from 9 and (12).³

Now we are going to prove that (i) is best possible. Put $g(N) = c \log \log \log N$, $n/2 < N < n$. Further let A_1, A_2, \dots, A_r , $r = [2^{-1} \log \log \log n]$ be relatively prime integers all of whose prime factors are less than $2^{-1} \log n$ and for which

$$1/4 < \phi(A_i)/A_i < 1/2, \quad i = 1, 2, \dots, r.$$

This is obviously possible since

$$\prod_{p < (\log n)/2} \left(1 - \frac{1}{p}\right) < \frac{c}{\log \log n} < \left(\frac{1}{4}\right)^{(\log \log \log n)/2}.$$

Now choose $n/2 < N < n$ so that $N + j \equiv 0 \pmod{A_i}, j \leq r$. This is possible since by the prime number theorem $A_1 \cdot A_2 \cdot \dots \cdot A_r < n/2$. (In all cases where we refer to the prime number theorem a more elementary result would be sufficient.) Clearly

$$\sum_{m=N+1}^{N+(\log \log \log n)/2} \frac{\phi(m)}{m} < \frac{\log \log \log n}{4}.$$

From (9) we have

$$(13) \quad \sum_{N+(\log \log \log n)/2}^{N+o(N)} \frac{\phi(m)}{m} < (1 + o(1)) \frac{6}{\pi^2} \left(g(N) - \frac{\log \log \log n}{2}\right).$$

Thus finally from (10) and (11) we obtain by a simple calculation

$$\sum_{m=N}^{N+o(N)} \frac{\phi(m)}{m} < (1 - c) \frac{6}{\pi^2} g(N),$$

which shows that (i) is best possible.⁴

THEOREM 7. *Let $g_1(n)/\log \log n \rightarrow \infty$. Then we have*

$$(i) \quad \sum_{m=n}^{n+o_1(n)} \frac{\sigma(m)}{m} = (1 + o(1)) \frac{\pi^2}{6} g_1(n).$$

(ii) *Let $g_2(n)/\log \log \log n \rightarrow \infty$. The number of integers m in $(n, n + g_2(n))$ which satisfy $\sigma(n+1)/(n+1) > \sigma(n)/n$ equals $(1 + o(1)) \cdot g(n)/2$.*

³ This proof is similar to a proof in P. Erdős, J. London Math. Soc. vol. 10 (1935) pp. 128-131.

⁴ This proof is similar to a proof of Chowla and Pillai, J. London Math. Soc. vol. 5 (1930) pp. 95-101.

(iii) *The number of integers m in $(n, n+g(n))$ which satisfy $\sigma(m)/m < c$ equals $(1+o(1)) g(n) f_2(c)$.⁵ All these results are best possible.*

We omit the proof of Theorem 7, since it is similar to that of Theorem 6. We must allow $g_1(n)/\log \log n \rightarrow \infty$, since it is well known that for some $m \leq n$, $\sigma(m) > c \log \log n$ (for example, $m = \prod_{p < (\log n)^{1/2}} p$).

Let $f(n) \leq 1$ and $F(n) \geq 1$ be multiplicative functions with

$$\sum_p \frac{1 - f(p)}{p} < \infty \quad \text{and} \quad \sum_p \frac{F(p) - 1}{p} < \infty.$$

Then we have:

THEOREM 8. *Let $A = A(n)$ tend to infinity arbitrarily slowly, then*

$$\frac{1}{A} \sum_{m=n}^{n+A} f(m) < (1 + o(1)) \frac{1}{n} \sum_{m=1}^n f(m)$$

and

$$\frac{1}{A} \sum_{m=n}^{n+A} F(m) > (1 + o(1)) \frac{1}{n} \sum_{m=1}^n F(m).$$

The proof is quite trivial; it is similar to that of (9). It can be shown that $\lim (1/n) \sum_{m=1}^n f(m)$ and $\lim (1/n) \sum_{m=1}^n F(m)$ exist.

Denote by $V(n)$ the number of prime factors of n and by $d(n)$ the number of divisors of n . We can prove analogs to Theorem 6 for these functions. But the results are very unsatisfactory since for $v(n)$ we have to choose $g(n) = n^{\epsilon/\log \log n}$ and for $d(n)$, $g(n) = n^c$ for some suitable c . These results are probably very far from best possible.

(3) Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$, $p_1^{\alpha_1} < p_2^{\alpha_2} < \cdots < p_k^{\alpha_k}$. Put $(p_i^{\alpha_i})^{b_i} = p_i^{\alpha_i + 1}$. We prove the following theorem.

THEOREM 9. *Let $1 < x$, then for almost all n the number of b 's greater than x equals*

$$x^{-1} \log \log n + o(\log \log n).$$

REMARK. We immediately obtain that every interval $(x, x + \epsilon)$ contains $(1 + o(1)) (\epsilon/x(x + \epsilon)) \log \log n$ b 's.

We are going to give only an outline of the proof. First of all we can assume that all the α 's are 1, since for large r the number of integers not greater than n for which r or more of the α 's is greater than 1 is less than ϵn , since the number of these integers is clearly less than

$$\left(\sum_p \frac{1}{p^2} \right)^r / r! < \epsilon n.$$

⁵ This result has been stated previously, see footnote 4.

Denote by $F(n)$ the number of prime factors p of n such that no prime q in (p, p^x) divides n . $F(n)$ is thus the number of b 's not less than x . We have

$$(14) \quad \sum_{m=1}^n F(m) = \frac{1}{x} \log \log n + o(\log \log n).$$

We now give a sketch of the proof. Clearly

$$\sum_{m=1}^n F(m) = \sum_p f_p(n)$$

where $f_p(n)$ denotes the number of integers $m \leq n$, with $m \equiv 0 \pmod{p}$ and $m \not\equiv 0 \pmod{q}$, $p < q < p^x$. It is easy to see that for $p < n^\epsilon$

$$f_p(n) = (1 + o(1))n/px \quad (p \text{ large}).$$

Also for all p

$$f_p(n) \leq n/p.$$

Thus

$$\begin{aligned} \sum_{m=1}^n F(m) &= \sum_{p \leq n^\epsilon} \frac{n}{px} + O \sum_{n^\epsilon < p < n} \frac{n}{p} + o(\log \log n) \\ &= (1 + o(1)) \frac{\log \log n}{x}, \end{aligned}$$

which proves (14). Now we have to show that

$$F(m) = (1 + o(1))(\log \log n)/x$$

for almost all $m \leq n$. We use Turán's method.⁶ We have

$$\begin{aligned} \sum_{m=1}^n \left(F(m) - \frac{1}{x} \log \log n \right)^2 \\ = \sum_{m=1}^n F^2(m) - \frac{2}{x} \log \log n \sum_{m=1}^n F(m) + n \left(\frac{\log \log n}{x} \right)^2. \end{aligned}$$

Now

$$(15) \quad \sum_{m=1}^n F^2(m) = (1 + o(1))n \left(\frac{\log \log n}{x} \right)^2.$$

We omit the proof of (15), it is similar to the proof of (14). Thus

$$\sum_{m=1}^n \left(F(m) - \frac{1}{x} \log \log n \right)^2 = o(n(\log \log n)^2)$$

which proves Theorem 9.

⁶ P. Turán, J. London Math. Soc. vol. 9 (1934) pp. 274-276.

THEOREM 10. *For almost all n we have*

$$\sum_{p_i|n} b_i = (1 + o(1)) \log \log n \log \log \log n.$$

THEOREM 11. *Let $1 < x$ be any number. For almost all n there exist intervals (m, m^x) , $m^x \leq n$, such that for every $m \leq y \leq m^x$, $n \not\equiv 0 \pmod{y}$.*

We omit the proofs of Theorems 10 and 11. They are similar to that of Theorem 9.

For some time I have not been able to decide the following question: Is it true that almost all integers n have divisors d_1 and d_2 , such that $d_1 < d_2 < 2d_1$.

(4) Let $f(n)$ be an additive function which has a distribution function. Then it is well known that⁷

$$(16) \quad \sum_p \frac{f(p)'}{p} < \infty, \quad \sum_p \frac{(f(p)')^2}{p} < \infty,$$

$f(p)' = f(p)$ if $|f(p)| \leq 1$ and $f(p)' = 1$ if $|f(p)| > 1$. Assume now that $|f(p^\alpha)| \leq C(f(n))$ is assumed to be real valued). We prove the following theorem.

THEOREM 12. *Let $|f(p^\alpha)| \leq c$. Denote by $F(x)$ the distribution function of $f(x)$. We have*

$$F(x) > 1 - \exp(-cx),$$

for every c and sufficiently large x . In other words the density of integers with $f(n) \geq x$ is less than $\exp(-cx)$.

Put $g(n) = \exp(2cf(n))$, $g(n)$ is multiplicative and clearly has a distribution function. Define

$$f_k(n) = \sum_{p|n, p \leq k} f(p), \quad g_k(n) = \exp(2cf_k(n)).$$

For sake of simplicity we assume that $f(p^\alpha) = f(p)$. It is well known that the distribution function $F_k(x)$ of $f_k(n)$ converges to $F(x)$, thus the distribution function $G_k(x)$ of $g_k(x)$ converges to $G(x)$ ($G(x)$ is the distribution function of $g(x)$). Suppose now that Theorem 12 is false, then there exists a constant c and infinitely many x_r with $x_r \rightarrow \infty$ and

$$F(x_r) > 1 - \exp(-cx_r).$$

Therefore for any r there exists a k so large that

$$F_k(x_r) > 1 - \exp(-cx_r).$$

⁷ P. Erdős and A. Wintner, Amer. J. Math. vol. 61 (1939) pp. 713-721.

Thus the density of integers with $g_k(n) > \exp(2cx_r)$ is greater than $\exp(-cx_r)$ and hence

$$\sum_{m \leq n} g_k(m) > (1 - \epsilon) \exp(cx_r) \cdot n$$

for n sufficiently large. Thus for any A there exists k and n_0 , such that for all $n > n_0$

$$(17) \quad \sum_{m \leq n} g_k(m) > An.$$

On the other hand

$$\sum_{m \leq n} g_k(m) = \sum_{m=1}^n \prod_{p|m} g_k(p) = \sum_{m=1}^n \prod_{p|m} (1 + (g_k(p) - 1)).$$

Put $g_k(p) - 1 = h_k(p)$. Clearly

$$\sum_{m=1}^n g_k(m) = \sum_{m=1}^n \prod_{p|m} (1 + h_k(p)) = \sum_d \left[\frac{n}{d} \right] h_k(d)$$

where $h_k(d) = \prod_{p|d} h_k(p)$. Thus

$$\sum_{m=1}^n g_k(m) \leq n \sum_d \frac{h_k(d)}{d} = n \prod_p \left(1 + \frac{h_k(p)}{p} \right).$$

From the fact that $g(n)$ has a distribution function and that $f(p^\alpha)$ is bounded, it easily follows that (we shall give the details in the proof of Theorem 13)

$$\sum_p \frac{h(p)}{p} < \infty, \quad \sum_p \frac{(h(p))^2}{p} < \infty, \quad h(p) = g(p) - 1.$$

Thus finally

$$\sum_{m=1}^n g_k(m) < c_1 n \prod_p \left(1 + \frac{h(p)}{p} \right) < c_2 n,$$

which contradicts (17), and this contradiction establishes the theorem.

It is easy to see that Theorem 12 is best possible. Let $\phi(x)$ tend to infinity arbitrarily slowly; then there exists an additive function $f(n)$ such that its distribution function $F(x)$ satisfies $F(x_i) < 1 - \exp(-\phi(x_i)x_i)$ for an infinite sequence x_i with $x_i \rightarrow \infty$. We omit the proof.

THEOREM 13. *Let $g(n) \geq 0$ be multiplicative. Then the necessary and sufficient condition for the existence of a distribution function is that*

$$(18) \quad \sum_p \frac{(g(p) - 1)'}{p} < \infty, \quad \sum_p \frac{((g(p) - 1)')^2}{p} < \infty,$$

where $(g(p) - 1)' = g(p) - 1$ if $|g(p) - 1| \leq 1$ and 1 otherwise.

The proof follows very easily from (16). Put $\log(g(n)) = f(n)$. $g(n)$ has a distribution function if and only if $f(n)$ has a distribution function. Thus from (16)

$$(19) \quad \sum_p \frac{(\log g(p))'}{p} < \infty, \quad \sum_p \frac{((\log g(p))')^2}{p} < \infty.$$

Now it follows from (19) that if we neglect a sequence of primes q with $\sum 1/q < \infty$ that $|g(p) - 1| < 1/2$. Thus

$$\log g(p) = \log(1 + (g(p) - 1)) = (g(p) - 1) + (1/2)(g(p) - 1)^2 + \dots$$

Also simple computation shows that $(\log g(p))' > (1/4)(g(p) - 1)^2$. Thus from (19)

$$\sum_p \frac{(g(p) - 1)^2}{p} < \infty$$

and

$$\sum_p ((1/2)(g(p) - 1)^2 + (g(p) - 1)^3 + \dots) < \infty.$$

Thus $\sum_p (g(p) - 1)/p < \infty$, which shows that (18) is necessary.

If the two series in (18) converge, then clearly

$$\sum_p \frac{\log g(p)}{p} = \sum_p \left(\frac{(g(p) - 1)}{p} + \frac{(1/2)(g(p) - 1)^2}{p} + \dots \right) < \infty$$

and

$$\sum_p \frac{(\log g(p))^2}{p} < c \sum_p \frac{(g(p) - 1)^2}{p} < \infty,$$

which shows that $f(n)$, and therefore $g(n)$, has a distribution function. Thus (18) is necessary, which completes the proof of Theorem 13.

These results suggest that if $g(n)$ is multiplicative, satisfies (18), $|g(p^\alpha)| < c$, then $g(n)$ has a mean value, that is, $\lim_{x \rightarrow \infty} (1/x) \sum_{n \leq x} f(n)$ exists. I have not yet been able to prove this.