

quences, $g(u) - \overline{D}'(p(u))$ in (2') being replaced by $g(u)$, as it may be under conditions on the $A_n(\sigma)$ discussed above. For by their theorem, given

$$\int^{\infty} p(\sigma) \exp \left[-\frac{1}{2} \int^{\sigma} \frac{du}{g(u)} \right] d\sigma < \infty,$$

there exists a function $F(s)$ holomorphic in Δ , not identically zero, such that $|F(s) - \sum_1^n 0e^{-\lambda_k s}| < e^{-p(\sigma)}$, hence $|F(s) - \sum_1^n 0e^{-\lambda_k s}| < A_n(\sigma)$ if $\{A_n(\sigma)\}$ is any asymptotic sequence with $\text{g.l.b.}_{n \geq 1} A_n(\sigma) = A(\sigma) = e^{-p(\sigma)}$; so that $F(s)$ is represented asymptotically in Δ by the series $\sum d_k e^{-\lambda_k s}$ with $d_k = 0$ ($k \geq 1$) with respect to the asymptotic sequence $\{A_n(\sigma)\}$, without being identically zero.

BIBLIOGRAPHY

1. S. Mandelbrojt, *Sur une inégalité fondamentale*, Ann. École Norm. (3) vol. 43 (1946) pp. 351-378.
2. S. Mandelbrojt and G. R. MacLane, *On functions holomorphic in a strip region, and an extension of Watson's problem*, Trans. Amer. Math. Soc. vol. 61 (1947) pp. 454-467.
3. S. Mandelbrojt, *Analytic continuation and infinitely differentiable functions*, Bull. Amer. Math. Soc. vol. 54 (1948) pp. 239-248.

RICE INSTITUTE

ERRATA

R. D. Carmichael, *On Euler's ϕ function*, vol. 13, pp. 241-243; vol. 54, p. 1192.

Vol. 54, p. 1192, lines 2 and 9. For "Hedburg" read "Hedberg."

Vol. 54, p. 1192, line 10. For " $2^{28}+1$ and $2^{29}+1$ " read " $2^{28}+1$ and $2^{29}+1$."