

ON NORMAL METRICS, AND A THEOREM OF COHN-VOSSEN

BY ROBERT FINN

Communicated by S. S. Chern, July 10, 1964

1. Let \mathfrak{G} denote the domain consisting of the complex z -plane with $n+1$ points $p_1, \dots, p_n, p_0 = \infty$ deleted. Let $\mu(E)$ be a measure over \mathfrak{G} , of finite total variation. Let

$$u(x, y) = \int_{\mathfrak{G}} \log \left| 1 - \frac{z}{\zeta} \right| d\mu_{\zeta} + h(z),$$

where $h(z)$ is harmonic in \mathfrak{G} . The conformal metric $e^{u(z)}|dz|$ will be said to be *normal* in \mathfrak{G} , provided $h(z)$ has the form

$$h(z) = \sum_1^n \beta_j \log |z - p_j| + \text{const.}$$

The metric is said to be *complete* if any path tending to one of the $\{p_j\}$ has infinite length. The *curvatura integra* is $C = -2\pi \int_{\mathfrak{G}} d\mu_{\zeta}$, and the *flux* at p_j is defined by

$$\Phi_j = \lim_{\gamma_j \rightarrow p_j} \frac{1}{2\pi} \oint_{\gamma_j} \frac{\partial u}{\partial n} |dz|,$$

for curves γ_j enclosing p_j .

Let Γ_j, γ_j be concentric circumferences surrounding p_j , and let $\mathcal{Q}_j(\Gamma, \gamma)$ be the area of the enclosed annulus in the given metric. Let $\mathcal{L}_j(\gamma)$ be the length of γ_j .

THEOREM. *If the metric is complete, then $\Phi_0 \geq -1, \Phi_j \geq 1, j > 0$.¹ The quantities*

$$\nu_j = \lim_{\gamma_j \rightarrow p_j} \frac{\mathcal{L}_j(\gamma)}{4\pi \mathcal{Q}_j(\Gamma, \gamma)}$$

exist and are finite for each j , and $\nu_0 = \Phi_0 + 1, \nu_j = \Phi_j - 1, j > 0$. There holds

$$C = 2\pi \left(\chi - \sum_0^n \nu_j \right),$$

where $\chi = 1 - n$ is the Euler characteristic of \mathfrak{G} .

¹ This assertion follows alternatively from the work of Huber [2]. The demonstrations differ.

This result contains, for the case considered, a theorem of Cohn-Vossen [1] (see also Huber [2]), according to which $C \leq 2\pi\chi$. Cases for which equality holds are easily constructed, but one can also give examples for which the $\{\nu_j\}$ are arbitrarily large.

Asymptotic estimates for $\mathcal{L}_j(\gamma)$ and $\mathcal{Q}_j(\Gamma, \gamma)$ are given. The curves $\{\gamma_j\}$ are shown to behave as approximate geodesic circles in the metric, thus permitting *a posteriori* an interpretation of the result in terms of the explicitly given geometry. The asymptotic lengths of the radial segments emanating from the $\{p_j\}$ are also estimated, and these curves are shown to approximate geodesics in the metric.

2. Let S be a finitely connected, open Riemann surface on which a conformal metric $e^{u(z)}|dz|$ is defined, and suppose the *curvatura integra* associated with the metric is finite. Suppose in addition that the region of positive curvature has compact support on S .

THEOREM. *If the metric is complete, then a doubly connected neighborhood of each boundary component can be mapped conformally onto the exterior of the unit disc. Defining Γ_j, γ_j as above, the limit*

$$\nu_j = \lim_{\gamma_j \rightarrow \infty} \frac{\mathcal{L}_j(\gamma)}{4\pi\mathcal{Q}_j(\Gamma, \gamma)}$$

exists for each such mapping, and $\nu_j = \Phi_j + 1$. There holds

$$C = 2\pi \left(\chi - \sum_0^n \nu_j \right).$$

The essential step in the proof consists in reducing the problem to the study of metrics, whose local structure is normal.

The reader will very likely share my discomfort over the hypothesis that the region $K > 0$ has compact support. I have been unable to prove the result under significantly weaker assumptions.

REFERENCES

1. S. Cohn-Vossen, *Kürzeste Wege und Totalkrümmung auf Flächen*, *Compositio Math.* **2** (1935), 69–133.
2. A. Huber, *On subharmonic functions and differential geometry in the large*, *Comment. Math. Helv.* **32** (1957), 13–72.