

FIXED POINT FREE INVOLUTIONS ON HOMOTOPY SPHERES

BY W. BROWDER AND G. R. LIVESAY¹

Communicated by J. Milnor, November 9, 1966

1. Introduction and statements of theorems. Let $T: \Sigma^{n+1} \rightarrow \Sigma^{n+1}$ be a smooth² (C^∞) fixed point free involution on a smooth manifold, Σ^{n+1} , homeomorphic to the $(n+1)$ -sphere, S^{n+1} . We wish to consider the following problem: does there exist an n -sphere, S^n , smoothly imbedded in Σ^{n+1} such that $TS^n = S^n$? If such an S^n exists, we will say that (T, Σ^{n+1}) *desuspends* to $(T|S^n, S^n)$ and that $(T|S^n, S^n)$ *suspends* to (T, Σ^{n+1}) . We claim (proofs are to appear later):

THEOREM 1. *If $n \geq 5$ is odd, then (T, Σ^{n+1}) desuspends to $(T|S^n, S^n)$ for some T -invariant $S^n \subset \Sigma^{n+1}$.*

If n is even, there are obstructions to desuspending (T, Σ^{n+1}) . There is a bilinear form, $B(x, y)$ defined on a certain subgroup of $H_*(M)$, where $\Sigma^{n+1} = A \cup TA$, A and TA are compact submanifolds of Σ^{n+1} with smooth boundary, and $\partial A = \partial TA = A \cap TA = M$. If $n \equiv 2 \pmod{4}$, then B is symmetric, and its signature, $\sigma(T, \Sigma^{n+1})$ is determined by (T, Σ^{n+1}) . If $n \equiv 0 \pmod{4}$, then B is skew-symmetric. Furthermore, if $n = 4k$, there is a map $\psi_0: H_{2k}(M; Z_2) \rightarrow Z_2$ such that $\psi_0(x+y) = \psi_0(x) + \psi_0(y) + B_2(x, y)$, where B_2 , defined on a subgroup of $H_{2k}(M; Z_2)$, corresponds to B , defined on a subgroup of $H_{2k}(M)$. The Arf invariant, $c(T, \Sigma^{n+1})$, [1], [4], corresponding to ψ_0 and B_2 , depends only on (T, Σ^{n+1}) . Regarding these invariants, we have

THEOREM 2. *If $n \equiv 2 \pmod{4}$ and $n > 5$, then (T, Σ^{n+1}) can be desuspended to $(T|S^n, S^n)$ if and only if $\sigma(T, \Sigma^{n+1}) = 0$.*

THEOREM 3. *If $n \equiv 0 \pmod{4}$ and $n > 4$, then (T, Σ^{n+1}) can be desuspended to $(T|S^n, S^n)$ if and only if $c(T, \Sigma^{n+1}) = 0$.*

At present, we have no example of (T, Σ^{n+1}) for which either $c(T, \Sigma^{n+1}) \neq 0$ for $n \equiv 0 \pmod{4}$, or $\sigma(T, \Sigma^{n+1}) \neq 0$ for $n \equiv 2 \pmod{4}$. An interesting example to study in connection with the possibility of

¹ The authors were partially supported by contracts GP 2425 and GP 3685, respectively, with the National Science Foundation.

² The results hold equally in the piecewise linear category with little change in the proofs.

a nonzero Arf invariant is the nonstandard involution of Hirsch and Milnor on S^5 , [3]. However, even if the Arf invariant is zero, our methods do not give a desuspension of this involution to the 4-sphere, because of the usual difficulties of finding a basis for $H_2(M^4)$ represented by imbedded spheres.

Regarding the uniqueness of the desuspension, we have

THEOREM 4. *If $n \geq 4$ is even, and (T, Σ^{n+1}) desuspends to $(T|S_0^n, S_0^n)$ and to $(T|S_1^n, S_1^n)$, then $(T|S_0^n, S_0^n)$ and $(T|S_1^n, S_1^n)$ are equivariantly concordant in $\Sigma^{n+1} \times I$.*

We say that (T_0, S_0^n) and (T_1, S_1^n) are *concordant* if there exists a fixed point free involution $T: S^n \times I \rightarrow S^n \times I$, where $I = [0, 1]$, such that $T(S^n \times 0) = S^n \times 0$, and equivariant diffeomorphisms $i_0: (T_0, S_0^n) \rightarrow (T|S^n \times 0, S^n \times 0)$ and $i_1: (T_1, S_1^n) \rightarrow (T|S^n \times 1, S^n \times 1)$. If $\bar{T}: \Sigma^{n+1} \rightarrow \Sigma^{n+1}$ is a smooth, fixed point free involution, then (T_0, S_0^n) and (T_1, S_1^n) are *concordant in $\Sigma^{n+1} \times I$* if they are concordant, and $(T, S^n \times I)$ is equivariantly imbedded in $(\bar{T} \times 1, \Sigma^{n+1} \times I)$ with $S^n \times 0 \subset \Sigma^{n+1} \times 0$, $S^n \times 1 \subset \Sigma^{n+1} \times 1$. If $n > 4$ is odd, the signature and Arf invariant, which appeared as obstructions to desuspending (T, Σ^k) , now appear as obstructions to obtaining a concordance in $(T \times 1, \Sigma^{n+1} \times I)$ between two given desuspensions, $(T|S_0^n, S_0^n)$ and $(T|S_1^n, S_1^n)$. If S_0^{4k-1} and S_1^{4k-1} are two invariant spheres in (T, Σ^{4k}) , then $\sigma(T, \Sigma^{4k}, S_0^{4k-1}, S_1^{4k-1})$, the signature of a certain bilinear form, is defined. We then have

THEOREM 5. *S_0^{4k-1} and S_1^{4k-1} are concordant in $(T \times 1, \Sigma^{4k} \times I)$ if and only if $\sigma(T, \Sigma^{4k}, S_0^{4k-1}, S_1^{4k-1}) = 0$. In particular, if $\sigma = 0$, then $(T|S_0^{4k-1}, S_0^{4k-1})$ and $(T|S_1^{4k-1}, S_1^{4k-1})$ are equivariantly diffeomorphic.*

Now suppose S_0^{4k+1} and S_1^{4k+1} are invariant spheres in (T, Σ^{4k+2}) . Then $c(T, \Sigma^{4k+2}, S_0^{4k+1}, S_1^{4k+1})$, an Arf invariant, is defined.

THEOREM 6. *S_0^{4k+1} and S_1^{4k+1} are concordant in $(T \times 1, \Sigma^{4k+2} \times I)$ if and only if $c(T, \Sigma^{4k+2}, S_0^{4k+1}, S_1^{4k+1}) = 0$.*

COROLLARY. *If $n \equiv 1 \pmod{4}$, there are at most two invariant n spheres in Σ^{n+1} , up to equivariant diffeomorphism.*

It is planned to present detailed proofs later. We will, however, indicate briefly some of the ideas involved.

2. Characteristic submanifolds. Let $T: \Sigma^{n+1} \rightarrow \Sigma^{n+1}$ be a fixed point free smooth involution. A *characteristic submanifold* $M^n \subset \Sigma^{n+1}$ is an n -manifold smoothly imbedded in Σ^{n+1} such that $\Sigma^{n+1} = A \cup TA$ with $A \cap TA = M^n$. We have a commutative square

$$\begin{array}{ccc} \Sigma^{n+1} & \longrightarrow & S^N \\ \downarrow \pi & & \downarrow \\ \Sigma^{n+1}/T & \xrightarrow{f} & P^N \end{array}$$

where N is large, P^N is a real projective N -space, and f classifies the principal Z_2 -bundle $\Sigma^{n+1} \xrightarrow{\pi} \Sigma^{n+1}/T$. By making f transverse-regular [5] on P^{N-1} , $\pi^{-1}f^{-1}P^{N-1}$ will be a characteristic submanifold. It is easy to see that all characteristic submanifolds arise in this way. Any two characteristic submanifolds are equivariantly cobordant in $(T \times 1, \Sigma^{n+1} \times I)$. (The definition is analogous to that of concordance in $\Sigma^{n+1} \times I$.) It is this fact that makes the signature and Arf invariant independent of the choice of characteristic submanifold.

3. The signature and Arf invariant. Let M be a characteristic submanifold in Σ^{n+1} . Then $\Sigma^{n+1} = A \cup TA$ with $A \cap TA = M$. We have the Mayer-Vietoris sequence

$$\dots \rightarrow H_{p+1}(\Sigma^{n+1}) \rightarrow H_p(M) \xrightarrow{(i_A, i_{TA})} H_p(A) \oplus H_p(TA) \rightarrow H_p(\Sigma^{n+1}) \rightarrow \dots$$

If $n = 2k$, $k > 0$, and $p = k$, this becomes

$$0 \rightarrow H_k(M^{2k}) \xrightarrow{(i_A, i_{TA})} H_k(A) \oplus H_k(TA) \rightarrow 0$$

and so $H_k(M^{2k}) = \ker i_A \oplus \ker i_{TA}$, and $T_* \ker i_A = \ker i_{TA}$. Since $M^{2k} \subset \Sigma^{2k+1}$, M is orientable, and a bilinear form $B(x, y) = x \cdot T_* y$ is defined, for x and y in $\ker i_A$. Since T preserves orientation in Σ^{2k+1} , it reverses orientation in M^{2k} , and the bilinear form B is symmetric (skew-symmetric) when the intersection form $x \cdot y$ is skew-symmetric (symmetric). Therefore, given (T, Σ^{n+1}) and a characteristic submanifold M^n , if $n \equiv 2 \pmod{4}$, the signature of the form $B(x, y)$ is determined, and turns out to be independent of the choice of characteristic submanifold. The reason for considering the signature of B is the following. If $x \in \ker i_A \subset H_k(M^{2k})$, and M^{2k} is $(k-1)$ -connected, (which we achieve by exchanging handles between A and TA) then x is represented by an imbedded $S^k \subset M^{2k}$, which bounds a cell $D^{k+1} \subset A$. (This statement may be false for $k = 3$, [2], but a different argument applies in this case.) Supposing M^{2k} is totally geodesic near D^{k+1} , we take a tubular neighborhood N of D^{k+1} , replace A by $A - N$, and replace TA by $TA \cup \bar{N}$. This will reduce the rank of $H_k(M)$. However, $(A - N) \cap (TA \cup \bar{N}) = M'$ is no longer T -invariant. We may obtain an invariant M' if we replace A by $(A - N) \cup T\bar{N}$, and replace TA by $(TA \cup \bar{N}) - TN$. However, to do this we need $S^k \cap TS^k = \emptyset$. It is to accomplish this that we need $\sigma = 0$

when k is odd and $c=0$ when k is even. The distinction between the two cases arises since if S^k and TS^k intersect transversally in M^{2k} at a point p with intersection number 1, then they intersect at Tp with intersection number $(-1)^{k+1}$.

The cohomology operation, $\psi(x)$, used to define the Arf invariant, merely serves to count, mod 2, the number of pairs (q, Tq) of points in $S^p \cap TS^p$, where S^p represents the Poincaré dual of x , and the intersection is transverse.

REFERENCES

1. C. Arf, *Untersuchungen über quadratische Formen in Körpern der Charakteristik 2*, J. Reine Angew. Math. **183** (1941), 148–167.
2. A. Haefliger, *Knotted $(4k-1)$ -spheres in $6k$ -space*, Ann. of Math. **75** (1962), 452–466.
3. M. W. Hirsch and J. W. Milnor, *Some curious involutions of spheres*, Bull. Amer. Math. Soc. **70** (1964), 372–377.
4. M. A. Kervaire and J. W. Milnor, *Groups of homotopy spheres. I*, Ann. of Math. **77** (1963), 504–537.
5. R. Thom, *Quelques propriétés global des variétés différentiables*, Comment. Math. Helv. **28** (1954), 17–86.

INSTITUTE FOR ADVANCED STUDY,
PRINCETON UNIVERSITY AND
CORNELL UNIVERSITY