

INDUCTIVELY DEFINED SETS OF REALS

BY DOUGLAS CENZER

Communicated by Alistair H. Lachlan, October 8, 1973

1. Introduction. The concept of inductive definability has become of great interest to recursion theorists in recent years. Recursion over natural numbers, ordinals, and higher type objects may itself be defined by an inductive operator—see for example [7] and [9]. Many results have been obtained characterizing the closures of inductive operators over the natural numbers, and relating lengths of inductive definitions to various interesting ordinals; see [3] for a brief summary.

The purpose of this note is to present results on the closure ordinals and sets of inductive operators over the continuum. Details will appear later in [2], [4], and [5].

2. Basic definitions and notation. An inductive operator Γ over a set X is a map from $P(X)$ to $P(X)$ such that for all A , $A \subseteq \Gamma(A)$. Γ determines a transfinite sequence $\{\Gamma^\sigma : \sigma \in \text{ORD}\}$, where $\Gamma^\sigma = U\{\Gamma^\tau : \tau < \sigma\}$ for $\sigma = 0$ or σ a limit and $\Gamma^{\sigma+1} = \Gamma(\Gamma^\sigma)$. Γ is monotone if, for all A, B in $P(X)$, $A \subseteq B$ implies $\Gamma(A) \subseteq \Gamma(B)$. Γ is positive if its application to a set A involves only the positive part of χ_A (the characteristic function of A).

The closure ordinal $|\Gamma|$ of Γ is the least ordinal σ such that $\Gamma^{\sigma+1} = \Gamma^\sigma$; clearly $|\Gamma|$ always has cardinality less than or equal to $\text{card}(X)$. The closure $\bar{\Gamma}$ of Γ is $\Gamma^{|\Gamma|}$, the set inductively defined by Γ .

For a class C of inductive operators, the closure ordinal $|C| = \sup\{|\Gamma| : \Gamma \in C\}$ and the closure algebra $\bar{C} = \{A : A \text{ is 1-1 reducible to } \bar{\Gamma} \text{ for some } \Gamma \text{ in } C\}$. We write $C\text{-mon}$ for the class of monotone operators in C .

In studying inductive operators over the continuum, we follow the usual convention that a real number is a function from the set ω of natural numbers to itself; thus the real line is ${}^\omega\omega$.

3. Main results. The central result of our research is the following theorem.

- THEOREM 1.** (a) $|\Pi_1^0\text{-mon}| = |\Pi_1^1\text{-mon}| = |\Sigma_1^1\text{-mon}| = \aleph_1$;
 (b) $(\Pi_1^0\text{-mon})^- = (\Pi_1^1\text{-mon})^- = \Pi_1^1$;
 (c) $(\Sigma_1^1\text{-mon})^- = \Sigma_2^1$.

AMS (MOS) subject classifications (1970). Primary 02F29; Secondary 02F27, 02F35, 02K30.

Key words and phrases. Inductive definition, continuum, ordinal, closure, inductively parametrized.

Copyright © American Mathematical Society 1974

One direction of these equalities is derived from the existence of a Π_1^0 monotone operator Γ with $|\Gamma| = \aleph_1$ and Γ equal to the set of codes for well-orderings.

The other direction depends heavily on first proving that both Π_1^1 and Σ_1^1 monotone operators can be put in positive form. See [1] for some related results.

Concerning nonmonotone inductive definitions, we obtain the following result by considering the essential countability of any computation on real numbers.

THEOREM 2. (a) $|\Sigma_1^1| = \aleph_1$;
 (b) $(\Delta_1^1)^- \subseteq \Sigma_2^1$.

It is interesting to compare these two theorems with the corresponding results for inductive definitions over the natural numbers, where

$$(1) \quad |\Pi_1^0\text{-mon}| = |\Pi_1^1\text{-mon}| = \omega_1 < |\Delta_1^1| < |\Sigma_1^1\text{-mon}|$$

and

$$(2) \quad (\Pi_1^0\text{-mon})^- = (\Pi_1^1\text{-mon})^- = \Pi_1^1 \subseteq (\Sigma_1^1\text{-mon})^- \subseteq \Sigma_2^1.$$

The simplest type of inductive definition over the reals would be one in which the real variable appeared only as a parameter. For example, if R is a Π_1^0 relation such that for all α , Γ_α , defined by $n \in \Gamma_\alpha(A) \leftrightarrow R(n, \alpha, A)$, is an inductive operator over ω , then there is an obvious Π_1^0 operator Γ such that for all n and α , $n \in (\Gamma_\alpha)^-$ iff $\langle n, \alpha \rangle \in \Gamma$. A similar fact holds if Π_1^0 is replaced by any other definability class.

We see by Theorem 1(b) and remark (2) that for Π_1^0 this process can be reversed: If Γ is a Π_1^0 monotone operator over ${}^\omega\omega$, then there is a Π_1^0 relation R such that for all α , Γ_α , defined by $n \in \Gamma_\alpha(A) \leftrightarrow R(n, \alpha, A)$, is a monotone operator over ω and $\alpha \in \Gamma$ iff $0 \in \Gamma_\alpha$. We say that the Γ_α parametrize Γ .

Comparing Theorem 1(c) with remark (2), we see that this cannot be done for Σ_1^1 operators over ${}^\omega\omega$. The surprising result here will be the extent of the definability class needed to parametrize Δ_1^0 nonmonotone operators.

In general, we say that $A \subseteq {}^\omega\omega$ can be *inductively parametrized* in C if there is a $\{\Gamma_\alpha : \alpha \in {}^\omega\omega\}$ in C and a recursive F such that for all α in ${}^\omega\omega$, $\alpha \in A$ iff $F(\alpha) \in \Gamma_\alpha$. Let $\text{PC1}(C)$ be the class of subsets of ${}^\omega\omega$ which can be inductively parametrized in C .

By the above remarks, $\text{PC1}(\Pi_1^0) = \Pi_1^1 = (\Pi_1^0\text{-mon})^-$. The power of recursive inductive operators over the continuum can be seen in the following result.

THEOREM 3. (a) $|\Delta_1^0| = \aleph_1$;
 (b) $\text{PC1}(\Pi_1^1) \cup \text{PC1}(\Sigma_1^1) \subseteq (\Delta_1^0)^-$.

Inductive definitions of length greater than \aleph_1 are closely related to the functionals 3E and ${}^3E^\#$ —see [8] for definitions. Let κ_1 be the ordinal of 3E , κ_2 that of ${}^3E^\#$; κ_1 is less than κ_2 . The following two results are implicit in [6].

PROPOSITION 4. $|\Pi_1^1| = \kappa_1$; $(\Pi_1^1)^- = \text{semirecursive in } {}^3E \text{ (hyperanalytic)}$.

PROPOSITION 5. For all $n > 1$, $|\Delta_n^1\text{-mon}| = |\Sigma_2^1| = \kappa_2$ and $(\Delta_n^1\text{-mon})^- = (\Sigma_2^1)^- = \text{semirecursive in } {}^3E^\# \text{ (hyperprojective)}$.

We can add the following.

THEOREM 6. $|\Pi_2^1| > \kappa_2$.

Hyperprojective inductive definitions can be analyzed in the manner of [3] and [10]. For example, we obtain the following analogue of a theorem of Richter from [10].

THEOREM 7. $|\Delta_2^1, \Delta_2^1|$ is the least ordinal which is recursively Mahlo in ${}^3E^\#$ and parameters from ${}^\omega\omega$.

All of the results in this paper are easily relativized with respect to a real parameter and have obvious boldface analogues.

REFERENCES

1. A. Blass and D. Cenzer, *Cores of Π_1^1 sets of reals* (to appear).
2. D. Cenzer, *Monotone inductive definitions over the continuum*, Notices Amer. Math. Soc. **21** (1974), A-19.
3. ———, *Ordinal recursion and inductive definitions*, J. Fenstad and P. Hinman (Ed.), Generalized Recursion Theory, Oslo, 1972, North-Holland, Amsterdam (to appear).
4. ———, *Ordinal recursion over the continuum* (to appear).
5. ———, *Parametric inductive definitions and recursive operators over the continuum* (to appear).
6. T. Grilliot, *Hierarchies based on objects of finite type*, J. Symbolic Logic **34** (1969), 177–182. MR **44** #75.
7. P. Hinman, *Recursion-theoretic hierarchies* (to appear).
8. P. Hinman and Y. Moschovakis, *Computability over the continuum*, Logic Colloquium '69, North-Holland, Amsterdam, 1971, pp. 77–105.
9. S. C. Kleene, *Recursive functionals and quantifiers of finite types. II*, Trans. Amer. Math. Soc. **108** (1963), 106–142. MR **27** #3521.
10. W. Richter, *Recursively Mahlo ordinals and inductive definitions*, Logic Colloquium '69 (Proc. Summer School and Colloq., Manchester, 1969), North-Holland, Amsterdam, 1971, pp. 273–288. MR **43** #7331.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32601