

JOINT SPECTRUM IN THE CALKIN ALGEBRA

BY A. T. DASH¹

Communicated by Robert Bartle, June 20, 1975

For a nice discussion pertaining to the essential spectrum of a single operator (bounded linear transformation) in a complex separable infinite dimensional Hilbert space H , the reader is referred to Fillmore, Stampfli and Williams [4]. The purpose of this note is to announce analogous results concerning the joint essential spectra of n -tuples of operators in H .

Joint essential spectrum. In the sequel $L(H)$ denotes the algebra of all operators on H and K denotes the ideal of compact operators on H . Let ν be the canonical homomorphism from $L(H)$ onto the Calkin algebra $L(H)/K = \mathcal{C}$. If $A = (A_1, \dots, A_n)$ is an n -tuple of operators on H , then we write $\nu(A_j) = a_j$, the coset containing A_j for each j , $1 \leq j \leq n$, and $a = (a_1, \dots, a_n)$.

The *joint essential spectrum* of an n -tuple of operators A denoted by $\sigma_e(A)$ is defined to be the *joint spectrum* $\sigma(a)$ of a .

Here $\sigma(a) = \sigma^l(a) \cup \sigma^r(a)$, where the *left (right) joint spectrum* $\sigma^l(a)$ ($\sigma^r(a)$) is defined as the set of all $z = (z_1, \dots, z_n)$ in \mathbb{C}^n (n -fold Cartesian product of the set of all complex numbers \mathbb{C}) such that $\{a_j - z_j\}_{1 \leq j \leq n}$ generates a proper left (right) ideal in the Calkin algebra \mathcal{C} . For this notion of joint spectrum, the reader may consult [1] and [5]. We call the set $\sigma^l(a)$ ($\sigma^r(a)$) as the *left (right) joint essential spectrum* and denote it by $\sigma_e^l(A)$ ($\sigma_e^r(A)$). Clearly, $\sigma_e^l(A) \subseteq \sigma^l(A)$, $\sigma_e^r(A) \subseteq \sigma^r(A)$; and hence $\sigma_e(A) \subseteq \sigma(A)$. Further, if $A = (A_1, \dots, A_n)$ is an n -tuple of *essentially commuting* (commuting modulo the compacts) operators, then $\sigma_e(A)$ is a nonempty compact subset of \mathbb{C}^n .

The following theorem describes the relationship between the joint spectrum and the joint essential spectrum of an n -tuple of operators.

THEOREM 1. *Let $A = (A_1, \dots, A_n)$ be an n -tuple of operators on H . Then $\sigma(A) = \sigma_e(A) \cup \sigma_p(A) \cup \sigma_p(A^*)^*$, where $A^* = (A_1^*, \dots, A_n^*)$ and star on the right represents complex conjugates.*

A point $z = (z_1, \dots, z_n)$ of \mathbb{C}^n is in $\sigma_p(A)$ (the *joint eigenvalue* of A) if

AMS (MOS) subject classifications (1970). Primary 47A05, 47A10, 47B05, 47B20, 47B30, 46H10.

Key words and phrases. Joint spectrum, joint essential spectrum, joint eigenvalues, Calkin algebra, hyponormal elements, compact operators, Fredholm operators.

¹ This research was supported by NRC Grant A07545.

and only if there exists a nonzero vector f in H such that $(A_j - z_j)f = 0$ for each $j, 1 \leq j \leq n$ (consult [3]).

COROLLARY 1. *Let $A = (A_1, \dots, A_n)$ be as given above. Then:*

(a) $\sigma^l(A)$ consists of $\sigma_e^l(A)$ together with the joint eigenvalues of finite multiplicity.

(b) $\sigma^r(A)$ consists of $\sigma_e^r(A)$ together with the set of all $z = (z_1, \dots, z_n)$ in C^n such that z^* is a joint eigenvalue of finite multiplicity of A^* .

The next theorem characterizes the joint essential spectrum of special operators.

THEOREM 2. *Let $A = (A_1, \dots, A_n)$ be an n -tuple of essentially hyponormal operators ($a_j a_j^* \leq a_j^* a_j, 1 \leq j \leq n$). Then $\sigma_e(A) = \sigma_e^r(A)$.*

COROLLARY 2 [2, LEMMA 2.1]. *Let $A = (A_1, \dots, A_n)$ be an n -tuple of essentially normal ($A_j^* A_j - A_j A_j^*$ is compact for each $j, 1 \leq j \leq n$) operators. Then $\sigma_e(A) = \sigma_e^l(A)$.*

Joint eigenvalues in the Calkin algebra. It is known that if $b \in C$ and $z \in \sigma(b)$, then there is a projection $p \neq 0$ such that $bp = zp$ or $pb = zp$ [4]. The following theorem is an extension of this result to n -tuples of elements in C .

THEOREM 3. *Let $a = (a_1, \dots, a_n)$ be an n -tuple of elements in the Calkin algebra C and $z = (z_1, \dots, z_n) \in \sigma(a)$. Then there is a projection $p \neq 0$ such that either $a_j p = z_j p$ for all $j, 1 \leq j \leq n$, or $p a_j = z_j p$ for all $j, 1 \leq j \leq n$.*

COROLLARY 3. *Let $A = (A_1, \dots, A_n)$ be an n -tuple of essentially commuting operators. Then there are orthogonal projections P and Q of infinite rank and nullity and a point $z = (z_1, \dots, z_n)$ of C^n such that $(A_j - z_j)P$ is compact for all $j, 1 \leq j \leq n$, and $Q(A_j - z_j)$ is compact for all $j, 1 \leq j \leq n$.*

COROLLARY 4. *Let $A = (A_1, \dots, A_n)$ be an n -tuple of essentially commuting operators. Then the operators A_1, \dots, A_n have a common invariant subspace "modulo the compacts".*

THEOREM 4. *Let $a = (a_1, \dots, a_n)$ be an n -tuple of hyponormal elements in the Calkin algebra C . Then:*

(a) $z = (z_1, \dots, z_n) \in \sigma(a)$ if and only if there is a projection $p \neq 0$ such that $a_j^* p = z_j^* p$ for all $j, 1 \leq j \leq n$.

(b) If p is a projection such that $a_j p = z_j p, 1 \leq j \leq n$, then $a_j^* p = z_j^* p, 1 \leq j \leq n$.

The essential key to most of the results above is the following:

THEOREM 5. *The following statements are equivalent:*

(1) $0 = (0, 0, \dots, 0) \in \sigma_e^l(A_1, \dots, A_n)$.

(2) $0 \in \sigma_e(\sum_{j=1}^n A_j^* A_j)$.

(3) *There exists an orthogonal sequence $\{e_k\}$ such that $\|A_j e_k\| \rightarrow 0$ as $k \rightarrow \infty$, for each j , $1 \leq j \leq n$.*

(4) *There exists an infinite dimensional projection P such that $A_j P$ is compact for each j , $1 \leq j \leq n$.*

(5) $\sum_{j=1}^n A_j^* A_j$ *is not Fredholm.*

(6) $0 \in \sigma^l(a_1, \dots, a_n)$.

(7) $0 \in \sigma(\sum_{j=1}^n a_j^* a_j)$.

REMARK. Most of the results above can be extended to sequences $\{A_n\}$ of operators with very little modifications in the proofs. However, for brevity, we have chosen to discuss them for n -tuples of operators in H .

REFERENCES

1. F. F. Bonsall and J. Duncan, *Numerical ranges of operators on normed spaces and of elements of normed algebras*, London Math. Soc. Lecture Note Series, 2, Cambridge Univ. Press, London and New York, 1971. MR 44 #5779.
2. L. G. Brown, R. G. Douglas and P. A. Fillmore, *Unitary equivalence modulo the compact operators and extensions of C^* -algebras*, Proc. Conf. on Operator Theory, Lecture Notes in Math., vol. 345, Springer-Verlag, New York, 1973, pp. 58–128.
3. A. T. Dash, *Joint spectra*, Studia Math. 45 (1973), 225–237. MR 49 #1156.
4. P. A. Fillmore, J. G. Stampfli and J. P. Williams, *On the essential numerical range, the essential spectrum, and a problem of Halmos*, Acta. Sci. Math. (Szeged) 33 (1972), 179–192. MR 48 #896.
5. R. Harte, *The spectral mapping theorem in several variables*, Bull. Amer. Math. Soc. 78 (1972), 871–875. MR 45 #7471.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF GUELPH,
GUELPH, ONTARIO, CANADA