

ON THE COHOMOLOGY OF H -SPACES OF EXCEPTIONAL TYPE

BY JAMES P. LIN

Communicated by S. Eilenberg, November 3, 1975

Since the 1950's, Araki [1], Borel [3], Bott [4], Cartan, and others investigated homological and homotopical properties of Lie groups. Using the differentiable structure of a Lie group, Araki [1] and Borel [3] calculated the mod p cohomology rings of the exceptional groups. In this note, we indicate that many of their results can be generalized to finite H -spaces. In particular, the proofs of these new results will be completely independent of the existence of an infinitesimal Lie structure.

Let G be a simply connected Lie group. Borel [3] used the classification theorem for simple Lie groups to prove that $H_*(G; Z)$ has no p -torsion for $p \geq 7$. The following result follows from a homological argument:

THEOREM 1. *Let X be a simply connected finite H -space. Suppose $H^*(X; Q)$ is isomorphic as algebras to the rational cohomology of an exceptional Lie group. Then $H_*(X; Z)$ has no p -torsion for $p \geq 7$ and has 3 or 5 torsion of order at most 3 or 5.*

Theorem 1 is not true in general for finite simply connected H -spaces. John Harper (unpublished) recently discovered finite simply connected H -spaces whose integral homology has p -torsion for any odd prime p .

Let G be a simply connected Lie group. One of the key results used to compute the homology of Lie groups is Bott's result that the homology of the loops on G , $H_*(\Omega G; Z)$, has no torsion. Bott uses Morse Theory [4] to prove this result. We prove the following analogous result for H -spaces:

THEOREM 2 *Let X be a simply connected finite H -space with $H^*(X; Q)$ isomorphic as algebras to the rational cohomology of an exceptional Lie group. Then $H_*(\Omega X; Z)$ has no odd torsion.*

More recently, Hodgkin [5] and Araki [2] proved that for G a simply connected Lie group, $K^*(G; Z)$ is torsion free. Their proofs depend heavily on the differential structure of a Lie group. Hodgkin uses the classification theorem and results of Borel and Araki on the cohomology of Lie groups. Araki uses the existence of a maximal torus.

The following result can be proven using homological methods:

AMS (MOS) subject classifications (1970). Primary 57F10, 57F25; Secondary 57F05, 55G20, 55D45.

Copyright © 1976, American Mathematical Society

THEOREM 3. *Let X be a simply connected finite H -space with $H^*(X; \mathbb{Q})$ isomorphic as algebras to the rational cohomology of an exceptional Lie group. Then $K^*(X; \mathbb{Z})$ has no odd torsion.*

Details and proofs will appear elsewhere.

REFERENCES

1. S. Araki, *Differential Hopf algebras and the cohomology mod 3 of the compact exceptional groups E_7 and E_8* , Ann. of Math. (2) **73** (1961), 404–436. MR **23** #A1372.
2. ———, *Hopf structures attached to K -theory: Hodgkin's theorem*, Ann. of Math. (2) **85** (1967), 508–525. MR **35** #3653.
3. A. Borel, *Topology of Lie groups and characteristic classes*, Bull. Amer. Math. Soc. **61** (1955), 397–432. MR **17**, 282.
4. R. Bott, *On torsion in Lie groups*, Proc. Nat. Acad. Sci. U.S.A. **40** (1954), 586–588. MR **16**, 12.
5. L. Hodgkin, *On the K -theory of Lie groups*, Topology **6** (1967), 1–36. MR **35** #4950.
6. J. P. Lin, *On the H -spaces with finitely generated cohomology algebras*, Thesis, Princeton University, 1974.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SAN DIEGO, LA JOLLA, CALIFORNIA 92037