

CLASSIFICATION OF SIMPLICIAL TRIANGULATIONS OF TOPOLOGICAL MANIFOLDS

BY DAVID E. GALEWSKI¹ AND RONALD J. STERN²

Communicated by S. Eilenberg, April 8, 1976

In this note we announce theorems which classify simplicial (not necessarily combinatorial) triangulations of a given topological n -manifold M , $n \geq 7$ (≥ 6 if $\partial M = \emptyset$), in terms of homotopy classes of lifts of the classifying map $\tau: M \rightarrow BTOP$ for the stable topological tangent bundle of M to a classifying space $BTRI_n$ which we introduce below. The (homotopic) fiber of the natural map $j: BTRI_n \rightarrow BTOP$ is described in terms of certain groups of PL homology 3-spheres. We also give necessary and sufficient conditions for a closed topological n -manifold M , $n \geq 6$, to possess a simplicial triangulation.

The proofs of these results incorporate recent geometric results of F. Ancel and J. Cannon [1], J. Cannon [2], R. D. Edwards [4], and D. Galewski and R. Stern [5].

In [8], R. Kirby and L. Siebenmann show that in each dimension greater than four there exist closed topological manifolds which admit no piecewise linear manifold structure and hence cannot be triangulated as a combinatorial manifold. Also, R. D. Edwards [3] has recently shown that the double suspension of the Mazer homology 3-sphere is homeomorphic to S^5 , thus showing that a simplicial triangulation of a topological manifold *need not* be combinatorial. But it is still unknown whether or not every topological manifold can be triangulated as a simplicial complex.

Our classification theorems for simplicial triangulations on a given topological manifold take the following forms:

Let $BTOP$ denote the classifying space for stable topological block bundles.

THEOREM 1. *There is a space $BTRI_n$ and a natural map $BTRI_n \rightarrow BTOP$ such that if M is a topological n -manifold, $n \geq 7$ (≥ 6 if $\partial M = \emptyset$) and $\tau: M \rightarrow BTOP$ classifies the stable topological tangent bundle of M , then there is a one-to-one correspondence between the set of concordance classes of simplicial triangulations of M and the set of vertical homotopy classes of lifts of τ to $BTRI_n$.*

AMS (MOS) subject classifications (1970). Primary 57C15.

¹This research was supported in part by National Science Foundation grant GP29585-A4

²This research was supported in part by a Faculty Research Grant at the University of Utah and by National Science Foundation grant MCS76-06393.

The obvious relative versions of Theorem 1 also hold true.

THEOREM 2. *The fiber TOP/TRI_n of $BTRI_n \rightarrow BTOP$ has only two non-zero homotopy groups, namely π_3 and π_4 , and the following sequence is exact.*

$$0 \rightarrow \pi_4 \rightarrow \ker(\alpha: \theta_3^H \rightarrow Z_2) \rightarrow \theta_3^{TRI_n} \rightarrow \pi_3 \rightarrow 0.$$

Here θ_3^H denotes the group of *PL* homology 3-spheres, modulo those which bound acyclic *PL* 4-manifolds, under the operation of connected sum; $\alpha: \theta_3^H \rightarrow Z_2$ is the Kervaire-Milnor-Rochlin map $\alpha(H^3) = I(H^3)/8 \pmod 2$, where $I(H^3)$ is the index of a parallelizable *PL* 4-manifold that H^3 bounds; and $\theta_3^{TRI_n}$ is the group of *PL* homology 3-spheres modulo those which bound acyclic homology 4-manifolds W with $W \times R^{n-4}$ a topological manifold, under the operation of connected sum. Note that if $\Sigma^{n-3} H^3$ is homeomorphic to S^n , then H^3 represents the zero element of $\theta_3^{TRI_n}$.

THEOREM 3. (i) $\pi_3(TOP/TRI_n) \subseteq Z_2$,

(ii) $\pi_3(TOP/TRI_n) = 0$ if and only if there exists a *PL* homology 3-sphere H^3 with $\alpha(H^3) = 1$ and the $(n - 3)$ -suspension of H^3 , $\Sigma^{n-3} H^3$, is homeomorphic to S^n .

(iii) $\pi_4(TOP/TRI_n) = 0$ if and only if every *PL* homology 3-sphere H^3 with $\alpha(H^3) = 0$ and which bounds an acyclic homology 4-manifold W with $W \times R^{n-4}$ a topological manifold, bounds an acyclic *PL* 4-manifold.

THEOREM 4. *There exists a *PL* homology 3-sphere H^3 such that*

- (i) $\alpha(H^3) = 1$,
- (ii) $H^3 \# H^3$ bounds an acyclic *PL* 4-manifold, and
- (iii) $\Sigma^{n-3} H^3$ is homeomorphic to S^n .

If and only if every closed topological n -manifold, $n \geq 6$, can be triangulated as a simplicial complex.

REMARK. For $M = 5$ and M^n oriented, Siebenmann [10] has shown under conditions (i) and (iii) that M is simplicially triangulable. M. Scharlemann has pointed out that if M^5 is unoriented, then (i), (iii) and the fact that $H^3 \# H^3$ bounds a contractible *PL* 4-manifold implies the result. For $6 \leq n \leq 8$, Theorem 4 was proven by M. Scharlemann [9], where in place of (ii) he has the orientability condition that the integral Bockstein of the Kirby-Siebenmann obstruction to putting a *PL* structure on M is zero. T. Matumoto has claimed a version of Theorem 4 under the stronger hypothesis that (iii) be replaced by the condition that $\Sigma^{n-4} H^3$ is homeomorphic to S^{n-1} .

We also investigate the question of whether a given topological n -manifold, $n \geq 6$, can be triangulated as a simplicial homotopy manifold. For example;

PROPOSITION 5. *Suppose that every *PL* homotopy 3-sphere bounds a contractible *PL* 4-manifold. Then there is a one-to-one correspondence between the set of concordance classes of simplicial homotopy manifold triangulations of*

a topological n -manifold M , $n \geq 6$, and concordance classes of PL manifold structures on M .

PROPOSITION 6. *Suppose there exists a bad counterexample to the 3 dimensional Poincaré conjecture; namely suppose there exists a PL homotopy 3-sphere H^3 , with*

(i) $\alpha(H^3) = 1$, and

(ii) $H^3 \# H^3$ bounds a contractible PL 4-manifold.

Then every topological n -manifold, $n \geq 6$, can be triangulated as a simplicial homotopy manifold.

Details of these and related results will appear in [6] and [7].

REFERENCES

1. F. Ancel and J. W. Cannon, *Any embedding of S^{n-1} in S^n ($n \geq 5$) can be approximated by locally flat embeddings*, Notices Amer. Math. Soc. **23** (1976), p. A-308. Abstract #732-G2.
2. J. Cannon, *Taming codimension one generalized manifolds* (preprint).
3. R. D. Edwards, *The double suspension of a certain homology 3-sphere is S^5* , Notices Amer. Math. Soc. **22** (1975), p. A-334. Abstract #75T-G33.
4. ———, *The double suspension of PL homology n -spheres*, Proc. Topology Conf. (Georgia, 1975).
5. D. Galewski and R. Stern. *The relationship between homology and topological manifolds via homology transversality* (preprint).
6. ———, *Classification of simplicial triangulations of topological manifolds* (preprint).
7. ———, *Surgery on compact simplicially triangulated topological manifolds* (preprint).
8. R. C. Kirby and L. C. Siebenmann, *On the triangulation of manifolds and the Hauptvermutung*, Bull. Amer. Math. Soc. **75** (1969), 742-749. MR **39** #3500.
9. M. Scharlemann, *Simplicial triangulations of non-combinatorial manifolds of dimension less than nine*, Inst. for Advanced Study, Princeton, N. J. (preprint).
10. L. C. Siebenmann, *Are nontriangulable manifolds triangulable?* Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga. 1969), (J. Cantrell and C. H. Edwards, Editors), Markham, Chicago, 1970, pp. 77-84. MR **42** #6837.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS,
GEORGIA 30601

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY,
UTAH 84112