

A CHARACTER FORMULA FOR THE DISCRETE SERIES OF A SEMISIMPLE LIE GROUP

BY JORGE VARGAS

ABSTRACT. For a semisimple Lie group G , we provide an explicit formula for the discrete series characters θ_λ restricted to the identity component of a split Cartan subgroup, whenever the parameter lies in a so-called Borel-de Siebenthal chamber and G has both a compact Cartan subgroup and a split Cartan subgroup.

Let G be a connected semisimple Lie group with finite center. The discrete series of G is, by definition, the set of equivalence classes of irreducible unitary representations π , such that π occurs discretely in the left (or right) regular representation of G . According to Harish-Chandra [3], G has a nonempty discrete series if and only if G contains a compact Cartan subgroup. Thus we fix a compact Cartan subgroup $B \subset G$, and a maximal compact subgroup $K \subset G$ which contains B . Let $\mathfrak{g}, \mathfrak{k}, \mathfrak{b}$ be the Lie algebras of G, K, B , and $\mathfrak{g}^{\mathbb{C}}, \mathfrak{k}^{\mathbb{C}}, \mathfrak{b}^{\mathbb{C}}$ their complexifications. Let $\Phi = \Phi(\mathfrak{g}^{\mathbb{C}}, \mathfrak{b}^{\mathbb{C}})$ be the root system of $(\mathfrak{g}^{\mathbb{C}}, \mathfrak{b}^{\mathbb{C}})$. A root $\alpha \in \Phi$ is called compact (respectively noncompact) if its root space lies in $\mathfrak{k}^{\mathbb{C}}$ (respectively the orthogonal complement of $\mathfrak{k}^{\mathbb{C}}$). The differentials of the characters of B form a lattice $\Lambda \subset i\mathfrak{b}^* (\mathfrak{b}^* = \text{dual space of } \mathfrak{b})$. The killing form induces a positive definite inner product $(,)$ on $i\mathfrak{b}^*$. An element $\lambda \in \Lambda$ is called nonsingular if $(\lambda, \alpha) \neq 0$ for every $\alpha \in \Phi$. We set $W = W(G, B) =$ Weyl group of B in G . Equivalently, W can be described as the group generated by the reflection about the compact roots in $i\mathfrak{b}^*$.

In order to state Harish-Chandra's enumeration of the discrete series [3], we assume, without loss of generality, that G is acceptable in the sense of Harish-Chandra. Then, for each nonsingular $\lambda \in \Lambda$, there exists exactly one tempered¹ invariant eigendistribution θ_λ on G , such that

$$\theta_\lambda|_{B \cap G'} = (-1)^q \frac{\sum_{w \in W} \text{sgn } w e^{w\lambda}}{\prod_{\alpha \in \Phi, (\alpha, \lambda) > 0} (e^{\alpha/2} - e^{-\alpha/2})}.$$

Here $q = \frac{1}{2} \dim G/K$, and G' = set of regular semisimple points in G . Every θ_λ is the character of a discrete series representation, and conversely. Moreover, $\theta_\lambda = \theta_\mu$ if and only if λ belongs to the W -orbit of μ .

Summary of doctoral dissertation, submitted to the Department of Mathematics at Columbia University in 1977; received by the editors October 2, 1979.

AMS (MOS) subject classifications (1970). Primary 22E45.

Key words and phrases. Semisimple Lie groups, representations, discrete series, character formulas.

¹A distribution θ on G is tempered if it extends to the Schwartz space of rapidly decreasing functions [3].

When G/K is a hermitian symmetric space, there exist positive root systems in Φ such that the sum of two noncompact positive roots is never a root. If λ is dominant with respect to such a positive root system, θ_λ is the character of one of the so-called holomorphic discrete series representations. In this special situation, S. Martens [5] and H. Hecht [4] have given explicit global formulas for the characters θ_λ . Whether or not G/K is hermitian symmetric, there exist positive root systems which satisfy the following condition [1]:

for each noncompact simple factor G_i of G , there exists exactly one noncompact simple root β_i , and this root β_i occurs at most twice in the highest root of G_i .

(Borel-de Siebenthal property). The problem of computing the discrete series characters θ_λ globally can be reduced, at least in principle, to the following rather special situation:

- (a) G is simple and has both a compact Cartan subgroup B and a split Cartan subgroup A .
- (b) Compute θ_λ restricted to the identity component of a split Cartan subgroup A .
- (c) The system of positive roots $\Psi = \{\alpha \in \Phi \mid \langle \alpha, \lambda \rangle > 0\}$ has the Borel-de Siebenthal property. (See Schmid [6].)

From now on, let G, λ, Ψ, A be as in (a)–(c), and d an inner automorphism of $\mathfrak{g}^{\mathbb{C}}$ such that $d : \mathfrak{b}^{\mathbb{C}} \xrightarrow{\sim} \mathfrak{a}^{\mathbb{C}}$.

We denote the identity component of A by A° and define

$$C = \{\exp X \mid X \in \mathfrak{a}, \langle \alpha, d^{-1}X \rangle < 0 \text{ for all } \alpha \in \Psi\}.$$

The closure of C and its conjugates cover A° . Our main result provides an explicit formula for the restrictions of θ_λ to A° . This formula involves a particular element t of the Weyl group of $(\mathfrak{g}^{\mathbb{C}}, \mathfrak{b}^{\mathbb{C}})$, whose description we defer until later.

THEOREM. *Let W_U be the subgroup of W generated by the compact simple roots for Ψ . Then*

$$\theta_\lambda|_C = (-1)^q \frac{|W|}{|W_U|} \frac{\sum_{w \in W} \text{sgn}(tw) e^{t w \lambda}}{\prod_{\alpha \in \Psi} (e^{\alpha/2} - e^{-\alpha/2})} \circ d^{-1}$$

Let β be the noncompact simple root for Ψ . Then β is as long as or longer than any noncompact root, and the system Φ' of the roots in Φ orthogonal to β has at most three irreducible components. Moreover if Φ' has more than one connected component, then all but perhaps one are of type A_1 , and all the A_1 -type components consist of noncompact roots. Two roots $\alpha_1, \alpha_2 \in \Phi$ are said to be strongly orthogonal if $\alpha_1 \pm \alpha_2 \notin \Phi$. For any strongly orthogonal subset $S \subset \Phi$ consisting of noncompact roots, we set $\Phi_S = \mathbb{Q}$ -linear span of S in Φ .

LEMMA 1. *Each irreducible component of $\Psi \cap \Phi_S$ has the Borel-de Siebenthal property.*

We now define a family of sub-root systems $\Phi = \Phi^0, \Phi^1, \dots, \Phi^m$ inductively, as follows: Φ^{i+1} is the set of roots in Φ^i orthogonal to the noncompact simple roots for $\Psi \cap \Phi^i$, until the process stops. Let S_0 be the set consisting of all positive noncompact roots that are simple roots for some $\Psi \cap \Phi^i, 0 \leq i \leq m$.

LEMMA 2. (a) S_0 is a strongly orthogonal set which spans Φ over \mathbf{Q} .

(b) S_0 contains at most one short root.

The next proposition describes the element t of the Weyl group of $(\mathfrak{g}^{\mathbf{C}}, \mathfrak{b}^{\mathbf{C}})$ which was used in the statement of the main result.

PROPOSITION. *There exists a unique t in the Weyl group of $(\mathfrak{g}^{\mathbf{C}}, \mathfrak{b}^{\mathbf{C}})$ such that*

- (1) t is a product of reflections about roots in S_0 .
- (2) If β occurs twice in the highest root, then $t \neq 1$.
- (3) t takes any long simple root into a noncompact root.
- (4) $(t w \lambda, \mu) \geq 0$ for every $w \in W_U$ and λ, μ dominant integral with respect to Ψ .

From the Proposition, one can deduce the following properties of t :

- (a) If α_1, α_2 are two adjacent long simple roots, then $\text{sgn } t \alpha_1 \neq \text{sgn } t \alpha_2$.
- (b) Assume β occurs twice in the highest root. Then t fixes any short root in S_0 .
- (c) If S_0 does not contain short roots, $t \alpha = \alpha$ for any short simple root α .
- (d) Again under the assumption that β occurs twice in the highest root, if S_0 does contain short roots, $t \alpha$ is noncompact for any short simple root α .

The proof of the theorem proceeds by induction on the dimension of G . The crux of the matter is to verify the consistency of our formula with Harish-Chandra's matching conditions [2]. Details will appear elsewhere.

Finally, I would like to express my deep gratitude to Professor Wilfried Schmid for his advice and interest.

REFERENCES

1. A. Borel, et J. de Siebenthal, *Les sous-groupes fermés de rang maximum des groupes de Lie clos*, Comm. Math. Helv. **23** (1949), 200–221.
2. Harish-Chandra, *Invariant eigendistributions on a semi-simple Lie group*, Trans. Amer. Math. Soc. **119** (1965), 457–508.
3. ———, *Discrete series for semi-simple Lie groups. II*, Acta Math. **116** (1966), 1–111.
4. H. Hecht, *The characters of Harish-Chandra representations*, Math. Ann. **219** (1976), 213–226.
5. S. Martens, *The characters of the holomorphic discrete series*, Proc. Nat. Acad. Sci. U.S.A. vol. 72, no. 9, 1975, pp. 3275–3276.
6. W. Schmid, *On the characters of the discrete series*, Invent. Math. **30** (1975), 47–144.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF CORDOBA,
CORDOBA, ARGENTINA