

# RESEARCH ANNOUNCEMENTS

BULLETIN (New Series) OF THE  
AMERICAN MATHEMATICAL SOCIETY  
Volume 12, Number 2, April 1985

## EQUIVARIANT $h$ -COBORDISMS AND FINITENESS OBSTRUCTIONS

BY MARK STEINBERGER AND JAMES WEST<sup>1</sup>

**ABSTRACT.** We classify up to topological equivalence those equivariant  $h$ -cobordisms which admit a handle structure, giving a topologically invariant Whitehead group and an  $s$ -cobordism theorem, and giving the comparison to the Diff and PL classification via an equivariant version of the controlled Whitehead groups of Chapman and Quinn. We also construct stably triangulable, compact  $G$ -manifolds with boundary which realize arbitrary controlled equivariant finiteness obstructions. The controlled finiteness obstructions of their boundaries are generic for closed  $G$ -manifolds whose product with  $\mathbf{R}$  is triangulable.

Here  $G$  is a finite group, and  $G$ -manifolds are assumed to be locally linear with codimension-3 gaps (proper inclusions of fixed-point components have codimension  $\geq 3$ ). By a  $G$ - $h$ -cobordism on  $M$  we mean a proper equivariant  $h$ -cobordism which admits a handle structure (equivariant) in which no handles are attached to fixed-point components of  $M$  of dimension less than 5.

Let  $\text{Wh}_G^{\text{PL}}(M)$  be the locally compact version as in [Si] of Illman's Whitehead group  $[\mathbf{II}_1]$ , and let  $\text{Wh}_G^{\text{PL},\rho}(M)$ , be the subgroup consisting of pairs  $(Y, M)$  such that  $Y_\alpha^H = M_\alpha^H \cup Y_\alpha^{>H}$  if either  $M_\alpha^H = M_\alpha^{>H}$  or  $\dim M_\alpha^H < 5$  for each component  $M_\alpha^H$  of  $M^H$ . With our assumptions,  $G$ - $h$ -cobordisms with an explicit handle structure are classified up to handle manipulation (or Cat isomorphism of  $\text{Cat} = \text{Diff}$  or PL) by  $\text{Wh}_G^{\text{PL},\rho}(M)$  (cf. [BQ, R]). This gives an  $s$ -cobordism theorem in Diff and PL, but as noted by [II<sub>2</sub>, BH, R] and [DR], any  $G$ - $h$ -cobordism whose torsion lies in  $\text{Wh}_G^{\text{PL}}(x) \subset \text{Wh}_G^{\text{PL}}(M)$ , for  $x \in M^G$ , is topologically trivial.

We say that two  $G$ -pairs  $(Y, X)$  and  $(Z, X)$  are stably homeomorphic if there is a homeomorphism  $Y \times Q_G \cong Z \times Q_G$  commuting up to proper  $G$ -homotopy with the inclusions of  $X$ , where  $Q_G$  is the product of infinitely many copies of the unit disc in the regular representation of  $G$  over  $\mathbf{R}$ . The

---

Received by the editors November 2, 1984.

1980 *Mathematics Subject Classification*. Primary 57S17, 57Q10.

<sup>1</sup>Partially supported by NSF.

©1985 American Mathematical Society  
0273-0979/85 \$1.00 + \$.25 per page

stable homeomorphism classes of relative CW pairs  $(Y, X)$ , with  $X \subset Y$  a proper  $G$ -equivalence give a quotient group  $\text{Wh}_G^{h\text{-Top}}(X)$  of  $\text{Wh}_G^{\text{PL}}(X)$ .

**THEOREM 1 (EQUIVARIANT TOPOLOGICAL  $s$ -COBORDISM THEOREM).** *Two  $G$ - $h$ -cobordisms on  $M$  (with handle structure as above) are  $G$ -homeomorphic rel  $M$  if and only if their torsions agree in  $\text{Wh}_G^{h\text{-Top}}(M)$ .*

By a  $G$ -CE map  $f: Y \rightarrow Z$  we mean a proper  $G$ -map for which  $f^{-1}z \subset U$  is  $G_z$ -nullhomotopic for each  $G_z$ -neighborhood  $U$  of  $f^{-1}z$  and each  $z \in Z$ . A  $G$ -CEPL map is one which is both  $G$ -CE and PL.

**COROLLARY.**  $\text{Wh}_G^{h\text{-Top}}(M) = \text{Wh}_G^{\text{PL}}(M) / \sim$ , where  $\sim$  is generated by  $G$ -CE maps rel  $M$ .

Let  $\text{WH}_G^{\text{CEPL}}(M) = \text{Wh}_G^{\text{PL}}(M) / \sim$ , where  $\sim$  is generated by  $G$ -CEPL maps rel  $M$ . Then  $\text{Wh}_G^{\text{PL}}(M) \rightarrow \text{Wh}_G^{h\text{-Top}}(M)$  factors through  $\text{Wh}_G^{\text{CEPL}}(M)$ , and the kernel of  $\text{Wh}_G^{\text{PL}}(M) \rightarrow \text{Wh}_G^{\text{CEPL}}(M)$  is  $\sum_{f: G/H \rightarrow M} f_* \text{Wh}_G^{\text{PL}}(G/H)$ , where the sum ranges over all maps of orbits into  $M$ . Thus all previously known examples of torsions of topologically trivial  $G$ - $h$ -cobordisms die in  $\text{Wh}_G^{\text{CEPL}}(M)$ .

Let  $\mathcal{E} = \text{PL}$  or  $\text{CEPL}$ . We compute the kernel of  $\text{Wh}_G^{\mathcal{E}}(M) \rightarrow \text{Wh}_G^{h\text{-Top}}(M)$  by an equivariant generalization of Chapman’s controlled Whitehead groups [C]. For a  $G$ -map  $p: X \rightarrow B$ , with  $B$  metric, let  $\text{Wh}_G^{\text{CEPL}}(X)_{p^{-1}\varepsilon}$  ( $\varepsilon$  a  $G$ -majorant of  $B$ ) be the fully equivariant analogue of Chapman’s group, and let  $\text{Wh}_G^{\text{PL}}(X)_{p^{-1}\varepsilon}$  be obtained by strengthening the basic equivalence relation from diagrams of  $G$ -CEPL maps to those of  $G$ -simple maps (point inverses have the equivariant simple homotopy type of points). When  $p = 1_X$  we denote these groups by  $\text{Wh}_G^{\mathcal{E}}(X)_{\varepsilon}$ . Restricted groups  $\text{Wh}_G^{\mathcal{E}, \rho}(M)_{\varepsilon}$  are defined as above.

**THEOREM 2.** *For  $\mathcal{E} = \text{PL}$  or  $\text{CEPL}$  there is an exact sequence*

$$\varinjlim_{\varepsilon} \text{Wh}_G^{\mathcal{E}, \rho}(M)_{\varepsilon} \rightarrow \text{Wh}_G^{\mathcal{E}, \rho}(M) \rightarrow \text{Wh}_G^{h\text{-Top}, \rho}(M) \rightarrow 0,$$

where  $\text{Wh}_G^{h\text{-Top}, \rho}(M)$  is the image of  $\text{Wh}_G^{\text{PL}, \rho}(M)$  in  $\text{Wh}_G^{h\text{-Top}}(M)$ .

We recover equivariant versions of all of the results of [C], with the obstructions groups for the Thin  $h$ -Cobordism Theorem and End Theorem given by the controlled PL Whitehead and  $\tilde{K}_0$  groups. However, unlike the inequivariant case,  $\varinjlim_{\varepsilon} \text{Wh}_G^{\mathcal{E}}(X)_{\varepsilon}$  is rarely trivial.

Here,  $\tilde{K}_{i_G}(X)_{p^{-1}\varepsilon} \subset \text{Wh}_G^{\text{PL}}(X \times T^{1-i})_{p^{-1}\varepsilon}$  (or, equivalently,  $\text{CEPL}$ ) is the subgroup of elements invariant under all of the transfers from the  $S^1$  factors of the  $(1-i)$ -torus  $T^{1-i}$  (with the trivial action) for  $i \leq 0$ . (The tilde is optional for  $i < 0$ .) The uncontrolled versions,  $\tilde{K}_{i_G}(X)$ , are isomorphic to those obtained from the Bass-Heller-Swan splitting when  $X$  is compact (cf. [Ran]). As in [Q<sub>1</sub>, Q<sub>2</sub>], all of these inverse systems are stable when  $p$  is an equivariant simplicial  $p$ -NDR [Q<sub>2</sub>] (e.g.,  $p = 1_X$ ,  $X$  a locally compact  $G$ -ANR, or  $p$  a  $G$ -simplicial map) and may be computed by a Leray spectral sequence with coefficients in the Whitehead and  $K$ -groups of  $p^{-1}$  (orbits) when  $p$  is simplicial. We obtain the following.

**THEOREM 3.** *Let  $X$  be a locally compact  $G$ -ANR. Then  $\varinjlim_{\varepsilon} K_{i_G}(X)_{\varepsilon} = 0$  for  $i < -1$ ,  $\varinjlim_{\varepsilon} K_{-1_G}(X)_{\varepsilon} \simeq H_0^{G,lf}(X : K_{-1_G})$ , and there is an exact sequence*

$$\begin{aligned} H_3^{G,lf}(X; K_{-1_G}) &\rightarrow H_1^{G,lf}(X; \tilde{K}_{0_G}) \rightarrow \varinjlim_{\varepsilon} \text{Wh}_G^{\text{CEPL}}(X)_{\varepsilon} \rightarrow H_2^{G,lf}(X; K_{-1_G}) \\ &\rightarrow H_0^{G,lf}(X; \tilde{K}_{0_G}) \rightarrow \varinjlim_{\varepsilon} \tilde{K}_{0_G}(X)_{\varepsilon} \rightarrow H_1^{G,lf}(X; K_{-1_G}) \rightarrow 0. \end{aligned}$$

Here, we take Bredon homology with locally finite chains with coefficient systems given by restriction of the functors  $\tilde{K}_{i_G}$  to orbits.

**COROLLARY.**  $\text{Wh}_G^{h\text{-Top}}(X)$  and  $\varinjlim_{\varepsilon} \tilde{K}_{0_G}(X)_{\varepsilon}$  are functors of the  $\pi_1$ -system of  $X$  for  $X$  compact.

As a sample computation, note that if  $X$  has the  $\pi_1$ -system of the  $n$ -torus with trivial action, then  $\text{Wh}_G^{h\text{-Top}}(X)$  is isomorphic to the Nil terms in

$$\text{Wh}_G^{\text{PL}}(X) \simeq \text{Wh}_G^{\text{PL}}(*) \oplus n\tilde{K}_{0_G}(*) \oplus \binom{n}{2} K_{-1_G}(*) \oplus \text{Nil terms}.$$

For any compact  $G$ -ANR  $X$  there is a well-defined controlled finiteness obstruction  $\varinjlim_{\varepsilon} \sigma_{\varepsilon}(X) \in \varinjlim_{\varepsilon} \tilde{K}_{0_G}(X)_{\varepsilon}$  which maps to the ordinary equivariant finiteness obstruction  $[\mathbf{A}] \sigma(X)$  under the natural map  $\varinjlim_{\varepsilon} \tilde{K}_{0_G}(X)_{\varepsilon} \rightarrow \tilde{K}_{0_G}(X)$ . Examples of compact  $G$ -manifolds  $M$  with  $\sigma(M) \neq 0$  have been given by **[Q<sub>1</sub>, DR]** and others. These examples all have finiteness obstruction in  $\tilde{K}_{0_G}(*) \subset \tilde{K}_{0_G}(M)$ ,  $* \in M^G$ .

**THEOREM 4.** *Let  $K$  be a finite  $G$ -complex and let  $x \in \varinjlim_{\varepsilon} \tilde{K}_{0_G}(K)_{\varepsilon}$ . Then there is a compact  $G$ -manifold  $M$  containing  $K$  as a  $\pi_1$ -equivalent retract,  $r: M \rightarrow K$ , such that  $r_*(\varinjlim_{\varepsilon} \sigma_{\varepsilon}(M)) = x$  and  $M \times \mathbf{R}$  is triangulable.*

For closed manifolds  $M$ , with  $M \times \mathbf{R}$  triangulable,  $\varinjlim_{\varepsilon} \sigma_{\varepsilon}(M)$  must have the form  $\tau \pm \bar{\tau}$  for some  $\tau \in \varinjlim_{\varepsilon} \tilde{K}_{0_G}(M)_{\varepsilon}$ , where  $\bar{\tau}$  is the suitably signed conjugate of  $\tau$ . This is satisfied generally by  $\partial(M \times I^n)$ ,  $n \geq 3$ , for the examples  $M$  of Theorem 4.

David Webb **[W]** has shown that  $\varinjlim_{\varepsilon} \tilde{K}_{0_G}(X)_{\varepsilon} \rightarrow \tilde{K}_{0_G}(X)$  is not always injective, so there exist  $G$ -finite  $G$ -manifolds which do not admit a handle structure. This provides a negative answer to Question 4.4 of **[Sch]**.

REFERENCES

**[A]** D. R. Anderson, *Torsion invariants and actions of finite groups*, Michigan Math. J. **29** (1982), 27–42.  
**[BH]** W. Browder and W.-C. Hsiang, *Some problems on homotopy theory, manifolds and transformation groups*, Proc. Sympos. Pure Math., vol. 32, part II, Amer. Math. Soc., Providence, R.I., 1978, pp. 251–267.  
**[BQ]** W. Browder and F. Quinn, *A surgery theory for  $G$ -manifolds and stratified sets*, Manifolds (Tokyo, 1973), Univ. of Tokyo Press, Tokyo, 1975, pp. 27–36.

[C] T. A. Chapman, *Controlled simple homotopy theory*, Lecture Notes in Math., vol. 1009, Springer-Verlag, 1983.

[DR] K. H. Doverman and M. Rothenberg, *An equivariant surgery sequence and equivariant diffeomorphism and homeomorphism classification* (preprint).

[Il<sub>1</sub>] S. Illman, *Whitehead torsion and group actions*, Ann. Acad. Sci. Fenn. Ser. A **588** (1974), 1–44.

[Il<sub>2</sub>] ———, Personal communication.

[Q<sub>1</sub>] F. Quinn, *Ends of maps. II*, Invent. Math. **68** (1982), 353–424.

[Q<sub>2</sub>] ———, *Geometric algebra and ends of maps*, Notes prepared for CBMS Conf., Notre Dame, 1984.

[Ran] A. Ranicki, *Algebraic and geometric splittings of the K- and L-groups of polynomial extensions* (preprint).

[R] M. Rothenberg, *Torsion invariants and finite transformation groups*, Proc. Sympos. Pure Math., vol. 32, Amer. Math. Soc., Providence, R.I., 1978, pp. 267–311.

[Si] L. C. Siebenmann, *Infinite simple homotopy types*, Indag. Math. **32** (1970), 479–495.

[Sch] R. Schultz (Editor), *Problems submitted to the AMS Summer Research Conference on Group Actions*, Boulder, 1983 (preprint).

[W] D. Webb, *Equivariantly finite manifolds with no handle structure* (in preparation).

DEPARTMENT OF MATHEMATICS, NORTHERN ILLINOIS UNIVERSITY, DEKALB,  
ILLINOIS 60115

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON,  
NEW JERSEY 08540