

On the Nonrelativistic Limit of the Dirac Theory

W. Hunziker

Institut für Theoretische Physik der ETH, Zürich, Switzerland

Received October 15, 1974

Abstract. The relation between a “nonrelativistic” Hamiltonian of the form $H^\infty = (A + B)^2 + C$ and a corresponding family of “Dirac-Hamiltonians” $H(c)$ in the limit $c \rightarrow \infty$ is investigated. It is shown that the resolvent $(z - H(c))^{-1}$ and the relativistic perturbation of isolated eigenvalues of H^∞ are analytic in $1/c$ for sufficiently large $|c|$.

1. Introduction

The Hamiltonian of a Dirac-electron of charge $e = 1$ and mass $m = 1/2$ may be written as

$$H(c) = c\alpha(\mathbf{p} - \mathbf{A}(\mathbf{x})) + \frac{1}{2}\beta c^2 + \varphi(\mathbf{x}), \quad (1)$$

where $\mathbf{p} = -i\mathbf{d}/d\mathbf{x}$ and with the 4×4 -matrices

$$\alpha = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} \mathbb{1} & 0 \\ 0 & \mathbb{1} \end{pmatrix},$$

whose elements are the 2×2 -matrices $\mathbb{1}$ and $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3) =$ set of Pauli spin-matrices. $\mathbf{A}(\mathbf{x})$ and $\varphi(\mathbf{x})$ are the potentials of the static electromagnetic field. The usual factor $1/c$ in front of $\mathbf{A}(\mathbf{x})$ is omitted on purpose since it must be kept fixed in the nonrelativistic limit $c \rightarrow \infty$. $H(c)$ acts on the Hilbertspace $C^4 \otimes L^2(\mathbb{R}^3)$ of square-integrable 4-component wave functions.

On a formal level, it is well understood that the nonrelativistic limit $c \rightarrow \infty$ is described by the Pauli-Hamiltonian

$$H^\infty = (\boldsymbol{\sigma}(\mathbf{p} - \mathbf{A}(\mathbf{x}))^2 + \varphi(\mathbf{x})) \quad (2)$$

on the smaller Hilbertspace $C^2 \otimes L^2(\mathbb{R}^3)$, and there exists a systematic scheme for obtaining corrections to H^∞ in the form of a power series in $1/c$ [1]. However, these “relativistic perturbations” of H^∞ are given by more and more singular operators which are by no means small with respect to H^∞ . One might therefore suspect that perturbation expansions in powers of $1/c$ are at best asymptotic.

Nevertheless, Titchmarsh [2] has proved analyticity in $1/c$ of eigenvalues and eigenfunctions for the spherically symmetric case without magnetic field: $\varphi = \varphi(r)$, $\mathbf{A} = 0$; and Veselić [3] has extended this result to the case without spherical symmetry: $\varphi = \varphi(\mathbf{x})$, $\mathbf{A} = 0$.

In this note we investigate the general case $\mathbf{A} \neq 0$ which poses essentially new problems—already in the nonrelativistic limit. One of the points we wish to make is that it is profitable to treat a general Hamiltonian of type $H^\infty = (A + B)^2 + C$ as a nonrelativistic limit of a corresponding Dirac-Hamiltonian $H(c)$.

In order to keep the conditions on A and φ fairly general, we here restrict ourselves to the discrete spectrum of H . The analyticity properties of the resolvent in $1/c$ will of course also be needed in the discussion of the continuum. However, additional assumptions for the electromagnetic field (like dilatation-analyticity or sufficiently rapid fall-off at infinity) are then necessary and the arguments become more technical [4].

2. The Hamiltonian and Its Spectrum

We summarize some (but not all!) known results on the selfadjointness and on the spectrum of $H(c)$. This is intended only as a background for the more general set-up introduced in Section 3.

H is of the form $H_0 + V$ with $H_0 = c\alpha p + \frac{1}{2}c^2\beta$, $V(x) = \varphi(x) - c\alpha A(x)$. $V(x)$ is a 4×4 -matrixvalued function on $R^3 \cdot L^p$ -norms of V may be defined with respect to any matrix-norm.

Theorem 1. *Let $V \in L^p + L^\infty$ for some $p > 3$. Then V is H_0 -bounded with arbitrarily small relative bound. Therefore, $H = H_0 + V$ is selfadjoint with domain $D(H_0)$.*

Theorem 2. *Suppose that $V \in L^p + \varepsilon L^\infty$ ($p > 3$), i.e. that the L^∞ -part of V can be chosen arbitrarily small in L^∞ -norm. Then the spectrum $\sigma(H)$ of H consists of the continuum $\sigma(H_0) = \{z \in R : |z| \geq \frac{1}{2}c^2\}$ and, in the complement of $\sigma(H_0)$, of isolated eigenvalues with finite multiplicities which can accumulate only at $+\frac{1}{2}c^2$.*

Remarks. Theorem 1 is proved in [5] and follows from the fact that an operator of the form

$$f(x) (1 + p^2)^{-1/2} \tag{3}$$

on $L^2(R^3)$ is bounded if $f \in L^p$, $p > 3$. The condition $p > 3$ excludes Coulomb-like singularities. However, $f(x) = |x|^{-1}$ is still relatively bounded with respect to $|p|$ [6].

Theorem 2 can be proved like its analogue for Schrödinger Hamiltonians [7]. The main point is that (3) is a compact operator if $f \in L^p + \varepsilon L^\infty$, i.e. V is relatively compact with respect to H_0 .

3. The Nonrelativistic Limit

We now pose the problem in a generalized form. Let A, B, C be symmetric operators on a Hilbertspace \mathcal{H} . On $C^2 \otimes \mathcal{H}$ we define the “Dirac-Hamiltonian”

$$H(c) = c\alpha \otimes (A + B) + \frac{c^2}{2} \beta \otimes \mathbb{1} + \mathbb{1} \otimes C,$$

where

$$\alpha = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \beta = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

We want to show that the limit $c \rightarrow \infty$ is described by the “Pauli-Hamiltonian”

$$H^\infty = (A + B)^2 + C \quad \text{on } \mathcal{H}.$$

Corresponding to the hypothesis of Theorem 1 we assume:

$$\begin{aligned} A &= A^*; B \text{ and } C \text{ are } A\text{-bounded, in particular} \\ B &\text{ has relative bound } < 1 \text{ with respect to } A. \end{aligned} \tag{4}$$

It follows easily that $H(c)$ is selfadjoint with domain $D(A)$ for c real and sufficiently large. For H^∞ we have

Lemma 1. H^∞ is selfadjoint with domain $D((A + B)^2)$ and bounded below.

Proof. By (4), $A + B$ is selfadjoint with domain $D(A)$ and A is $(A + B)$ -bounded. Hence C is $(A + B)$ -bounded and has therefore arbitrarily small relative bound with respect to $(A + B)^2$.

Remark. For H^∞ , the splitting

$$H^\infty = A^2 + (AB + BA + B^2 + C)$$

into unperturbed part plus perturbation is artificial and raises unnecessary domain questions. These will be avoided automatically by treating H^∞ as a limit of $H(c)$.

We first discuss the unperturbed resolvent $(z - H_0(c))^{-1}$ for

$$H_0(c) = c\alpha A + \frac{1}{2}c^2\beta,$$

where we have dropped the tensor-product notation. From $\alpha^2 = \beta^2 = 1$ and $\alpha\beta + \beta\alpha = 0$ it follows that

$$H_0^2(c) = c^2 A^2 + \frac{1}{4}c^4,$$

which shows that $\sigma(H_0)$ has at least the gap $(-\frac{1}{2}c^2, +\frac{1}{2}c^2)$. For $z \notin \sigma(H_0)$ we have

$$\begin{aligned} (z - H_0)^{-1} &= (z + H_0)(z^2 - H_0^2)^{-1} \\ &= (z + c\alpha A + \frac{1}{2}c^2\beta)(z^2 - \frac{1}{4}c^4 - c^2 A^2)^{-1}. \end{aligned}$$

Before taking the nonrelativistic limit $c \rightarrow \infty$ we must subtract from $H_0(c)$ or $H(c)$ the rest energy $\frac{1}{2}c^2$ or, equivalently, replace in the resolvents z by $z + \frac{1}{2}c^2$. This will always be assumed in the following. It is also convenient to use 2-component notation: $u = \begin{pmatrix} u^1 \\ u^2 \end{pmatrix}$, $u_k \in \mathcal{H}$, for vectors $u \in C^2 \otimes \mathcal{H}$ and the corresponding 2×2 -matrix notation for operators on $C^2 \otimes \mathcal{H}$. The unperturbed resolvent then takes the form

$$\begin{aligned} G_0(z, c) &= (z - H_0(c))^{-1} \\ &= \begin{pmatrix} 1 + \frac{z}{c^2} & \frac{A}{c} \\ \frac{A}{c} & z \end{pmatrix} \left(z + \frac{z^2}{c^2} - A^2 \right)^{-1}, \end{aligned}$$

which shows explicitly that, for $z \notin \sigma(A^2)$, $G_0(z, c)$ is analytic in $1/c$ in a z -dependent neighbourhood of $1/c = 0$. To construct the full resolvent $G(z, c) = (z - H(c))^{-1}$ we

start from the resolvent equation

$$G(z, c) = G_0(z, c) + K(z, c) G(z, c),$$

where

$$K(z, c) = G_0(z, c) (H(c) - H_0(c)) \\ = \left(z + \frac{z^2}{c^2} - A^2 \right)^{-1} \begin{pmatrix} \left(1 + \frac{z}{c^2} \right) C + AB & \left(1 + \frac{z}{c^2} \right) cB + \frac{AC}{c} \\ \frac{1}{c} (AC + zB) & AB + \frac{zC}{c^2} \end{pmatrix}.$$

This expression for K must be understood in the following sense: a term like $\left(z + \frac{z^2}{c^2} - A^2 \right)^{-1} AB$ is defined on $D(B)$ as the product of the bounded operator $A \left(z + \frac{z^2}{c^2} - A^2 \right)^{-1}$ with B . As a consequence of (4), K is therefore defined on $D(A)$ and bounded. In the following we denote with K the unique bounded extension of this operator to all of $C^2 \otimes \mathcal{H}$. For $B \neq 0$ we see that K diverges as $c \rightarrow \infty$. To control this divergence we set

$$S(c) = \begin{pmatrix} 1 & 0 \\ 0 & c \end{pmatrix}$$

and introduce

$$\tilde{G}_0(z, c) = S G_0 S^{-1} = \left(z + \frac{z^2}{c^2} - A^2 \right)^{-1} \begin{pmatrix} 1 + \frac{z}{c^2} & \frac{A}{c^2} \\ A & \frac{z}{c^2} \end{pmatrix} \\ = (z - A^2)^{-1} \begin{pmatrix} 1 & 0 \\ A & 0 \end{pmatrix} \quad \text{for } c = \infty. \quad (5)$$

$$\tilde{K}(z, c) = S K S^{-1} = \left(z + \frac{z^2}{c^2} - A^2 \right)^{-1} \begin{pmatrix} \left(1 + \frac{z}{c^2} \right) C + AB & \left(1 + \frac{z}{c^2} \right) B + \frac{AC}{c^2} \\ AC + zB & AB + \frac{zC}{c^2} \end{pmatrix} \\ = (z - A^2)^{-1} \begin{pmatrix} C + AB & B \\ AC + zB & AB \end{pmatrix} \quad \text{for } c = \infty. \quad (6)$$

We notice that for $z \notin \sigma(A^2)$, $\tilde{G}_0(z, c)$ and $\tilde{K}(z, c)$ are analytic in $(1/c)^2$ in a neighbourhood of zero. The resolvent equation transforms into

$$\tilde{G}(z, c) = \tilde{G}_0(z, c) + \tilde{K}(z, c) \tilde{G}(z, c) \quad (7)$$

for $\tilde{G} = S G S^{-1}$. Our next task is to connect this equation for $c = \infty$ with the Pauli Hamiltonian H^∞ .

Theorem 3. *Let A, B, C satisfy (4) and let $z \notin \sigma(A^2)$. Then the two equations*

$$(z - H^\infty)u = v \tag{8}$$

and

$$\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} (z - A^2)^{-1}v \\ A(z - A^2)^{-1}v \end{pmatrix} + \tilde{K}(z, \infty) \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \tag{9}$$

are equivalent in the following sense: (8) implies (9) for $\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} u \\ (A + B)u \end{pmatrix}$, (9) implies $u_1 \in D((A + B)^2)$, $u_2 = (A + B)u_1$ and (8) for $u = u_1$.

The straightforward but somewhat lengthy proof is given in Section 5. As a corollary we note that for $z \notin \sigma(A^2)$, $z \notin \sigma(H^\infty)$, $G^\infty(z) = (z - H^\infty)^{-1}$ satisfies

$$\begin{pmatrix} G^\infty(z) & 0 \\ (A + B)G^\infty(z) & 0 \end{pmatrix} = \begin{pmatrix} (z - A^2)^{-1} & 0 \\ A(z - A^2)^{-1} & 0 \end{pmatrix} + \tilde{K}(z, \infty) \begin{pmatrix} G^\infty(z) & 0 \\ (A + B)G^\infty(z) & 0 \end{pmatrix}. \tag{10}$$

Conversely, if $z \notin \sigma(A^2)$ and if $(1 - \tilde{K}(z, \infty))^{-1}$ exists and is bounded, it follows that $z \notin \sigma(H^\infty)$ and that

$$\begin{pmatrix} G^\infty(z) & 0 \\ (A + B)G^\infty(z) & 0 \end{pmatrix}$$

is the unique solution of (10) in $L(C^2 \otimes \mathcal{H})$. Therefore, (10) is a suitable resolvent equation for H^∞ which may also be used, incidentally, as a starting point for time-independent scattering theory [4]. Corresponding to the hypothesis of theorem 2 we now assume in addition to (4) that

$$B \text{ and } C \text{ are relatively compact with respect to } A. \tag{11}$$

As in Theorem 2 it then follows that $H(c)$ can only have isolated eigenvalues of finite multiplicities in the complement of $\sigma(H_0(c))$, in particular in the gap $(-c^2, 0)$ (rest energy subtracted). Since $\tilde{K}(z, \infty)$ is compact for $z \notin \sigma(A^2)$ we obtain a similar result for H^∞ :

Lemma 2. *In the complement of $\sigma(A^2)$, $\sigma(H^\infty)$ consists only of isolated eigenvalues of finite multiplicities which are bounded below.*

Proof. Suppose that for some $z \notin \sigma(A^2)$, the homogeneous equation

$$\tilde{K}(z, \infty)u = u \tag{12}$$

has a nontrivial solution. Then $u = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$ satisfies (9) with $v = 0$. By Theorem 3, it follows that $u_1 \in D((A + B)^2)$, $u_2 = (A + B)u_1$ and

$$(z - H^\infty)u_1 = 0. \tag{13}$$

We conclude that $u_1 \neq 0$ and that $z \in \sigma(H^\infty)$. By the Fredholm alternative, therefore, $(1 - \tilde{K}(z, \infty))^{-1}$ exists for $z \notin \sigma(H^\infty)$. Since $\tilde{K}(z, \infty)$ is analytic in $z \notin \sigma(A^2)$,

$(1 - \tilde{K}(z, \infty))^{-1}$ and therefore $G^\infty(z)$ are meromorphic in $z \notin \sigma(A^2)$. Conversely, (13) implies (12) for $u = \begin{pmatrix} u_1 \\ (A+B)u_1 \end{pmatrix}$, hence the eigenvalues of H^∞ in the complement of $\sigma(A^2)$ are of finite multiplicity. Boundedness below is obvious from Lemma 1. This concludes the proof.

We are now prepared to discuss the analyticity properties in $1/c$ of $G(z, c)$. Let $z \notin \sigma(A^2) \cup \sigma(H^\infty)$. Then $(1 - \tilde{K}(z, \infty))^{-1}$ exists. Since $\tilde{K}(z, c)$ is analytic in $(1/c)^2$ it follows that $(1 - \tilde{K}(z, c))^{-1}$ exists and is analytic in $(1/c)^2$ for $|c|$ sufficiently large. The same is true for $\tilde{G}_0(z, c)$ and therefore, by (7), for $\tilde{G}(z, c)$. From (5) and (10) we see that the power series of $\tilde{G}(z, c)$ in $(1/c)^2$ begins with

$$\tilde{G}(z, c) = \begin{pmatrix} G^\infty(z) & 0 \\ (A+B)G^\infty(z) & 0 \end{pmatrix} + o\left(\frac{1}{c^2}\right).$$

Due to the particular form of the leading term, $G(z, c) = S^{-1}(c)\tilde{G}(z, c)S(c)$ is still analytic in $1/c$ with an expansion

$$G(z, c) = \begin{pmatrix} G^\infty(z) & 0 \\ 0 & 0 \end{pmatrix} + \frac{1}{c} \begin{pmatrix} 0 & (A+B)G^\infty(z) \\ (A+B)G^\infty(z) & 0 \end{pmatrix} + o\left(\frac{1}{c^2}\right).$$

In general, the diagonal elements of $G(z, c)$ are even in $1/c$, the off-diagonal elements are odd. These analyticity properties of the resolvent are the basis from which the analyticity properties of eigenvalues and eigenfunctions follow in the usual way [6]. As has been remarked by Veselić, the fact that $G(z, \infty)$ is a pseudo-resolvent rather than a resolvent is thereby no obstacle. We only give the final result:

Theorem 4. *Let A, B, C satisfy (4) and (11). Let z be an eigenvalue of H^∞ in the complement of $\sigma(A^2)$ and m its (finite) multiplicity. Then z is the limit for $c \rightarrow \infty$ of eigenvalues $z_k(c)$ of $H(c)$ (with rest-energy subtracted) of total multiplicity m . The functions $z_k(c)$ are analytic in $(1/c)^2$ for $|c|$ sufficiently large. An orthonormal set of corresponding eigenvectors of $H(c)$ can be chosen such that each eigenvector is of the form $\begin{pmatrix} u_1(c) \\ u_2(c) \end{pmatrix}$ where $u_1(c)$ and $c^{-1}u_2(c)$ are analytic in $(1/c)^2$ and where $u_1(\infty)$ is an eigenvector of H^∞ with eigenvalue z .*

Remark. In general there will be other eigenvalues of $H(c)$ which for $c \rightarrow \infty$ will not converge to an eigenvalue of H^∞ in the complement of $\sigma(A^2)$. However, these eigenvalues will leave any compact not intersecting $\sigma(A^2)$, i.e. they will either join $\sigma(A^2)$ or disappear to $\pm\infty$. In fact it is equally possible to study the limit $c \rightarrow \infty$ of $-H(c) - \frac{1}{2}c^2$ which leads to a Pauli Hamiltonian $(A+B)^2 - C$.

4. Proof of Theorem 3

(8) \rightarrow (9):

We define $u_1 = u$ and $u_2 \in D(A)$ by

$$(A+B)u_1 - u_2 = 0. \tag{14}$$

Then (8) takes the form

$$(z - C)u_1 - (A + B)u_2 - v = 0, \quad (15)$$

and the combination $A(z - A^2)^{-1}(14) - (z - A^2)^{-1}(15)$ gives the upper component of (9)

$$u_1 = (z - A^2)^{-1}v + A(z - A^2)^{-1}Bu_1 + (z - A^2)^{-1}Cu_1 + (z - A^2)^{-1}Bu_2. \quad (16)$$

All terms in (16) are in $D(A)$. The combination $A(16) + (14)$ leads to the lower component of (9):

$$u_2 = A(z - A^2)^{-1}v + z(z - A^2)^{-1}Bu_1 + A(z - A^2)^{-1}Cu_1 + A(z - A^2)^{-1}Bu_2. \quad (17)$$

(9) → (8):

(9) is equivalent to the set (16) (17) with two important modifications. First, all operator products must be replaced by their bounded extensions to \mathcal{H} . To indicate this extension we write, for example, $[A(z - A^2)^{-1}B]$ for the extension of $A(z - A^2)^{-1}B$. Secondly, we start only with the information that u_1 and u_2 are in \mathcal{H} . The combination $A(z - A^2)^{-1}(16) + (z - A^2)^{-1}(17)$ gives

$$[(z - A^2)^{-1}(A + B)]u_1 = (z - A^2)^{-1}u_2, \quad (18)$$

where we have used identities like

$$A(z - A^2)^{-1}[(z - A^2)^{-1}C] = (z - A^2)^{-1}[A(z - A^2)^{-1}C],$$

which hold trivially on $D(A)$ and extend by continuity to all of \mathcal{H} . We now take the scalar product of (18) with an arbitrary $f \in \mathcal{H}$ and set $g = (\bar{z} - A^2)^{-1}f$. Using $[(z - A^2)^{-1}(A + B)]^* = (A + B)(\bar{z} - A^2)^{-1}$ we find

$$((A + B)g, u_1) = (g, u_2) \quad (19)$$

for all $g \in D(A^2)$. Since A is the closure of its restriction to $D(A^2)$ and since B is A -bounded, this extends by continuity to all $g \in D(A)$. Since $A + B$ is selfadjoint with domain $D(A)$, it follows that

$$u_1 \in D(A + B) \quad \text{and} \quad u_2 = (A + B)u_1. \quad (20)$$

Writing (20) in the form

$$u_1 = z(z - A^2)^{-1}u_1 + A(z - A^2)^{-1}Bu_1 - A(z - A^2)^{-1}u_2$$

and subtracting this from (16) we get

$$(z - A^2)^{-1}(z - C)u_1 = (z - A^2)^{-1}v + [(z - A^2)^{-1}(A + B)]u_2.$$

In the same way as before we conclude that

$$((A + B)g, u_2) = (g, (z - C)u_1 - v)$$

for all $g \in D(A)$. It follows that $u_2 \in D(A + B)$ and $(A + B)u_2 = (z - C)u_1 - v$. Combined with (20) this is the desired result:

$$u_1 \in D((A + B)^2), \quad u_2 = (A + B)u_1 \quad \text{and} \quad (z - H^\infty)u_1 = v.$$

References

1. See for instance Bjorken, J. D., Drell, S. D.: *Relativistic Quantum Mechanics*. New York: Mc Graw Hill Inc. 1964
2. Titchmarsh, E. C.: *Proc. Roy. Soc.* **266** A, 33 (1962)
3. Veselić, K.: *Commun. math. Phys.* **22**, 27 (1971)
4. Hunziker, W.: unpublished report, available from the Institute of theoretical physics of the ETH
5. Prosser, R. T.: *J. Math. Phys.* **4**, 1048 (1964)
6. Kato, T.: *Perturbation theory for linear operators*. Berlin-Heidelberg-New York: Springer 1966
7. Hunziker, W.: *Helv. Phys. Acta* **39**, 452 (1966)

Communicated by K. Hepp

W. Hunziker
Institut für Theoretische Physik der E. T. H.
CH-8049 Zürich, Switzerland