

HARMONIC MAPS OF COMPLETE RIEMANNIAN MANIFOLDS

SEOUNG DAL JUNG

1. Introduction

The theory of harmonic mappings of a Riemannian manifold into another has been initiated by J. Eells and J. H. Sampson([2]) and studied by many authors. In particular, R. M. Schoen and S. T. Yau([3]) proved the following theorem:

Theorem A. *Let M be a complete noncompact Riemannian manifold with nonnegative Ricci curvature and let N be a compact Riemannian manifold of nonpositive sectional curvature. Then every harmonic map of finite energy from M to N is constant.*

In this paper, we extend Theorem A under weaker assumptions by using Kato's inequality([1]) and characterize a harmonic map on complete Riemannian manifolds.

The author would like to thank the referee for his helpful and kind suggestions.

2. Preliminaries

Let $\pi : E \rightarrow M$ be a Riemannian vector bundle over an m -dimensional manifold M , i.e., E is a vector bundle over M equipped with a C^∞ -assignment of an inner product $\langle \cdot, \cdot \rangle$ to each fiber E_x of E over $x \in M$. Assume that a metric connection D is given on E , i.e., $D : A^p(E) \rightarrow A^{p+1}(E)$ is an \mathbb{R} -linear map such that if $f \in A^0$, $D(fs) = fDs + sdf$ and

$$(2.1) \quad d \langle s, t \rangle = \langle Ds, t \rangle + \langle s, Dt \rangle$$

1991 *Mathematics Subject Classification.* 58E20.

Key words and phrases. Harmonic maps, Kato's inequality, Tension field.

This work was partially supported by Development Fund of Cheju National University, 1997

for any $s, t \in A^p(E)$, where $A^p(E)$ is the set of all E -valued p -forms on M . Equivalently, for any $X \in TM$,

$$(2.2) \quad X \langle s, t \rangle = \langle D_X s, t \rangle + \langle s, D_X t \rangle.$$

The Laplacian for $A^*(E)$ is by definition the operator

$$(2.3) \quad \Delta = DD^* + D^*D,$$

where D^* is the formal adjoint of D . Let $\{V_1, \dots, V_m\}$ and $\{\omega^1, \dots, \omega^m\}$ be a locally defined frame field and its dual coframe field respectively. Then on $A^*(E)$, we have

$$(2.4) \quad D = \sum_{i=1}^m \omega^i \wedge D_{V_i}, \quad D^* = - \sum_{j=1}^m i(V_j) D_{V_j},$$

where $i(X)$ denotes the interior product operator with respect to X . Then on $A^*(E)$, we have the Weitzenböck formula

$$(2.5) \quad \Delta = - \sum_i D_{V_i}^2 + \sum_{k,\ell} \omega^k \wedge i(V_\ell) R_{V_k V_\ell},$$

where $D_{XY}^2 = D_X D_Y - D_{\nabla_X Y}$ and $R_{XY} = -[D_X, D_Y] + D_{[X,Y]}$ is the curvature tensor on $A^*(E)$. From this equation, we have

$$(2.6) \quad -\Delta^M |\Phi|^2 = 2 \sum_i |D_{V_i} \Phi|^2 + 2 \langle \Phi, \sum_i D_{V_i}^2 \Phi \rangle,$$

where $\langle \cdot, \cdot \rangle$ refers to the pointwise inner product, $|\Phi|^2 = \langle \Phi, \Phi \rangle$, and $\Delta^M \equiv d\delta + \delta d$ is the ordinary Hodge Laplacian. From (2.5) and (2.6), for any harmonic section $\Phi \in A^*(E)$, we have

$$(2.7) \quad -\Delta^M |\Phi|^2 = 2 \sum_i |D_{V_i} \Phi|^2 + 2 \langle \Phi, \sum_{k,\ell} \omega^k \wedge i(V_\ell) R_{V_k V_\ell} \Phi \rangle.$$

In particular, let $\Phi = \sum_a e_a \phi^a \in A^1(E)$, where $\{e_1, \dots, e_r\}$ is a local frame field of E and $\{\phi^a\}$ are 1-forms on M . Then we obtain

$$\begin{aligned} \langle \sum_{k,\ell} \omega^k \wedge i(V_\ell) R_{V_k V_\ell} \Phi, \Phi \rangle &= \sum_{a,b,k,\ell} \langle R_{V_k V_\ell}^E e_a, e_b \rangle \langle \omega^k \wedge i(V_\ell) \phi^a, \phi^b \rangle \\ &+ \sum_{a,k,\ell} \langle \omega^k \wedge i(V_\ell) R_{V_k V_\ell}^M \phi^a, \phi^a \rangle, \end{aligned}$$

where R^E and R^M are the curvature tensors on $\Gamma(E)$ and M , respectively. The second summand is equal to $\sum_a Ric^M(\phi_a, \phi_a)$, where Ric^M denotes the Ricci curvature tensor of M and ϕ_a stands for the vector field dual to ϕ^a . For the first summand, if we set $\phi^a = \sum \alpha_i^a \omega^i$, then $\langle \omega^i \wedge i(V_j)\phi^a, \phi^b \rangle = \alpha_j^a \alpha_i^b$, so that

$$\begin{aligned} \sum_{a,b,k,\ell} \langle R_{V_k V_\ell}^E e_a, e_b \rangle \langle \omega^k \wedge i(V_\ell)\phi^a, \phi^b \rangle &= \sum_{a,b,k,\ell} \langle R_{V_k V_\ell}^E e_a \alpha_\ell^a, e_b \alpha_k^b \rangle \\ &= \sum_{k,\ell} \langle R_{V_k V_\ell}^E \Phi(V_\ell), \Phi(V_k) \rangle. \end{aligned}$$

Hence for a harmonic E -valued 1-form Φ , we have

$$(2.8) \quad \begin{aligned} -\Delta^M |\Phi|^2 &= 2 \sum_i |D_{V_i} \Phi|^2 + 2 \sum_a Ric^M(\phi_a, \phi_a) \\ &\quad - 2 \sum_{i,j} \langle R_{V_i V_j}^E \Phi(V_i), \Phi(V_j) \rangle. \end{aligned}$$

3. Harmonic maps

Let M and N be Riemannian manifolds with Riemannian metrics g and h respectively, and let their Levi-Civita connections be ∇^M and ∇^N respectively. Let $f : M \rightarrow N$ be a C^∞ -map, and let $E \equiv f^*TN$ be the induced bundle over M . Then E has a naturally induced metric connection which we denote by $D \equiv f^*\nabla^N$. Also, the differential df of f gives naturally a cross section of the vector bundle $\text{Hom}(TM, E)$ over M . Since $\text{Hom}(TM, E)$ is canonically identified with $E \otimes T^*M$ (T^*M = the cotangent bundle of M), we see that df may be regarded as an E -valued 1-form, i.e., $df \in \Gamma(E \otimes T^*M) = A^1(E)$. Then $E \otimes T^*M$ has a naturally induced metric connection which we denote also by D . It is well known that $D(df) = 0$ ([2]). Hence we say that the map f is *harmonic* if $D^*(df) = 0$. Also, f is said to be *totally geodesic* if for any $X \in TM$, $D_X df = 0$. It follows that totally geodesic maps are harmonic but the converse is not true in general. Let R^E be the curvature tensor of D . Then R^E is related to the curvature tensor R^N of ∇^N in the following way: let $X, Y \in T_x M$ and $s \in \Gamma(E)$, then

$$(3.1) \quad R_{XY}^E s = R_{df_x(X)df_x(Y)}^N s.$$

When a function ρ , local or global, is given on N , we shall identify it throughout this paper with the function $\rho \circ f$ induced on M .

Let $\{\bar{V}_a\}$ and $\{V_i\}$ be local frame fields on N and M respectively, and let $\{\bar{\omega}^a\}$ and $\{\omega^i\}$ be the their dual coframe fields respectively. Let $\phi^a \equiv f^*\bar{\omega}^a$. Then we can write df as $df = \sum_a \bar{V}_a \otimes \phi^a$. Hence if we set $\Phi = df$ in (2.8), then from (3.1) we obtain

$$(3.2) \quad \begin{aligned} -\Delta^M |df|^2 = & 2 \sum_i |D_{V_i} df|^2 + 2 \sum_a Ric^M(\phi_a, \phi_a) \\ & - 2 \sum_{i,j} \langle R_{df(V_i)df(V_j)}^N df(V_i), df(V_j) \rangle, \end{aligned}$$

where ϕ_a is the vector field dual to ϕ^a .

4. Main Theorems

Let $\rho(x)$ denote the least eigenvalue of $\mathcal{R}_x \equiv \sum_{k,\ell} \omega^k \wedge i(V_\ell) R_{V_k V_\ell}$ in E_x , where $\{V_j\}$ is a local frame field and $\{\omega^j\}$ is the dual coframe field of $\{V_j\}$. To prove our theorem, we prepare the following Kato's inequality:

Lemma 4.1([1]). *If $s \in \Gamma(E)$ satisfies $\Delta s = 0$, then*

$$\Delta^M |s| \leq -\rho |s|.$$

Let λ_0 is the infimum of the spectrum of the Laplacian Δ^M on L^2 -functions on M . The proof of the following Theorem 4.2 is based on a method of P. Bérard ([1]).

Theorem 4.2. *Let M be a complete noncompact Riemannian manifold. If $\rho \geq -\lambda_0$ at all $x \in M$ and $\rho > -\lambda_0$ at some point x_0 , then every harmonic map $f : M \rightarrow N$ of finite energy, i.e., $df \in L^2(M)$, is constant.*

proof. Since $\Delta df = 0$, by Lemma 4.1, we have

$$\Delta^M |df| \leq -\rho |df|.$$

By the assumption, we obtain

$$(4.1) \quad \Delta^M |df| \leq -\rho |df| \leq \lambda_0 |df|.$$

Since M is complete, one can construct function ω_ℓ such that $\omega_\ell \in C_0^\infty(M)$ and $\omega_\ell \equiv 1$ on $B(x_0, \ell)$, $\text{supp} \omega_\ell \subset B(x_0, 2\ell)$ and $|d\omega_\ell| \leq C/\ell$ for some

constant C , where $\ell \in \mathbb{R}_+$, $x_0 \in M$ and $B(x_0, \ell)$ is the Riemannian open ball with radius ℓ and center x_0 . Put $\varphi \equiv |df|$. Then from (4.1), we have

$$(4.2) \quad \int_M \langle d\omega_\ell^2 \varphi, d\varphi \rangle \leq - \int_M \omega_\ell^2 \rho \varphi^2 \leq \lambda_0 \int_M \omega_\ell^2 \varphi^2.$$

Since $d\omega_\ell^2 \varphi = 2\omega_\ell d\omega_\ell \varphi + \omega_\ell^2 d\varphi$, the left hand side of (4.2) can be written as

$$(4.3) \quad \int_M 2 \langle d\omega_\ell \varphi, \omega_\ell d\varphi \rangle + \int_M \omega_\ell^2 |d\varphi|^2 = \int_M |d(\omega_\ell \varphi)|^2 - \int_M \varphi^2 |d\omega_\ell|^2.$$

From (4.2) and (4.3), we then obtain the inequality

$$(4.4) \quad \int_M |d(\omega_\ell \varphi)|^2 \leq \int_M \varphi^2 |d\omega_\ell|^2 - \int_M \rho \omega_\ell^2 \varphi^2 \leq \int_M \varphi^2 |d\omega_\ell|^2 + \lambda_0 \int_M (\omega_\ell \varphi)^2.$$

Since λ_0 is the infimum of the spectrum of the Laplacian Δ^M acting on functions, we have

$$(4.5) \quad \int_M |d(\omega_\ell \varphi)|^2 \geq \lambda_0 \int_M (\omega_\ell \varphi)^2.$$

From (4.4) and (4.5), we have

$$\lambda_0 \int_M (\omega_\ell \varphi)^2 \leq \int_M \varphi^2 |d\omega_\ell|^2 - \int_M \rho \omega_\ell^2 \varphi^2 \leq \int_M \varphi^2 |d\omega_\ell|^2 + \lambda_0 \int_M (\omega_\ell \varphi)^2.$$

Now, letting $\ell \rightarrow \infty$, we have

$$\lambda_0 \int_M \varphi^2 \leq - \int_M \rho \varphi^2 \leq \lambda_0 \int_M \varphi^2.$$

This implies that $\varphi = |df| = 0$ under our assumptions. Hence f is constant. \square

Corollary 4.3. *Let M be a complete Riemannian manifold and let N be a Riemannian manifold of nonpositive sectional curvature. If $\text{Ric}^M \geq -\lambda_0$ at all $x \in M$ and $\text{Ric}^M > -\lambda_0$ at some point x_0 , then every harmonic map $f : M \rightarrow N$ of finite energy, i.e., $df \in L^2(M)$, is constant.*

Proof. First, we recall the first Kato's inequality ([1]): for any $s \in \Gamma(E)$, $|d|s|| \leq |Ds|$. It follows from the first Kato's inequality that if $\Delta s = 0$, then

$|s|\Delta^M|s| \leq -\langle \mathcal{R}(s), s \rangle$ (cf. [1], p.263). Since $\Delta df = 0$ for a harmonic map $f : M \rightarrow N$, we get from the first Kato's inequality and (3.2)

$$\begin{aligned} |df|\Delta^M|df| &\leq -\langle \mathcal{R}(df), df \rangle \\ &= -\sum_a Ric^M(f^*\bar{w}^a, f^*\bar{w}^a) + \sum_{i,j} \langle R_{df(V_i)df(V_j)}^N df(V_i), df(V_j) \rangle. \end{aligned}$$

Note that if N has nonpositive sectional curvature, then we get

$$\sum_{i,j} \langle R_{df(V_i)df(V_j)}^N df(V_i), df(V_j) \rangle \leq 0.$$

Hence, if N has nonpositive sectional curvature, then we have

$$|df|\Delta^M|df| \leq -\sum_a Ric^M(f^*\bar{w}^a, f^*\bar{w}^a).$$

If we assume $Ric^M \geq -\lambda_0$, then we have

$$|df|\Delta^M|df| \leq \lambda_0|df|^2.$$

Hence we have $\Delta^M|df| \leq \lambda_0|df|$. Then the rest of the proof goes as in the proof of Theorem 4.2. \square

Now we are going to prove a characterization of harmonic maps on complete Riemannian manifolds. Take coordinate neighborhoods $\{U; x^j\}$ of M and $\{V; y^a\}$ of N in such a way that $f(U) \subset V$, where $(x^j) = (x^1, \dots, x^m)$ and $(y^a) = (y^1, \dots, y^n)$ are local coordinates of M and N respectively. The indices i, j, k run over the range $\{1, \dots, m\}$ and the indices a, b, c the range $\{1, \dots, n\}$. The summation convention will be used with respect to these two systems of indices. Suppose that $f : M \rightarrow N$ is represented by equation $y^a = y^a(x^1, \dots, x^m)$ with respect to $\{U; x^j\}$ and $\{V; y^a\}$. We put $A_i^a = \partial_i y^a(x^1, \dots, x^m)$, where $\partial_i = \partial/\partial x^i$. Then the differential df of the mapping f is represented by the matrix (A_i^a) with respect to the local coordinates (x^j) and (y^a) of M and N . Let $X = X^j \partial_j$ and $Y = Y^j \partial_j$ are vector fields on M . If we put in U

$$\begin{aligned} A_{j;i}^a &= \nabla_j A_i^a, \\ \nabla_j A_i^a &= \partial_j A_i^a + \Gamma_{bc}^a A_j^b A_i^c - \Gamma_{ji}^k A_k^a, \end{aligned}$$

where Γ_{bc}^a and Γ_{ji}^k are Christoffel symbols with respect to ∇^N and ∇^M respectively, then $(A_{ji}^a X^j Y^i) \partial_a$ is the local expression of a vector field B defined along $f(M)$ and $A_{ji}^a = A_{ij}^a$. We now put

$$A^a = g^{ij} A_{ij}^a,$$

where $(g^{ij}) = (g_{ij})^{-1}$. Then the vector field T with component A^a defined along $f(M)$ is called the *tension field* of $f : M \rightarrow N$. It is well known that $f : M \rightarrow N$ is harmonic if and only if $T = 0$ ([2]). If we put $\eta = (A_i^c A^b g_{bc}) dx^i$, then η is a 1-form in M . Hence $\ast(\omega_\ell^2 \eta)$ is an $(m-1)$ -form with compact support in $B(2\ell)$, where \ast is the Hodge star operator. Then by Stokes' theorem, we have

$$\int_M d(\ast(\omega_\ell^2 \eta)) = 0.$$

Since $d(\ast(\omega_\ell^2 \eta)) = -\ast \delta(\omega_\ell^2 \eta)$, we have

$$\int_M \ast \delta(\omega_\ell^2 \eta) = \int_{B(2\ell)} \ast \delta(\omega_\ell^2 \eta) = 0.$$

Moreover, we have

$$\delta(\omega_\ell^2 \eta) = \omega_\ell^2 \delta \eta - \ast(2\omega_\ell d\omega_\ell \wedge \ast \eta)$$

on M . Also, for a 1-form η , $\delta \eta = -\nabla^i \eta_i$. Hence we have

$$\begin{aligned} \delta(\omega_\ell^2 \eta) &= \omega_\ell^2 \{-\nabla^i (A_i^c A^b g_{bc})\} - \ast(2\omega_\ell d\omega_\ell \wedge \ast \eta) \\ &= -\omega_\ell^2 \{(\nabla^i A_i^c) A^b g_{bc} + A_i^c \nabla^i A^b g_{bc}\} - \ast(2\omega_\ell d\omega_\ell \wedge \ast \eta) \\ &= -\omega_\ell^2 \{A^c A_b g_{bc}\} - \omega_\ell^2 A_i^c \nabla^i A^b g_{bc} - \ast(2\omega_\ell d\omega_\ell \wedge \ast \eta) \\ &= -|\omega_\ell T|^2 - \langle \omega_\ell df, \omega_\ell \nabla T \rangle - \ast(2\omega_\ell d\omega_\ell \wedge \ast \eta). \end{aligned}$$

Since $\ast(d\omega_\ell \wedge \ast \eta) = (d\omega_\ell)_i \eta^i = \langle d\omega_\ell, \eta \rangle$, we have

$$\delta(\omega_\ell^2 \eta) = -|\omega_\ell T|^2 - \langle \omega_\ell df, \omega_\ell \nabla T \rangle - 2 \langle d\omega_\ell, \omega_\ell \eta \rangle.$$

By integration, we have

$$0 = \|\omega_\ell T\|_{B(2\ell)}^2 + \langle \langle \omega_\ell df, \omega_\ell \nabla T \rangle \rangle_{B(2\ell)} + 2 \langle \langle d\omega_\ell, \omega_\ell \eta \rangle \rangle_{B(2\ell)}.$$

By Schwartz's inequality and $|d\omega_\ell| \leq C/\ell$, we get

$$0 \geq \|\omega_\ell T\|_{B(2\ell)}^2 - \|\omega_\ell df\| \|\omega_\ell \nabla T\|_{B(2\ell)} - \frac{C}{\ell} \|\omega_\ell \eta\|_{B(2\ell)}.$$

For any finite energy map f , i.e., $|df| \in L^2(M)$, letting $\ell \rightarrow \infty$, we have

$$0 \geq \|T\|^2 - \|df\| \|\nabla T\|.$$

From this inequality, we have the following:

Theorem 4.4. *Let M be a complete Riemannian manifold. If a mapping $f : M \rightarrow N$ has a finite energy and $\nabla T = 0$, then f is harmonic.*

REFERENCES

- [1] P. Bérard, *A note on Bochner type theorems for complete manifolds*, Manuscripta Math. **69** (1990), 261-266.
- [2] J. Eells and J. H. Sampson, *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. **86** (1964), 109-160.
- [3] R. M. Schoen and S. T. Yau, *Harmonic maps and the topology of stable hypersurfaces and manifolds of nonnegative Ricci curvature*, Comm. Math. Helv. **51** (1976), 333-341.
- [4] H. H. Wu, *The Bochner technique in differential geometry*, Mathematical reports **3** (1988), 289-538.

DEPARTMENT OF MATHEMATICS, CHEJU NATIONAL UNIVERSITY, CHEJU 690-756,
KOREA

Received February 21, 1997

Revised May 13, 1997