

COMMUTING FAMILIES IN HECKE AND TEMPERLEY-LIEB ALGEBRAS

TOM HALVERSON, MANUELA MAZZOCCO AND ARUN RAM

Abstract. We define analogs of the Jucys-Murphy elements for the affine Temperley-Lieb algebra and give their explicit expansion in terms of the basis of planar Brauer diagrams. These Jucys-Murphy elements are a family of commuting elements in the affine Temperley-Lieb algebra, and we compute their eigenvalues on the generic irreducible representations. We show that they come from Jucys-Murphy elements in the affine Hecke algebra of type A, which in turn come from the Casimir element of the quantum group $U_h \mathfrak{gl}_n$. We also give the explicit specializations of these results to the finite Temperley-Lieb algebra.

§1. Introduction

The “Jucys-Murphy elements” are a family of commuting elements in the group algebra of the symmetric group. In characteristic 0, these elements have enough distinct eigenvalues to give a full analysis of the representation theory of the symmetric group [OV]. Even in positive characteristic these elements are powerful tools [K]. Similar elements are used in the Hecke algebras of type A and, in a strong sense, it is these elements that control the beautiful connections between the modular representation theory of Hecke algebras of type A and the Fock space representations of the affine quantum group (see [Ar] and [Gr]).

Since the Temperley-Lieb algebra is a quotient of the Hecke algebra of type A it inherits a commuting family of elements from the Hecke algebra (these elements can be viewed as the image, under a surjective homomorphism, of the commutative subalgebra in the Bernstein presentation of the affine Hecke algebra). In order to use these elements for modular representation theory of the Temperley-Lieb algebra it is important to have good

Received October 5, 2007.

Revised November 18, 2008.

Accepted February 4, 2009.

2000 Mathematics Subject Classification: Primary 20G05; Secondary 16G99, 81R50, 82B20.

control of their expansion in terms of the standard basis of noncrossing diagrams. In this paper we study this question, in the more general setting of the affine Temperley-Lieb algebras. Specifically, we analyze a convenient choice of a commuting family of elements in the affine Temperley-Lieb algebra. Our main result, Theorem 2.9, is an explicit expansion of these elements in the standard basis. The fact that, in the Temperley-Lieb algebra, these elements have integral coefficients is made explicit in Remark 2.10. The import of this result is that this commuting family can be used to attack questions in modular representation theory.

In Section 3 we review the Schur-Weyl duality setup of Orellana and Ram [OR] which (following the ideas in [Re]) explains how commuting families in centralizer algebras arise naturally from Casimir elements. We explain, in detail, the cases that lead to commuting families in the affine Hecke algebras of type A and the affine Temperley-Lieb algebra. One new consequence of our analysis is an explanation of the “special” relation that is used in one of the Temperley-Lieb algebras of Graham and Lehrer [GL2]. In our context, this relation appears naturally from the Schur-Weyl duality (see Proposition 3.2). Using the knowledge of eigenvalues of Casimir elements we compute the eigenvalues of the commuting families in the affine Hecke algebra and in the affine Temperley-Lieb algebra in the generic irreducible representations (analogues of the Specht, or Weyl, modules).

The representation theory of Temperley-Lieb algebras and affine Temperley-Lieb algebras is ‘well understood’. In particular, in the paper [GL1], the cell modules and the irreducible modules of the affine Temperley-Lieb algebra are analyzed in detail and the eigenvalues of a commuting family of ‘Murphy elements’ on these modules are determined. They show ([GL1, Thm. (9.6)]) that the representations of the affine Hecke algebra which appear as affine Temperley-Lieb algebra representations are those representations that correspond, under the Deligne-Langlands correspondence [KL], to unipotent elements in GL_n with at most two Jordan blocks. The commuting family of elements we use in this paper does not differ significantly from those used in [GL1]. Our results provide an expansion of these elements in terms of the basis of noncrossing diagrams and place the representations studied in [GL1] into a Schur-Weyl duality context.

The research of M. Mazzocco was partially supported by a Mark Mensink Honors Research Award at the University of Wisconsin, Madison. She was an undergraduate researcher participating in research and teaching initiatives partly supported by the National Science Foundation under

grant DMS-0353038. The research of A. Ram was also partially supported by this award. A significant portion of this research was done during a residency of A. Ram at the Max-Planck-Institut für Mathematik (MPI) in Bonn. He thanks the MPI for support, hospitality, and a wonderful working environment. The research of T. Halverson was partially supported by the National Science Foundation under grant DMS-0100975.

§2. Affine braid groups, Hecke and Temperley-Lieb algebras

2.1. The affine braid group $\tilde{\mathcal{B}}_k$

The *affine braid group* is the group $\tilde{\mathcal{B}}_k$ of affine braids with k strands (braids with a flagpole). The group $\tilde{\mathcal{B}}_k$ is presented by generators T_1, T_2, \dots, T_{k-1} and X^{ε_1} ,

$$(2.1) \quad \begin{aligned} T_i &= \text{flagpole} \parallel \text{strand } i \text{ crosses over strand } i+1 \parallel \text{strand } i+1 \parallel \text{strand } i \parallel \text{strand } i+1 \parallel \text{strand } i \parallel \text{strand } i+1 \text{ and} \\ X^{\varepsilon_1} &= \text{flagpole} \text{ with a loop on the left side} \parallel \text{strand } 1 \parallel \text{strand } 2 \parallel \text{strand } 3 \parallel \text{strand } 4 \parallel \text{strand } 5 \parallel \text{strand } 6 \parallel \text{strand } 7 \end{aligned}$$

with relations

$$(2.2) \quad \begin{aligned} X^{\varepsilon_1} T_1 X^{\varepsilon_1} T_1 &= T_1 X^{\varepsilon_1} T_1 X^{\varepsilon_1} \\ X^{\varepsilon_1} T_i &= T_i X^{\varepsilon_1}, & \text{for } i > 1, \\ T_i T_j &= T_j T_i, & \text{if } |i - j| > 1, \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}, & \text{if } 1 \leq i \leq k - 2. \end{aligned}$$

For $1 \leq i \leq k$ define

$$(2.3) \quad \begin{aligned} X^{\varepsilon_i} &= T_{i-1} T_{i-2} \cdots T_2 T_1 X^{\varepsilon_1} T_1 T_2 \cdots T_{i-1} \\ &= \text{flagpole} \text{ with a loop on the left side} \parallel \text{strand } 1 \parallel \text{strand } 2 \parallel \text{strand } 3 \parallel \text{strand } 4 \parallel \text{strand } 5 \parallel \text{strand } 6 \parallel \text{strand } 7 \text{ with strand } i \text{ crossing over strand } i+1. \end{aligned}$$

By drawing pictures of the corresponding affine braids it is easy to check that


$$(2.4) \quad X^{\varepsilon_i} X^{\varepsilon_j} = X^{\varepsilon_j} X^{\varepsilon_i}, \quad \text{for } 1 \leq i, j \leq k,$$

so that the elements $X^{\varepsilon_1}, \dots, X^{\varepsilon_k}$ are a commuting family for $\tilde{\mathcal{B}}_k$. Thus $X = \langle X^{\varepsilon_i} \mid 1 \leq i \leq k \rangle$ is an abelian subgroup of $\tilde{\mathcal{B}}_k$. The free abelian group generated by $\varepsilon_1, \dots, \varepsilon_k$ is \mathbb{Z}^k and

$$(2.5) \quad X = \{X^\lambda \mid \lambda \in \mathbb{Z}^k\} \quad \text{where} \quad X^\lambda = (X^{\varepsilon_1})^{\lambda_1} (X^{\varepsilon_2})^{\lambda_2} \dots (X^{\varepsilon_k})^{\lambda_k},$$

for $\lambda = \lambda_1 \varepsilon_1 + \dots + \lambda_k \varepsilon_k$ in \mathbb{Z}^k .

Remark 2.1. An alternate presentation of $\tilde{\mathcal{B}}_k$ can be given using the generators T_0, T_1, \dots, T_{k-1} and τ where



Remark 2.2. The affine braid group $\tilde{\mathcal{B}}_k$ is the affine braid group of type GL_k . The affine braid groups of type SL_k and PGL_k are the subgroup

$$\tilde{\mathcal{B}}_Q = \langle T_0, T_1, \dots, T_{k-1} \rangle \quad \text{and the quotient} \quad \tilde{\mathcal{B}}_P = \frac{\tilde{\mathcal{B}}_k}{\langle \tau^k \rangle}, \quad \text{respectively.}$$

Then $\tau^k = X^{-\varepsilon_1} X^{-\varepsilon_2} \dots X^{-\varepsilon_k}$ is a central element of $\tilde{\mathcal{B}}_k$, $\tau T_i \tau^{-1} = T_{i+1}$ (where the indices are taken mod k), and

$$Z(\tilde{\mathcal{B}}_k) = \langle \tau^k \rangle, \quad \tilde{\mathcal{B}}_k = \langle \tau \rangle \rtimes \tilde{\mathcal{B}}_Q, \quad \tilde{\mathcal{B}}_P = \langle \bar{\tau} \rangle \rtimes \tilde{\mathcal{B}}_Q.$$

In $\tilde{\mathcal{B}}_k$ we have $\langle \tau \rangle \cong \mathbb{Z}$, and $\bar{\tau} \in \tilde{\mathcal{B}}_P$ is defined to be the image of τ under the homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}/k\mathbb{Z}$ so that $\langle \bar{\tau} \rangle \cong \mathbb{Z}/k\mathbb{Z}$.

2.2. The Temperley-Lieb algebra $TL_k(n)$


A Temperley-Lieb diagram on k dots is a graph with k dots in the top row, k dots in the bottom row, and k edges pairing the dots such that the graph is planar (without edge crossings). For example,



are Temperley-Lieb diagrams on 7 dots. The composition $d_1 \circ d_2$ of two diagrams $d_1, d_2 \in T_k$ is the diagram obtained by placing d_1 above d_2 and identifying the bottom vertices of d_1 with the top dots of d_2 removing any connected components that live entirely in the middle row. If T_k is the set of Temperley-Lieb diagrams on k dots then the Temperley-Lieb algebra $TL_k(n)$ is the associative algebra with basis T_k ,

$$TL_k(n) = \text{span}\{d \in T_k\} \text{ with multiplication defined by } d_1 d_2 = n^\ell (d_1 \circ d_2),$$

where ℓ is the number of blocks removed from the middle row when constructing the composition $d_1 \circ d_2$ and n is a fixed element of the base ring. For example, using the diagrams d_1 and d_2 above, we have



The algebra $TL_k(n)$ is presented by generators

$$(2.6) \quad e_i = \begin{array}{c} \bullet \quad \dots \quad \bullet \quad \overset{i}{\curvearrowright} \quad \overset{i+1}{\curvearrowleft} \quad \bullet \quad \dots \quad \bullet \\ | \quad \dots \quad | \quad \bullet \quad \bullet \quad | \quad \dots \quad | \\ \bullet \quad \dots \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \dots \quad \bullet \end{array}, \quad 1 \leq i \leq k - 1,$$

and relations

$$(2.7) \quad e_i^2 = ne_i, \quad e_i e_{i\pm 1} e_i = e_i, \quad \text{and} \quad e_i e_j = e_j e_i, \quad \text{if } |i - j| > 1$$

(see [GHJ, Lemma 2.8.4]).

Remark 2.3. In the definition of the Temperley-Lieb algebra, and for other algebras defined in this paper, the base ring could be any one of several useful rings (e.g. \mathbb{C} , $\mathbb{C}(q)$, $\mathbb{C}[[h]]$, $\mathbb{Z}[q, q^{-1}]$, $\mathbb{Z}[n]$ or localizations of these at special primes). The most useful approach is to view the results of computations as valid over any ring R with $n, q, h \in R$ such that the formulas make sense.

2.3. The surjection $\tilde{H}_k(q) \mapsto TL_k(n)$

The affine Hecke algebra \tilde{H}_k is the quotient of the group algebra of the affine braid group $\mathbb{C}\tilde{\mathcal{B}}_k$ by the relations

$$(2.8) \quad T_i^2 = (q - q^{-1})T_i + 1, \quad \text{so that } \mathbb{C}\tilde{\mathcal{B}}_k \longrightarrow \tilde{H}_k$$

is a surjective homomorphism (q is a fixed element of the base ring). The affine Hecke algebra \tilde{H}_k is the affine Hecke algebra of type GL_k . The *affine Hecke algebras of types SL_k and PGL_k* are, respectively, the quotients \tilde{H}_Q and \tilde{H}_P of the group algebras of $\tilde{\mathcal{B}}_Q$ and $\tilde{\mathcal{B}}_P$ (see Remark 2.2) by the relations (2.8).

The *Iwahori-Hecke algebra* is the subalgebra H_k of \tilde{H}_k generated by T_1, \dots, T_{k-1} . In the Iwahori-Hecke algebra H_k , define

$$(2.9) \quad e_i = q - T_i, \quad \text{for } i = 1, 2, \dots, k-1.$$

Direct calculations show that $e_i^2 = (q + q^{-1})e_i$ and that $e_1 e_2 e_1 = e_1$ and $e_2 e_1 e_2 = e_2$ if and only if

$$(2.10) \quad q^3 - q^2 T_1 - q^2 T_2 + q T_1 T_2 + q T_2 T_1 - T_1 T_2 T_1 = 0.$$

Thus, setting $n = [2] = q + q^{-1}$, there are surjective algebra homomorphisms given by

$$(2.11) \quad \begin{array}{ccccc} \psi : \tilde{H}_k(q) & \longrightarrow & H_k(q) & \longrightarrow & TL_k(n) \\ X^{\varepsilon_1} & \longmapsto & 1 & \longmapsto & 1 \\ T_i & \longmapsto & T_i & \longmapsto & q - e_i. \end{array}$$

The kernel of ψ is generated by the element on the left hand side of equation (2.10). In the notation of Theorem 4.1, the representations of H_k correspond to the case when $\mu = \emptyset$. Writing $\tilde{H}_k^{\lambda/\emptyset}$ as \tilde{H}_k^λ , the element from (2.10) acts as 0 on the irreducible Iwahori-Hecke algebra modules $\tilde{H}_3^{\mathbf{00}}$ and $\tilde{H}_3^{\mathbf{01}}$, and (up to a scalar multiple) it is a projection onto $\tilde{H}_3^{\mathbf{10}}$.

Remark 2.4. There is an alternative surjective homomorphism that instead sends $T_i \mapsto e_i - q^{-1}$. This alternative surjection has kernel generated by

$$q^{-3} + q^{-2} T_1 + q^{-2} T_2 + q^{-1} T_1 T_2 + q^{-1} T_2 T_1 + T_2 T_1 T_2.$$

This element is 0 on $\tilde{H}_3^{\mathbf{10}}$ and $\tilde{H}_3^{\mathbf{01}}$, and (up to a scalar multiple) it is a projection onto $\tilde{H}_3^{\mathbf{00}}$.

Remark 2.5. A priori, there are two different kinds of integrality for the Temperley-Lieb algebra: coefficients in $\mathbb{Z}[n]$ or coefficients in $\mathbb{Z}[q, q^{-1}]$ (in terms of the basis of Temperley-Lieb diagrams). The relation between these is as follows. If

$$[2] = q + q^{-1} = n \quad \text{then} \quad q = \frac{1}{2}(n + \sqrt{n^2 - 4}), \quad q^{-1} = \frac{1}{2}(n - \sqrt{n^2 - 4}),$$

since $q^2 - nq + 1 = 0$. Then

$$[k] = \frac{q^k - q^{-k}}{q - q^{-1}} = \frac{1}{2^{k-1}} \sum_{m=1}^{(k+1)/2} \binom{k}{2m-1} n^{k-2m+1} (n^2 - 4)^{m-1}$$


so that $[k]$ is a polynomial in n . The polynomials

$$n^k = (q + q^{-1})^k \quad \text{and} \quad \{k\} = q^k + q^{-k} \quad \text{and} \quad [k] = \frac{q^k - q^{-k}}{q - q^{-1}},$$

all form bases of the ring $\mathbb{C}[(q + q^{-1})]$. The transition matrix B between the $[k]$ and the $\{k\}$ is triangular (with 1s on the diagonal) and the transition matrix C between the n^k and the $\{k\}$ is also triangular (the non zero entries are binomial coefficients). Hence, the transition matrix BC^{-1} between $[k]$ and n^k has integer entries and so $[k]$ is, in fact, a polynomial in n with integer coefficients.

2.4. Affine Temperley-Lieb algebras

The *affine Temperley-Lieb algebra* T_k^a is the diagram algebra generated by



The generators of T_k^a satisfy $e_i^2 = ne_i$, $e_i e_{i\pm 1} e_i = e_i$, $\tau e_i \tau^{-1} = e_{i+1}$ (where the indices are taken mod k) and

(2.12)

$$\tau^2 e_{k-1} =$$
 $= e_1 e_2 \cdots e_{k-1}$

(see [GL2, 4.15(iv)]). In T_k^a , we let $X^{\varepsilon_1} = T_1^{-1} T_2^{-1} \cdots T_{k-1}^{-1} \tau^{-1}$ (see Remark 2.1).

Graham and Lehrer [GL2, §4.3] define four slightly different affine Temperley-Lieb algebras, the diagram algebra T_k^a and the algebras defined as follows:

$$\begin{aligned} \text{Type } GL_k: & \quad \widehat{TL}_k^a \text{ is } \tilde{H}_k \text{ with the relation (2.10),} \\ \text{Type } SL_k: & \quad TL_k^a \text{ is } \tilde{H}_Q \text{ with the relation (2.10),} \\ \text{Type } PGL_k: & \quad \widetilde{TL}_k^a \text{ is } \tilde{H}_P \text{ with the relation (2.10).} \end{aligned}$$

For each invertible element α in the base ring there is a surjective homomorphism

$$(2.13) \quad \begin{array}{ccccc} \tilde{H}_k & \longrightarrow & \widehat{TL}_k^a & \longrightarrow & T_k^a \\ \tau & \longmapsto & \tau & \longmapsto & \alpha\tau \\ T_i & \longmapsto & q - e_i & \longmapsto & q - e_i \end{array}$$

and every irreducible representation of \widehat{TL}_k^a factors through one of these homomorphisms (see [GL2, Prop. 4.14(v)]). In Proposition 3.2 we shall see that these homomorphisms arise naturally in the Schur-Weyl duality setting.

Remark 2.6. Following [GL1, Cor. (5.11) and Thm. (6.1)], for any irreducible \widehat{TL}_k^a -module there is a constant x such that

$$(2.14) \quad e_1 X^{-\varepsilon_1} e_1 = x e_1.$$

In other words, the representation theory (at least the full information of the cell modules, see [GL1, §6], and irreducible modules) is completely contained in the algebras obtained by adding the relation (2.14) to \widehat{TL}_k^a .

2.5. A commuting family in the affine Temperley-Lieb algebra

View the elements $X^{-\varepsilon_i}$ in the affine Temperley-Lieb algebra \widehat{TL}_k^a via the surjective algebra homomorphism of (2.13). Define

$$(2.15) \quad \begin{aligned} (q - q^{-1})m_1 &= q^{-1}X^{-\varepsilon_1} \quad \text{and} \\ (q - q^{-1})m_i &= q^{i-2}(X^{-\varepsilon_i} - q^{-2}X^{-\varepsilon_{i-1}}), \end{aligned}$$

for $i = 2, 3, \dots, k$. Since $X^{-\varepsilon_i}X^{-\varepsilon_j} = X^{-\varepsilon_j}X^{-\varepsilon_i}$ for all $1 \leq i, j \leq k$, and the m_i are linear combinations of the $X^{-\varepsilon_i}$,

$$m_i m_j = m_j m_i \text{ in } \widehat{TL}_k^a, \quad \text{for all } 1 \leq i, j \leq k.$$

PROPOSITION 2.7. For $1 \leq i \leq k$,

- (a) $X^{-\varepsilon_i} = q^{-(i-2)}(q - q^{-1})(m_i + q^{-1}m_{i-1} + q^{-2}m_{i-2} + \cdots + q^{-(i-1)}m_1)$,
- (b) $X^{-\varepsilon_1} + \cdots + X^{-\varepsilon_i} = q^{-(i-2)}(q - q^{-1})(m_i + [2]m_{i-1} + \cdots + [i]m_1)$.

Proof. Rewrite (2.15) as

$$X^{-\varepsilon_i} = q^{-(i-2)}(q - q^{-1})m_i + q^{-2}X^{-\varepsilon_{i-1}}$$

and use induction,

$$\begin{aligned} X^{-\varepsilon_i} &= q^{-(i-2)}(q - q^{-1})m_i \\ &\quad + q^{-2}(q^{-(i-1-2)}(q - q^{-1})(m_{i-1} + q^{-1}m_{i-2} + \cdots + q^{-(i-2)}m_1)), \end{aligned}$$

to obtain the formula for $X^{-\varepsilon_i}$ in (a). Summing the formula in (a) over i gives

$$\begin{aligned} \sum_{j=1}^i X^{-\varepsilon_j} &= \sum_{j=1}^i \left(q^{-(j-2)}(q - q^{-1}) \sum_{\ell=0}^{j-2} q^{-\ell} m_{j-\ell} \right) \\ &= q^{-(i-2)}(q - q^{-1}) \sum_{j=1}^i \sum_{\ell=0}^{j-1} q^{i-j-\ell} m_{j-\ell} \end{aligned}$$

and, thus, formula (b) follows from

$$\begin{aligned} \sum_{j=1}^i \sum_{\ell=0}^{j-1} q^{i-j-\ell} m_{j-\ell} &= \sum_{j=1}^i \sum_{r=1}^j q^{i-j-(j-r)} m_r = \sum_{r=1}^i \sum_{j=r}^i q^{i+r-2j} m_r \\ &= \sum_{r=1}^i [i - r + 1] m_r. \end{aligned}$$

□

The following Lemma is a transfer of the recursion $X^{\varepsilon_i} = T_{i-1}X^{\varepsilon_{i-1}}T_{i-1}$ to the m_i . The following are the base cases of Lemma 2.8.

$$m_1 = \frac{q^{-1}}{q - q^{-1}}X^{-\varepsilon_1} \quad \text{and} \quad m_2 = \frac{x}{q - q^{-1}}e_1 - (e_1m_1 + m_1e_1)$$

LEMMA 2.8. *Following Remark 2.6, assume that $e_1 X^{-\varepsilon_1} e_1 = x e_1$ for some constant x . For $2 \leq i \leq k$,*

$$m_i = \frac{q^{i-2}x}{q - q^{-1}} e_{i-1} - (e_{i-1} m_{i-1} + m_{i-1} e_{i-1}) - \sum_{\ell=1}^{i-2} ([i - \ell] - [i - \ell - 2]) m_\ell e_{i-1}.$$

Proof. From (2.3) and (2.9) we have $X^{-\varepsilon_i} = (q^{-1} - e_{i-1}) X^{-\varepsilon_{i-1}} (q^{-1} - e_{i-1})$. Substituting this into the definition of m_i gives

$$\begin{aligned} (q - q^{-1}) m_i &= q^{i-2} (X^{-\varepsilon_i} - q^{-2} X^{-\varepsilon_{i-1}}) \\ &= q^{i-2} (q^{-1} - e_{i-1}) X^{-\varepsilon_{i-1}} (q^{-1} - e_{i-1}) - q^{i-4} X^{-\varepsilon_{i-1}} \\ &= q^{i-2} e_{i-1} X^{-\varepsilon_{i-1}} e_{i-1} - q^{i-3} (e_{i-1} X^{-\varepsilon_{i-1}} + X^{-\varepsilon_{i-1}} e_{i-1}). \end{aligned}$$

Use Proposition 2.7 (a) to substitute for $X^{-\varepsilon_{i-1}}$,

$$\begin{aligned} (q - q^{-1}) m_i &= (q - q^{-1}) q^{-(m+i-3)} (q^{i-2} e_{i-1} m_{i-1} e_{i-1} - q^{i-3} (e_{i-1} m_{i-1} + m_{i-1} e_{i-1})) \\ &\quad + (q - q^{-1}) q^{-(i-3)} (q^{-1} m_{i-2} + \cdots + q^{-(i-2)} m_1) (q^{i-2} e_{i-1}^2 - 2q^{i-3} e_{i-1}) \\ &= (q - q^{-1}) (q e_{i-1} m_{i-1} e_{i-1} - (e_{i-1} m_{i-1} + m_{i-1} e_{i-1})) \\ &\quad + (q - q^{-1}) (m_{i-2} + \cdots + q^{-(i-3)} m_1) (q + q^{-1} - 2q^{-1}) e_{i-1} \\ &= (q - q^{-1}) \left(\begin{array}{c} q e_{i-1} m_{i-1} e_{i-1} - (e_{i-1} m_{i-1} + m_{i-1} e_{i-1}) \\ + (q - q^{-1}) (m_{i-2} + \cdots + q^{-(i-3)} m_1) e_{i-1} \end{array} \right), \end{aligned}$$

which gives

$$(2.16) \quad m_i = q e_{i-1} m_{i-1} e_{i-1} - (e_{i-1} m_{i-1} + m_{i-1} e_{i-1}) + (q - q^{-1}) (m_{i-2} + q^{-1} m_{i-3} + q^{-2} m_{i-4} + \cdots + q^{-(i-3)} m_1) e_{i-1}.$$

Using induction, substitute for the first m_{i-1} in this equation to get

$$\begin{aligned} m_i &= -(e_{i-1} m_{i-1} + m_{i-1} e_{i-1}) + (q - q^{-1}) \sum_{\ell=1}^{i-2} q^{-(i-2-\ell)} m_\ell e_{i-1} \\ &\quad + q \left(\frac{q^{i-3}x}{q - q^{-1}} e_{i-1} - 2m_{i-2} e_{i-1} - \sum_{\ell=1}^{i-3} ([i - \ell - 1] - [i - \ell - 3]) m_\ell e_{i-1} \right) \\ &= \frac{q^{i-2}x}{q - q^{-1}} e_{i-1} - (e_{i-1} m_{i-1} + m_{i-1} e_{i-1}) - \sum_{\ell=2}^{i-2} ([i - \ell] - [i - \ell - 2]) m_\ell e_{i-1}. \end{aligned}$$

□

2.6. Diagram representation of Murphy elements

Label the vertices from left to right in the top row of a diagram $d \in T_k$ with $1, 2, \dots, k$, and label the corresponding vertices in the bottom row with $1', 2', \dots, k'$. The *cycle type* of a diagram $d \in T_k$ is the set partition $\tau(d)$ of $\{1, 2, \dots, k\}$ obtained from d by setting $1 = 1', 2 = 2', \dots, k = k'$. If $\tau(d)$ is a set partition of the form $\{\{1, 2, \dots, \gamma_1\}, \{\gamma_1 + 1, \mu_1 + 2, \dots, \gamma_1 + \gamma + 2\}, \dots, \{\gamma_1 + \dots + \gamma_{\ell-1} + 1, \dots, k\}\}$, where $(\gamma_1, \dots, \gamma_\ell)$ is a composition of k , then we simplify notation by writing $\tau(d) = (\gamma_1, \dots, \gamma_\ell)$. For example




There are diagrams whose cycle type cannot be written as a composition (for example $d = \begin{matrix} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \end{matrix}$ has cycle type $\{\{1, 4\}, \{2, 3\}\}$) but all of the diagrams needed here have cycle types that are compositions.

If $\gamma = (\gamma_1, \dots, \gamma_\ell)$ is a composition of k define

$$(2.17) \quad d_\gamma = \sum_{\tau(d)=\gamma} d$$

as the sum of the Temperley-Lieb diagrams on k dots with cycle type γ . Define d_γ^* be the sum of diagrams obtained from the summands of d_γ by wrapping the first edge in each row around the pole, with the orientation coming from $X^{-\varepsilon_1}$ as shown in the examples below. When the first edge in the top row connects to the first vertex in the bottom row only one new diagram is produced, otherwise there are two. For example, in \widehat{TL}_4^a ,



View d_γ and d_γ^* as elements of \widehat{TL}_k^a by setting

$$d_\gamma = d_{\gamma_{1^{k-i}}}, \quad \text{if } \gamma \text{ is a composition of } i \text{ with } i < k.$$

With this notation, expanding the first few m_i in terms of diagrams gives

$$\begin{aligned}
(q - q^{-1})m_1 &= q^{-1}d_1^*, & (q - q^{-1})m_2 &= xd_2 - q^{-1}d_2^*, \\
(q - q^{-1})m_3 &= qxd_{1,2} - q^{-1}[2]d_{1,2}^* - xd_3 + q^{-1}d_3^*, \\
(q - q^{-1})m_4 &= q^2xd_{1^2,2} - q^{-1}([3] - [1])d_{1^2,2}^* - x[2]d_{2,2} + q^{-1}[2]d_{2,2}^* \\
&\quad - qxd_{1,3} + q^{-1}[2]d_{1,3}^* + xd_4 - q^{-1}d_4^*, \\
(q - q^{-1})m_5 &= q^3xd_{1^3,2} - q^{-1}([4] - [2])d_{1^3,2}^* - q^2xd_{1^2,3} + q^{-1}([3] - [1])d_{1^2,3}^* \\
&\quad + qxd_{1,4} - q^{-1}[2]d_{1,4}^* - qx[2]d_{1,2,2} + q^{-1}[2]^2d_{1,2,2}^* + x[2]d_{2,3} \\
&\quad - q^{-1}[2]d_{2,3}^* - x([3] - [1])d_{2,1,2} + q^{-1}([3] - [1])d_{2,1,2}^* \\
&\quad + x[2]d_{3,2} - q^{-1}[2]d_{3,2}^* - xd_5 + q^{-1}d_5^*,
\end{aligned}$$

where, as in Lemma 2.8, x is the constant defined by the equation $e_1X^{-\varepsilon_1}e_1 = xe_1$.

THEOREM 2.9. *Following Remark 2.6, assume that $e_1X^{-\varepsilon_1}e_1 = xe_1$ for some constant x . Then $(q - q^{-1})m_1 = q^{-1}d_1^*$, $(q - q^{-1})m_2 = xd_2 - q^{-1}d_2^*$ and, for $i \geq 2$,*

$$m_i = \sum_{\text{compositions } \gamma} (m_i)_\gamma d_\gamma + (m_i)_\gamma^* d_\gamma^*,$$

where the sum is over all compositions $\gamma = 1^{b_1}r_11^{b_2}r_2 \cdots 1^{b_\ell}r_\ell$ of i with $r_\ell > 1$, and

$$\begin{aligned}
(m_i)_\gamma &= (-1)^{|\gamma| - \ell(\gamma) - 1} \frac{q^{b_1}x}{q - q^{-1}} \prod_{j>1} ([b_j + 2] - [b_j]), \quad \text{and} \\
(m_i)_\gamma^* &= (-1)^{|\gamma| - \ell(\gamma)} \frac{q^{-1}}{q - q^{-1}} ([b_1 + 1] - [b_1 - 1]) \prod_{j>1} ([b_j + 2] - [b_j]),
\end{aligned}$$

with $\ell(\gamma) = \ell + b_1 + \cdots + b_\ell$.

Proof. From our computations above, $m_1 = Ad_1^*$ and $m_2 = Bd_2 - Ad_2^*$, where

$$A = \frac{q^{-1}}{q - q^{-1}} \quad \text{and} \quad B = \frac{x}{q - q^{-1}}.$$

Let $m_1 = Ad_1^*$. For $i > 2$ the recursion in Lemma 2.8 gives

$$\begin{aligned} m_i &= q^{i-2} B e_{i-1} - (e_{i-1} m_{i-1} + m_{i-1} e_{i-1}) - \sum_{\ell=1}^{i-2} ([i-\ell] - [i-\ell-2]) m_\ell e_{i-1} \\ &= q^{i-2} B d_{1^{i-2}, 2} - \left(\sum_{\gamma=\gamma' r} ((m_{i-1})_{\gamma' r} d_{\gamma', r+1} + (m_{i-1})_{\gamma' r}^* d_{\gamma', r+1}^*) \right) \\ &\quad - ([i-1] - [i-3]) A d_{1^{i-2}, 2} \\ &\quad + \sum_{\ell=2}^{i-2} -([i-\ell] - [i-\ell-2]) \sum_{\gamma'} ((m_\ell)_{\gamma'} d_{\gamma' 1^{i-2-\ell_2}} + (m_\ell)_{\gamma'}^* d_{\gamma' 1^{i-2-\ell_2}}^*). \end{aligned}$$

So if d has cycle type $\gamma = 1^{b_1} r_1 1^{b_2} r_2 \cdots 1^{b_\ell} r_\ell$ with $r_\ell > 0$, then

- (a) Each part of size r ($r > 1$) contributes $(-1)^{r-1}$ to the coefficient. Thus, there is a total contribution of $(-1)^{|\gamma|-\ell(\gamma)}$ from these parts.
- (b) Each inner 1^b ($b \geq 0$) contributes a factor of $[b+2] - [b]$ to the coefficient.
- (c) The first 1^b ($b > 0$) contributes a $-q^b B$ in a nonstarred class,
- (c') The first 1^b ($b = 0$) contributes a $-B$ in a nonstarred class, which is the same as case (c) with $b = 0$.
- (d) The first 1^b ($b > 0$) contributes a $([b+1] - [b-1])A$ in a starred class.
- (d') The first 1^b ($b = 0$) contributes an A in a starred class, which is the same as case (d) with $b = 0$ assuming $[-1] = 0$. \square

Remark 2.10. To view m_1, \dots, m_k in the (nonaffine) Temperley-Lieb algebra $TL_k(n)$ (via (2.11)) let $X^{-\varepsilon_1} = 1$ so that $x = q + q^{-1}$. If $b_1 > 1$ then $d_\gamma^* = d_\gamma$ and if $b_1 = 0$ then $d_\gamma^* = 2d_\gamma$. In both cases the coefficients in Theorem 2.9 specialize to

$$(m_i)_\gamma + (m_i)_\gamma^* = (-1)^{|\gamma|-\ell(\gamma)-1} [b_1 + 1] \prod_{b_j \geq 0, j > 1} ([b_j + 2] - [b_j])$$

and

$$m_i = \sum_{\gamma} ((m_i)_\gamma + (m_i)_\gamma^*) d_\gamma,$$

where the sum is over compositions $\gamma = 1^{b_1}r_1 1^{b_2}r_2 \dots 1^{b_\ell}r_\ell$ of i with $r_\ell > 1$. The first few examples are

$$\begin{aligned}
 m_1 &= \frac{q^{-1}}{q - q^{-1}} = \frac{q^{-1}}{q - q^{-1}}d_1, & m_2 &= e_2 = d_2, & m_3 &= [2]d_{12} - d_3, \\
 m_4 &= [3]d_{1^2,2} - [2]d_{2,2} - [2]d_{1,3} + d_4, \\
 m_5 &= [4]d_{1^3,2} - [3]d_{1^2,3} + [2]d_{1,4} - [2]^2d_{1,2,2} + [2]d_{2,3} - ([3] - [1])d_{2,1,2} \\
 & \qquad \qquad \qquad + [2]d_{3,2} - d_5.
 \end{aligned}$$

§3. Schur functors

3.1. R -matrices and quantum Casimir elements

Let $U_h\mathfrak{g}$ be the Drinfeld-Jimbo quantum group corresponding to a finite dimensional complex semisimple Lie algebra \mathfrak{g} . We shall use the notations and conventions for $U_h\mathfrak{g}$ as in [LR] and [OR]. There is an invertible element $\mathcal{R} = \sum a_i \otimes b_i$ in (a suitable completion of) $U_h\mathfrak{g} \otimes U_h\mathfrak{g}$ such that, for two $U_h\mathfrak{g}$ modules M and N , the map

$$\begin{aligned}
 \check{R}_{MN}: M \otimes N &\longrightarrow N \otimes M & \begin{array}{c} M \otimes N \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ N \otimes M \end{array} \\
 m \otimes n &\longmapsto \sum b_i n \otimes a_i m & \begin{array}{c} \bullet \quad \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ N \otimes M \end{array}
 \end{aligned}$$

is a $U_h\mathfrak{g}$ module isomorphism. In order to be consistent with the graphical calculus these operators should be written *on the right*. The element \mathcal{R} satisfies ‘‘quasitriangularity relations’’ (see [LR, (2.1)–(2.3)]) which imply that, for $U_h\mathfrak{g}$ modules M, N, P and a $U_h\mathfrak{g}$ module isomorphism $\tau_M: M \rightarrow M$,

$$\begin{aligned}
 \begin{array}{c} M \otimes N \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ N \otimes M \end{array} &= \begin{array}{c} M \otimes N \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ N \otimes M \end{array} \\
 \check{R}_{MN}(\text{id}_N \otimes \tau_M) &= (\tau_M \otimes \text{id}_N)\check{R}_{MN}, \\
 \begin{array}{c} M \otimes (N \otimes P) \\ \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \bullet \\ (N \otimes P) \otimes M \end{array} &= \begin{array}{c} M \otimes N \otimes P \\ \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \bullet \\ N \otimes P \otimes M \end{array} \\
 \check{R}_{M,N \otimes P} &= (\check{R}_{MN} \otimes \text{id}_P)(\text{id}_N \otimes \check{R}_{MP})
 \end{aligned}$$

$$\begin{array}{c}
 (M \otimes N) \otimes P \\
 \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ P \otimes (M \otimes N) \end{array}
 \end{array}
 =
 \begin{array}{c}
 M \otimes N \otimes P \\
 \begin{array}{c} \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \\ \diagup \quad \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \\ P \otimes M \otimes N \end{array}
 \end{array}$$

$$\check{R}_{M \otimes N, P} = (\text{id}_M \otimes \check{R}_{NP})(\check{R}_{MP} \otimes \text{id}_N),$$

which, together, imply the braid relation

$$\begin{array}{c}
 M \otimes N \otimes P \\
 \begin{array}{c} \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \\ \diagup \quad \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \\ P \otimes N \otimes M \end{array}
 \end{array}
 =
 \begin{array}{c}
 M \otimes N \otimes P \\
 \begin{array}{c} \bullet \quad \bullet \quad \bullet \\ \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \\ \diagup \quad \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \\ P \otimes N \otimes M \end{array}
 \end{array}$$

$$\begin{aligned}
 (\check{R}_{MN} \otimes \text{id}_P)(\text{id}_N \otimes \check{R}_{MP})(\check{R}_{NP} \otimes \text{id}_M) \\
 = (\text{id}_M \otimes \check{R}_{NP})(\check{R}_{MP} \otimes \text{id}_N)(\text{id}_P \otimes \check{R}_{MN}).
 \end{aligned}$$

Let ρ be such that $\langle \rho, \alpha_i \rangle = 1$ for all simple roots α_i . As explained in [LR, (2.14)] and [Dr], there is a *quantum Casimir element* $e^{-h\rho u}$ in the center of $U_h\mathfrak{g}$ and, for a $U_h\mathfrak{g}$ module M we define a $U_h\mathfrak{g}$ module isomorphism

$$C_M: M \longrightarrow M \qquad \begin{array}{c} M \\ \bullet \\ \bullet \\ M \end{array} C_M \\
 m \longmapsto (e^{-h\rho u})m$$

and the elements C_M satisfy

$$(3.1) \quad C_{M \otimes N} = (\check{R}_{MN} \check{R}_{NM})^{-1}(C_M \otimes C_N), \quad \text{and} \quad C_M = q^{-\langle \lambda, \lambda + 2\rho \rangle} \text{id}_M$$

if M is a $U_h\mathfrak{g}$ module generated by a highest weight vector v^+ of weight λ (see [LR, Prop. 2.14] or [Dr, Prop. 3.2]). Note that $\langle \lambda, \lambda + 2\rho \rangle = \langle \lambda + \rho, \lambda + \rho \rangle - \langle \rho, \rho \rangle$ are the eigenvalues of the classical Casimir operator [Dx, 7.8.5]. From the relation (3.1) it follows that if $M = L(\mu)$, $N = L(\nu)$ are finite dimensional irreducible $U_h\mathfrak{g}$ modules then $\check{R}_{MN} \check{R}_{NM}$ acts on the λ isotypic component $L(\lambda)^{\oplus c_{\mu\nu}^\lambda}$ of the decomposition

$$\begin{aligned}
 (3.2) \quad L(\mu) \otimes L(\nu) &= \bigoplus_{\lambda} L(\lambda)^{\oplus c_{\mu\nu}^\lambda} \\
 &\text{by the constant } q^{\langle \lambda, \lambda + 2\rho \rangle - \langle \mu, \mu + 2\rho \rangle - \langle \nu, \nu + 2\rho \rangle}.
 \end{aligned}$$

3.2. The $\tilde{\mathcal{B}}_k$ module $M \otimes V^{\otimes k}$

Let $U = U_h \mathfrak{g}$ be a quantum group and let M and V be U -modules such that the operators \check{R}_{MV} , \check{R}_{VM} and \check{R}_{VV} are well defined. Define \check{R}_i , $1 \leq i \leq k - 1$, and \check{R}_0^2 in $\text{End}_U(M \otimes V^{\otimes k})$ by

$$\begin{aligned} \check{R}_i &= \text{id}_M \otimes \text{id}_V^{\otimes(i-1)} \otimes \check{R}_{VV} \otimes \text{id}_V^{\otimes(k-i-1)} \quad \text{and} \\ \check{R}_0^2 &= (\check{R}_{MV} \check{R}_{VM}) \otimes \text{id}_V^{\otimes(k-1)}. \end{aligned}$$

Then the braid relations

$$\check{R}_i \check{R}_{i+1} \check{R}_i = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} = \check{R}_{i+1} \check{R}_i \check{R}_{i+1}$$

and

$$\check{R}_0^2 \check{R}_1 \check{R}_0^2 \check{R}_1 = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} = \begin{array}{c} \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \end{array} = \check{R}_1 \check{R}_0^2 \check{R}_1 \check{R}_0^2.$$

imply that there is a well defined map

$$(3.3) \quad \begin{array}{lll} \Phi: \tilde{\mathcal{B}}_k & \longrightarrow & \text{End}_U(M \otimes V^{\otimes k}) \\ T_i & \longmapsto & \check{R}_i, \quad 1 \leq i \leq k - 1, \\ X^{\varepsilon_1} & \longmapsto & \check{R}_0^2, \end{array}$$

which makes $M \otimes V^{\otimes k}$ into a right $\tilde{\mathcal{B}}_k$ module. By (3.1) and the fact that (3.4)

$$\Phi(X^{\varepsilon_i}) = \check{R}_{M \otimes V^{\otimes(i-1)}, V} \check{R}_{V, M \otimes V^{\otimes(i-1)}} = \begin{array}{c} \text{---} \quad \text{---} \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \quad \diagup \quad \diagdown \quad \diagup \quad \diagdown \quad \diagup \quad \diagdown \\ \text{---} \quad \text{---} \end{array}.$$

the eigenvalues of $\Phi(X^{\varepsilon_i})$ are related to the eigenvalues of the Casimir. The *Schur functors* are the functors

$$(3.5) \quad \begin{array}{ll} F_V^\lambda: \{U\text{-modules}\} & \longrightarrow \{\tilde{\mathcal{B}}_k\text{-modules}\} \\ M & \longmapsto \text{Hom}_U(M(\lambda), M \otimes V^{\otimes k}) \end{array}$$

where $\text{Hom}_U(M(\lambda), M \otimes V^{\otimes k})$ is the vector space of highest weight vectors of weight λ in $M \otimes V^{\otimes k}$.

are the contents of the boxes for the partition in (3.8).

If ν is a partition and

$$(3.10) \quad V = L(\square) \quad \text{then} \quad L(\nu) \otimes V = \bigoplus_{\lambda \in \nu^+} L(\lambda),$$


where the sum is over all partitions λ with $\leq n$ rows that are obtained by adding a box to ν [Mac, I App. A (8.4) and I (5.16)], Hence, the $U_h \mathfrak{gl}_n$ -module decompositions of

$$(3.11) \quad L(\mu) \otimes V^{\otimes k} = \bigoplus_{\lambda} L(\lambda) \otimes \tilde{H}_k^{\lambda/\mu}, \quad k \in \mathbb{Z}_{\geq 0},$$

are encoded by the graph \hat{H}^{μ} with

$$(3.12) \quad \begin{array}{ll} \text{vertices on level } k: & \{\text{skew shapes } \lambda/\mu \text{ with } k \text{ boxes}\} \\ \text{edges:} & \lambda/\mu \longrightarrow \gamma/\mu, \text{ if } \gamma \text{ is obtained from } \lambda \\ & \text{by adding a box} \\ \text{labels on edges:} & \text{content of the added box.} \end{array}$$


For example if $\mu = (3, 3, 3, 2) = \square$, then the first 4 rows of \hat{H}^{μ} are



The following result is well known (see [Ji] or [LR, (4.4)]).


PROPOSITION 3.1. *If $U = U_h \mathfrak{gl}_n$ and $V = L(\varepsilon_1) = L(\square)$ is the n -dimensional “standard” representation of \mathfrak{gl}_n then the map Φ of (3.3) factors through the surjective homomorphism (2.8) to give a representation of the affine Hecke algebra.*

For a skew shape λ/μ with k boxes identify paths from μ to λ/μ in \hat{H}^μ with *standard tableaux* of shape λ/μ by filling the boxes, successively, with $1, 2, \dots, k$ as they appear. In the example graph \hat{H}^μ above



3.4. The quantum group $U_h\mathfrak{gl}_2$

In the case when $n = 2$, $U = U_h\mathfrak{gl}_2$ and the partitions which appear in (3.11) and in the graph \hat{H}^μ all have ≤ 2 rows. For example if $\mu = (42) = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$ then the first few rows of \hat{H}^μ are




and this is the graph which describes the decompositions in (3.11).

PROPOSITION 3.2. *If $U = U_h\mathfrak{gl}_2$, $M = L(\mu)$ where μ is a partition of m with ≤ 2 rows, and $V = L(\varepsilon_1) = L(\square)$ is the 2-dimensional “standard” representation of \mathfrak{gl}_2 then the map Φ of (3.3) factors through the surjective homomorphism of (2.13) with $\alpha^2 = -q^{2m-1}$ to give a representation of the affine Temperley-Lieb algebra T_k^a .*

Proof. The proof that the kernel of Φ contains the element (2.12) is exactly as in the proof of [OR, Thm. 6.1(c)]: The element e_1 in T_2^a acts on

$V^{\otimes 2}$ as $(q + q^{-1}) \cdot \text{pr}$ where pr is the unique $U_h \mathfrak{g}$ -invariant projection onto $L(\mathfrak{H})$ in $V^{\otimes 2}$. Using $e_1 T_1 = -q^{-1} e_1$ and the pictorial equalities



it follows that $\Phi_2(e_1 X^{\varepsilon_1} T_1 X^{\varepsilon_1})$ acts as $-(q + q^{-1})q^{-1} \cdot \check{R}_{L(\mathfrak{H}), L(\mu)} \check{R}_{L(\mu), L(\mathfrak{H})} (\text{id}_{L(\mu)} \otimes \text{pr})$. By (3.1), this is equal to

$$\begin{aligned}
 & -q^{-1} (C_{L(\mu)} \otimes C_{L(\mathfrak{H})}) C_{L(\mu) \otimes L(\mathfrak{H})}^{-1} \Phi_2(\text{id}_{L(\mu)} \otimes e_1) \\
 & = -q^{-1} q^{-\langle \mu, \mu + 2\rho \rangle} q^{-\langle \varepsilon_1 + \varepsilon_2, \varepsilon_1 + \varepsilon_2 + 2\rho \rangle} C_{L(\mu + \varepsilon_1 + \varepsilon_2)}^{-1} \Phi_2(\text{id}_{L(\mu)} \otimes e_1),
 \end{aligned}$$

and the coefficient $-q^{-1} q^{-\langle \mu, \mu + 2\rho \rangle} q^{-\langle \varepsilon_1 + \varepsilon_2, \varepsilon_1 + \varepsilon_2 + 2\rho \rangle} C_{L(\mu + \varepsilon_1 + \varepsilon_2)}^{-1}$ simplifies to

$$\begin{aligned}
 & -q^{-1} q^{-\langle \mu, \mu + 2\rho \rangle} q^{-\langle \varepsilon_1 + \varepsilon_2, \varepsilon_1 + \varepsilon_2 + 2\rho \rangle} q^{\langle \mu + \varepsilon_1 + \varepsilon_2, \mu + \varepsilon_1 + \varepsilon_2 + 2\rho \rangle} \\
 & = -q^{-1} q^{2(\mu_1 + \mu_2)} = -q^{2m-1},
 \end{aligned}$$

where $m = \mu_1 + \mu_2 = |\mu|$. □

3.5. The quantum group $U_h \mathfrak{sl}_2$


The restriction of an irreducible representation $L(\lambda)$ of $U_h \mathfrak{gl}_n$ to $U_h \mathfrak{sl}_n$ is irreducible and all irreducible representations of $U_h \mathfrak{sl}_n$ are obtained in this fashion. Since the “determinant” representation is trivial as an $U_h \mathfrak{sl}_n$ module it follows from (3.7) that the irreducible representations $L_{\mathfrak{sl}_n}(\lambda)$ of $U_h \mathfrak{sl}_n$ are indexed by partitions $\lambda = (\lambda_1, \dots, \lambda_n)$ with $\lambda_n = 0$. Hence, the graph which describes the $U_h \mathfrak{sl}_2$ -module decompositions of

$$(3.15) \quad L(\mu) \otimes V^{\otimes k} = \bigoplus_{\lambda} L(\lambda) \otimes \tilde{T}_k^{\lambda/\mu}, \quad k \in \mathbb{Z}_{\geq 0},$$

is exactly the same as the graph for $U_h \mathfrak{gl}_2$ except with all columns of length 2 removed from the partitions. More precisely, the decompositions are encoded by the graph $\hat{T}^{\lambda/\mu}$ with

$$(3.16) \quad \begin{aligned}
 \text{vertices on level } k: & \quad \{\mu_1 - \mu_2 + k, \mu_1 - \mu_2 + k - 2, \dots, \mu_1 - \mu_2 - k\} \cap \mathbb{Z}_{\geq 0} \\
 \text{edges:} & \quad \ell \longrightarrow \ell \pm 1.
 \end{aligned}$$

For example if $m = 7$ and $\mu_1 - \mu_2 = 3$ then the first few rows of $\hat{T}^{\lambda/\mu}$ are



Paths in (3.17) correspond to paths in (3.14) which correspond to standard tableaux T of shape λ/μ .

§4. Eigenvalues

4.1. Eigenvalues of the X^{ε_i} in the affine Hecke algebra

Recall, from (2.8), that the *affine Hecke algebra* \tilde{H}_k is the quotient of the group algebra of the affine braid group $\mathbb{C}\tilde{B}_k$ by the relations

$$(4.1) \quad T_i^2 = (q - q^{-1})T_i + 1.$$

As observed in Proposition 3.2 the map Φ in (3.3) makes the module $L(\mu) \otimes V^{\otimes k}$ in (3.11) into an \tilde{H}_k module. Thus the vector spaces $\tilde{H}_k^{\lambda/\mu}$ in (3.11) are the \tilde{H}_k -modules given by

$$\tilde{H}_k^{\lambda/\mu} = F_V^\lambda(L(\mu)), \quad \text{where } F_V^\lambda \text{ are the Schur functors of (3.5).}$$

The following theorem is well known (see, for example, [Ch]).

THEOREM 4.1. (a) *The X^{ε_i} , $1 \leq i \leq k$, mutually commute in the affine Hecke algebra \tilde{H}_k .*

(b) The eigenvalues of X^{ε_i} are given by the graph $\hat{H}^{\lambda/\mu}$ of (3.13) in the sense that if

$$\begin{aligned} \hat{H}_k^{\lambda/\mu} &= \{\text{skew shapes } \lambda/\mu \text{ with } k \text{ boxes}\} \quad \text{and} \\ \hat{H}_k^{\lambda/\mu} &= \{\text{standard tableaux } T \text{ of shape } \lambda/\mu\} \end{aligned}$$

for $\lambda/\mu \in \hat{H}_k^{\lambda/\mu}$, then

$$\begin{aligned} \hat{H}_k^{\lambda/\mu} &\text{ is an index set for the simple } \tilde{H}_k \text{ modules } \tilde{H}_k^{\lambda/\mu} \\ &\text{ appearing in } L(\mu) \otimes V^{\otimes k}, \end{aligned}$$

and

$$\tilde{H}_k^{\lambda/\mu} \text{ has a basis } \{v_T \mid T \in \hat{H}_k^{\lambda/\mu}\} \text{ with } X^{\varepsilon_i} v_T = q^{2c(T(i))} v_T,$$

where $c(T(i))$ is the content of box i of T .

(c) $\kappa = X^{\varepsilon_1} \cdots X^{\varepsilon_k}$ is a central element of \tilde{H}_k and

$$\kappa \text{ acts on } \tilde{H}_k^{\lambda/\mu} \text{ by the constant } q^{2\sum_{b \in \lambda/\mu} c(b)}.$$

Proof. (a) is a restatement of (2.4). (b) Since the \tilde{H}_k action and the $U_h \mathfrak{gl}_n$ action commute on $L(\mu) \otimes V^{\otimes k}$ it follows that the decomposition in (3.11) is a decomposition as $(U_h \mathfrak{gl}_n, \tilde{H}_k)$ bimodules, where the $\tilde{H}_k^{\lambda/\mu}$ are some \tilde{H}_k -modules. Comparing the $L(\lambda)$ components on each side of

$$\begin{aligned} \bigoplus_{\lambda} L(\lambda) \otimes \tilde{H}_{\ell}^{\lambda/\mu} &\cong L(\mu) \otimes V^{\otimes \ell} = L(\mu) \otimes V^{\otimes(\ell-1)} \otimes V \\ &\cong \left(\bigoplus_{\nu} L(\nu) \otimes \tilde{H}_{\ell-1}^{\nu/\mu} \right) \otimes V \\ &\cong \bigoplus_{\lambda} \bigoplus_{\lambda/\nu=\square} L(\lambda) \otimes \tilde{H}_{\ell-1}^{\nu/\mu} \\ &\cong \bigoplus_{\lambda} \left(L(\lambda) \otimes \left(\bigoplus_{\lambda/\nu=\square} \tilde{H}_{\ell-1}^{\nu/\mu} \right) \right) \end{aligned}$$

gives

$$(4.2) \quad \tilde{H}_{\ell}^{\lambda/\mu} \cong \bigoplus_{\lambda/\nu=\square} \tilde{H}_{\ell-1}^{\nu/\mu},$$

for any $\ell \in \mathbb{Z}_{\geq 0}$ and skew shape λ/μ with ℓ boxes. Iterate (4.2) (with $\ell = k, k - 1, \dots$) to produce a decomposition

$$\tilde{H}_k^{\lambda/\mu} = \bigoplus_{T \in \hat{H}_k^{\lambda/\mu}} \tilde{H}_1^T,$$

where the summands \tilde{H}_1^T are 1-dimensional vector spaces. This determines a basis (unique up to multiplication of the basis vectors by constants) $\{v_T \mid T \in \hat{H}_k^{\lambda/\mu}\}$ of $\tilde{H}_k^{\lambda/\mu}$ which respects the decompositions in (4.2) for $1 \leq \ell \leq k$.

Combining (3.1), (3.2) and (3.4) gives that X^{ε_i} acts on the $L(\lambda)$ component of the decomposition (3.10) by the constant

$$q^{\langle \lambda, \lambda + 2\rho \rangle - \langle \nu, \nu + 2\rho \rangle - \langle \varepsilon_1, \varepsilon_1 + 2\rho \rangle} = q^{2c(\lambda/\nu)}$$

since if $\lambda = \nu + \varepsilon_j$, so that λ is the same as ν except with an additional box in row j , then $\nu \subseteq \lambda$, $\lambda/\nu = \square$ and

$$\begin{aligned} & \langle \lambda, \lambda + 2\rho \rangle - \langle \nu, \nu + 2\rho \rangle - \langle \varepsilon_1, \varepsilon_1 + 2\rho \rangle \\ &= \langle \nu + \varepsilon_j, \nu + \varepsilon_j + 2\rho \rangle - \langle \nu, \nu + 2\rho \rangle - (1 + 2(n - 1)) \\ &= 2\nu_j + \langle \varepsilon_j, \varepsilon_j + 2\rho \rangle - 2n + 1 = 2\nu_j + (1 + 2(n - j)) - 2n + 1 \\ &= 2(\nu_j + 1) - 2j = 2c(\lambda/\nu). \end{aligned}$$

Hence,

$$X^{\varepsilon_i} v_T = q^{2c(T(i))} v_T, \quad \text{for } 1 \leq i \leq k,$$

where $T(i)$ is the box containing i in T .

The remainder of the proof, including the simplicity of the \tilde{H}_k -modules $\tilde{H}_k^{\lambda/\mu}$, is accomplished as in [R, Thm. 4.1].

(c) The element $X^{\varepsilon_1} \dots X^{\varepsilon_k}$ is central in \tilde{B}_k (it is a full twist) and hence its image is central in \tilde{H}_k . The constant describing its action on $\tilde{H}_k^{\lambda/\mu}$ follows from the formula $X^{\varepsilon_i} v_T = q^{2c(T(i))} v_T$. \square

4.2. Eigenvalues of the m_i in T_k^a

Let m_1, m_2, \dots, m_k be the commuting family in the affine Temperley-Lieb algebra as defined in (2.15). We will use the results of Theorem 4.1 to determine the eigenvalues of the m_i in the (generically) irreducible representations.

THEOREM 4.2. (a) *The elements m_i , $1 \leq i \leq k$, mutually commute in T_k^a .*

- (b) The eigenvalues of the elements m_i are given by the graph $\hat{T}^{\lambda/\mu}$ of (3.17) in the sense that if the set of vertices on level k is

$$\hat{T}_k^{\lambda/\mu} = \{\mu_1 - \mu_2 + k, \mu_1 - \mu_2 + k - 2, \dots, \mu_1 - \mu_2 - k\} \cap \mathbb{Z}_{\geq 0}, \text{ and}$$

$$\hat{T}_k^{\lambda/\mu} = \{\text{paths } p = (\mu = p^{(0)} \rightarrow p^{(1)} \rightarrow \dots \rightarrow p^{(k)} = \lambda/\mu) \text{ to } \lambda/\mu \text{ in } \hat{T}^{\lambda/\mu}\},$$

for $\lambda/\mu \in \hat{T}_k^{\lambda/\mu}$ then

$$\hat{T}_k^{\lambda/\mu} \text{ is an index set for the simple } T_k^a \text{ modules } T_k^{\lambda/\mu} \text{ appearing in } L(\mu) \otimes V^{\otimes k},$$

and

$$T_k^{\lambda/\mu} \text{ has a basis } \{v_p \mid p \in \hat{T}_k^{\lambda/\mu}\}$$

with

$$m_i v_p = \begin{cases} \pm [p^{(i-1)} + 1] v_p, & \text{if } p^{(i-1)} \pm 1 = p^{(i-2)} = p^{(i)}, \\ 0, & \text{otherwise.} \end{cases}$$

where $p^{(i)}$ is the partition (a single part in this case) on level i of the path p .

- (c) $\kappa = m_k + [2]m_{k-1} + \dots + [k]m_1$ is a central element of T_k^a and κ acts on $T_k^{\lambda/\mu}$ by the constant

$$[k] \frac{q^{-(\mu_1 + \mu_2) - (\mu_1 - \mu_2) + 1}}{q - q^{-1}} + q^{-(\mu_1 + \mu_2)} ([\lambda_1 - \lambda_2 + 2] + [\lambda_1 - \lambda_2 + 4] + \dots + [\mu_1 - \mu_2 + k]).$$

Proof. (a) The elements X^{ε_i} commute with one another in the affine Hecke algebra (see (2.4) and the m_j are by definition linear combinations of the X^{ε_i} (see 2.15), so they commute.

(b) Let p be a path to λ/μ in $\hat{T}^{\lambda/\mu}$ and let T be the corresponding standard tableau on 2 rows. If $p^{(i)} = p^{(i-1)} - 1 = p^{(i-2)} - 2$ or If $p^{(i)} = p^{(i-1)} + 1 = p^{(i-2)} + 2$ then $c(T(i-1)) = c(T(i)) - 1$ and, from (2.15) and Theorem 4.1(b),

$$\begin{aligned} m_i v_T &= q^{i-2} \frac{q^{-2c(T(i))} - q^{-2} q^{-2c(T(i-1))}}{q - q^{-1}} v_T \\ &= q^{i-2} \frac{q^{-2c(T(i))} - q^{-2} q^{-2c(T(i))+2}}{q - q^{-1}} v_T = 0. \end{aligned}$$

If $p^{(i)} = p^{(i-2)} = p^{(i-1)} - 1$ with $T^{(i-1)} = (a, b)$ then $c(T(i)) = a$ and $c(T(i-1)) = b - 2$ and

$$\begin{aligned} m_i v_T &= q^{i-2} \frac{q^{-2a} - q^{-2} q^{-2b+4}}{q - q^{-1}} v_T \\ &= q^i \frac{q^{-(a+b+1)} (q^{-(a-b+1)} - q^{(a-b+1)})}{q - q^{-1}} = -q^{-m} [a - b + 1] v_T, \end{aligned}$$

where $m = |\mu| = a + b - i + 1$. If $p^{(i)} = p^{(i-2)} = p^{(i-1)} + 1$ with $T^{(i-1)} = (a, b)$ then $c(T(i-1)) = a - 1$ and $c(T(i)) = b - 1$ and

$$\begin{aligned} m_i v_T &= q^{i-2} \frac{q^{-2b+2} - q^{-2} q^{-2a+2}}{q - q^{-1}} v_T \\ &= q^i \frac{q^{-(a+b+1)} (q^{(a-b+1)} - q^{-(a-b+1)})}{q - q^{-1}} = q^{-m} [a - b + 1] v_T, \end{aligned}$$

where $m = |\mu| = a + b - i + 1$.

(c) Let $k = |\lambda/\mu|$. The identity

$$q^{\lambda_1 + \lambda_2 - 2} \sum_{b \in \lambda/\mu} q^{-2c(b)} = \left(\sum_{i=\mu_2}^{\lambda_2-1} [\lambda_1 + \lambda_2 - 2i] (q - q^{-1}) \right) + [k] q^{\mu_2 - \mu_1 + 1},$$

is best visible in an example: With $\lambda = (10, 6)$ and $\mu = (4, 2)$,

$$\begin{aligned} & q^{16-2} \begin{pmatrix} 0 & +0 & +0 & +0 & +q^{-8} & +q^{-10} & +q^{-12} & +q^{-14} & +q^{-16} & +q^{-18} \\ +0 & +0 & +q^{-2} & +q^{-4} & +q^{-6} & +q^{-8} & & & & \end{pmatrix} \\ &= \begin{pmatrix} 0 & +0 & +0 & +0 & +q^6 & +q^4 & +q^2 & +q^0 & +q^{-2} & +q^{-4} \\ +0 & +0 & +q^{12} & +q^{10} & +q^8 & +q^6 & & & & \end{pmatrix} \\ &= \begin{pmatrix} 0 & +0 & +0 & +0 & +0 & +0 & +0 & +0 & +0 & +0 \\ +0 & +0 & (q^{12} - q^{-12}) & +(q^{10} - q^{-10}) & +(q^8 - q^{-8}) & +(q^6 - q^{-6}) & +0 & +0 & +0 & +0 \end{pmatrix} \\ & \quad + \begin{pmatrix} 0 & +0 & +0 & +0 & q^{-4} & +q^{-2} & +q^0 & +q^2 & +q^4 & +q^6 \\ +0 & +0 & +q^{-12} & q^{-10} & +q^{-8} & +q^{-6} & & & & \end{pmatrix} \\ &= \left(\sum_{i=2}^{6-1} q^{16-2i} - q^{-(16-2i)} \right) + [10] q^{4-2+1}. \end{aligned}$$

Then Proposition 2.7 says

$$X^{-\varepsilon_1} + \dots + X^{-\varepsilon_k} = q^{-(k-2)} (q - q^{-1}) (m_k + [2]m_{k-1} + \dots + [k]m_1),$$

and so $m_k + [2]m_{k-1} + \cdots + [k]m_1$ acts on $T_k^{\lambda/\mu}$ by the constant

$$\begin{aligned}
(q - q^{-1})^{-1} q^{k-2} \sum_{b \in \lambda/\mu} q^{-2c(b)} &= (q - q^{-1})^{-1} q^{-(\mu_1 + \mu_2)} q^{\lambda_1 + \lambda_2 - 2} \sum_{b \in \lambda/\mu} q^{-2c(b)} \\
&= (q - q^{-1})^{-1} q^{-(\mu_1 + \mu_2)} \left([k] q^{\mu_2 - \mu_1 + 1} + \sum_{i=\mu_2}^{\lambda_2 - 1} [\lambda_1 + \lambda_2 - 2i] (q - q^{-1}) \right) \\
&= [k] \frac{q^{-m - p^{(0)} + 1}}{q - q^{-1}} + \sum_{i=\mu_2}^{\lambda_2 - 1} q^{-m} [m + k - 2i] \\
&= [k] \frac{q^{-m - p^{(0)} + 1}}{q - q^{-1}} + q^{-m} ([p^{(k)} + 2] + [p^{(k)} + 4] + \cdots \\
&\quad \cdots + [p^{(0)} + k - 2] + [p^{(0)} + k]),
\end{aligned}$$

since $\mu_1 + \mu_2 = m$, $\mu_1 - \mu_2 = p^{(0)}$, $\lambda_1 + \lambda_2 = m + k$ and $\lambda_1 - \lambda_2 = p^{(k)}$. \square

REFERENCES

- [Ar] S. Ariki, Representations of quantum algebras and combinatorics of Young tableaux, University Lecture Series **26**, Amer. Math. Soc., Providence, RI, 2002. ISBN: 0-8218-3232-8, MR1911030 (2004b:17022)
- [Ch] I. Cherednik, *A new interpretation of Gel'fand-Tsetlin bases*, Duke Math. J., **54** (1987), no. 2, 563–577.
- [Dr] V. G. Drinfel'd, *Almost cocommutative Hopf algebras*, Leningrad Math. J., **1** (1990), 321–342.
- [Dx] J. Dixmier, Enveloping algebras, Graduate Studies in Mathematics **11**, Amer. Math. Soc., Providence, RI, 1996. ISBN: 0-8218-0560-6, MR1393197 (97c:17010)
- [FH] W. Fulton and J. Harris, Representation theory. A first course, Graduate Texts in Mathematics **129**, Springer-Verlag, New York 1991. ISBN: 0-387-97527-6; 0-387-97495-4, MR1153249 (93a:20069)
- [GHJ] F. Goodman, P. de la Harpe, and V. Jones, Coxeter graphs and towers of algebras, Mathematical Sciences Research Institute Publications **14**, Springer-Verlag, New York, 1989.
- [GL1] J. Graham and G. Lehrer, *Diagram algebras, Hecke algebras and decomposition numbers at roots of unity*, Ann. Scient. Éc. Norm. Sup. 4^e série **36** (2003), no. 4, 479–524. MR2013924 (2004k:20007)
- [GL2] J. Graham and G. Lehrer, *Cellular and diagram algebras in representation theory*, Representation theory of algebraic groups and quantum groups, Adv. Stud. Pure Math. **40**, Math. Soc. Japan, Tokyo, 2004, pp. 141–173. MR2074593 (2005i:20005)
- [Gr] I. Grojnowski, *Affine \mathfrak{sl}_p controls the representation theory of the symmetric group and related Hecke algebras*, [arxiv:math.RT/9907129](https://arxiv.org/abs/math/9907129).

- [Ji] M. Jimbo, *A q -analogue of $U(\mathfrak{gl}(N+1))$, Hecke algebra, and the Yang-Baxter equation*, Lett. Math. Phys., **11** (1986), no. 3, 247–252. MR0841713 (87k:17011)
- [K] A. Kleshchev, *Linear and projective representations of symmetric groups*, Cambridge Tracts in Mathematics **163**, Cambridge University Press, Cambridge, 2005. ISBN: 0-521-83703-0, MR2165457 (2007b:20022)
- [KL] D. Kazhdan and G. Lusztig, *Proof of the Deligne-Langlands conjecture for Hecke algebras*, Invent. Math., **87** (1987), no. 1, 153–215. MR0862716 (88d:11121)
- [LR] R. Leduc and A. Ram, *A ribbon Hopf algebra approach to the irreducible representations of centralizer algebras: the Brauer, Birman-Wenzl, and type A Iwahori-Hecke algebras*, Adv. Math., **125** (1997), 1–94. MR1427801 (98c:20015)
- [Mac] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Second Edition, Oxford University Press, 1995.
- [OR] R. Orellana and A. Ram, *Affine braids, Markov traces and the category O* , Proceedings of the International Colloquium on Algebraic Groups and Homogeneous Spaces, Mumbai 2004, Tata Institute of Fundamental Research (TIFR), Mumbai, Narosa Publishing House, 2007, pp. 423–473.
- [OV] A. Okounkov and A. Vershik, *A new approach to representation theory of symmetric groups*, Selecta Math. (N.S.), **2** (1996), no. 4, 581–605. MR1443185 (99g:20024)
- [R] A. Ram, *Skew shape representations are irreducible*, Combinatorial and Geometric representation theory (S.-J. Kang and K.-H. Lee, eds.), Contemp. Math. **325**, Amer. Math. Soc., 2003, pp. 161–189. MR1988991 (2004f:20014)
- [Re] N. Yu. Reshetikhin, *Quantized universal enveloping algebras, the Yang-Baxter equation, and invariants of links I*, LOMI Preprint E-4-87 (1987).

Tom Halverson
Department of Mathematics
Macalester College
Saint Paul, MN 55105
U.S.A.

halverson@macalester.edu

Manuela Mazzocco
Department of Mathematics
University of Wisconsin
Madison, WI 53706
U.S.A.

Arun Ram

Department of Mathematics and Statistics

University of Melbourne

Parkville VIC 3010

Australia

`aram@unimelb.edu.au`

and

Department of Mathematics

University of Wisconsin

Madison, WI 53706

U.S.A.