

## ON THE COMMUTATIVITY OF THE RADICAL OF THE GROUP ALGEBRA OF A FINITE GROUP

KAORU MOTOSE AND YASUSHI NINOMIYA

(Received December 27, 1978)

Let  $K$  be an algebraically closed field of characteristic  $p > 0$ , and  $G$  a finite group of order  $p^a m$  where  $(p, m) = 1$  and  $a > 0$ . We denote by  $J(KG)$  the radical of the group algebra  $KG$ . In case  $p$  is odd, D.A.R. Wallace [6] proved that  $J(KG)$  is commutative if and only if  $G$  is abelian or  $G/P$  is a Frobenius group with complement  $P$  and kernel  $G'$ , where  $P$  is a Sylow  $p$ -subgroup of  $G$  and  $G'$  the commutator subgroup of  $G$ . On the other hand, in case  $p = 2$ , S. Koshitani [1] has recently given a necessary and sufficient condition for  $J(KG)$  to be commutative. In this paper, we shall give alternative conditions for  $J(KG)$  to be commutative.

If  $J(KG)$  is commutative, then  $G$  is a  $p$ -nilpotent group and a Sylow  $p$ -subgroup of  $G$  is abelian ([6], Theorem 2). We may therefore restrict our attention to a  $p$ -nilpotent group. Now, we put  $N = O_p(G)$ . For a central primitive idempotent  $\varepsilon$  of  $KN$ , we put  $G_\varepsilon = \{g \in G \mid g\varepsilon g^{-1} = \varepsilon\}$ . Let  $a_i$  ( $i = 1, 2, \dots, s$ ) be a complete residue system of  $G(\text{mod } G_\varepsilon)$

$$G = G_\varepsilon a_1 \cup G_\varepsilon a_2 \cup \dots \cup G_\varepsilon a_s.$$

Then K. Morita [2] proved the following:

**Theorem 1.** *If  $G$  is a  $p$ -nilpotent group, then  $e = \sum_{i=1}^s \varepsilon^{a_i}$  is a central primitive idempotent of  $KG$  and  $KGe$  is isomorphic to the matrix ring  $(KP_\varepsilon)_f$  of degree  $f$  over  $KP_\varepsilon$  for some  $f$ , where  $P_\varepsilon$  is a Sylow  $p$ -subgroup of  $G_\varepsilon$ .*

In what follows, for a subset  $S$  of  $G$ , we denote by  $\hat{S}$  the element  $\sum_{x \in S} x$  of  $KG$ . By [5], Theorem, it holds that  $J(KG)^2 = 0$  if and only if  $p^a = 2$ . When this is the case,  $J(KG)$  is trivially commutative. Therefore we may restrict our attention to the case  $p^a \geq 3$ . The following proposition contains [1], Theorem 2.

**Proposition.** *If  $G$  is a non-abelian group and  $p^a \geq 3$ , then the following conditions are equivalent:*

- (1)  $J(KG)$  is commutative.
- (2)  $(G'P)' = G'$  and  $J(KG'P)$  is commutative.
- (3) (i)  $G'$  is a  $p'$ -group, and

(ii) each block of  $KG'P$ , which is not the principal block, is of defect 0 if  $p \neq 2$  and of defect 1 or 0 if  $p = 2$ .

(4) (i)  $G'$  is a  $p'$ -group, and

(ii) for each  $x \in G' - 1$ ,  $C_{G'P}(x)$  is a  $p'$ -group if  $p \neq 2$  and its order is not divisible by 4 if  $p = 2$ .

Proof. (1) $\Rightarrow$ (2): We put  $H = G'P$ . Since  $H$  is a normal subgroup of  $G$ , we have  $J(KH) \subset J(KG)$ . Hence  $J(KH)$  is commutative, and so, by [6], Theorem 2,  $|H'|$  is not divisible by  $p$ . Since  $J(KG)$  is commutative and  $J(KG) \supset J(KH) \supset J(KH'P) \supset \hat{H}'J(KP)$ , by [6], Lemma 3, we have  $\hat{G}'KG \supset J(KG)^2 \supset \hat{H}'^2J(KP)^2 = \hat{H}'J(KP)^2 \ni \hat{H}'\hat{P}$ . Thus, we have  $G' \subset H'P$ . Since  $G'$  is a  $p'$ -group by [6], Theorem 2, we have  $G' = H'$ .

(2) $\Rightarrow$ (3): Since  $J(KG'P)$  is commutative and  $(G'P)' = G'$ ,  $G'$  is a  $p'$ -group by [6], Theorem 2. Now, we put  $e_1 = |G'|^{-1}\hat{G}'$ , and  $e_2 = 1 - e_1$ . Then  $e_1$  and  $e_2$  are central idempotents of  $KG'P$ . Thus we have  $J(KG'P) = e_1J(KG'P) \oplus e_2J(KG'P)$ . Since  $J(KG'P)$  is commutative, by [6], Lemma 3, we have  $J(KG'P)^2 = e_1J(KG'P)^2 \oplus e_2J(KG'P)^2 \subset (\widehat{G'P})'KG'P = \hat{G}'KG'P = e_1KG'P$ . Therefore  $e_2J(KG'P)^2 = 0$ , and so by Theorem 1, every non-simple block of  $e_2KG'P$  is isomorphic to the matrix ring over  $KD$ , where  $K$  is of characteristic 2 and  $D$  is a group of order 2. Hence  $e_2KG'P$  is a direct sum of blocks of defect 0 or of defect 1 or 0 according as  $p$  is odd or 2. Since  $e_1KG'P (= e_1KP)$  is the principal block, we obtain (3).

(3) $\Rightarrow$ (4): This is easy by [3], Theorem 4.

(4) $\Rightarrow$ (3) is trivial.

(3) $\Rightarrow$ (1): Since  $G'P$  is a normal subgroup of  $G$  and  $[G:G'P]$  is not divisible by  $p$ , we have  $J(KG) = J(KG'P)KG$ . We put  $e_1 = |G'|^{-1}\hat{G}'$ , and  $e_2 = 1 - e_1$ . Then  $e_1$  and  $e_2$  are central idempotents of  $KG$  and  $J(KG) = e_1J(KG'P) \cdot KG \oplus e_2J(KG'P)KG$ . Since  $e_1J(KG'P)KG \subset \hat{G}'KG$ ,  $e_1J(KG'P)KG$  is a central ideal of  $KG$  by [4], Lemma 5. By Theorem 1, every block of  $e_2KG'P$  is isomorphic to the matrix ring over  $KD$ , where  $D$  is a  $p$ -group. From our assumption, every non-simple block of  $e_2KG'P$  has the radical of square zero. Hence  $e_2[J(KG'P)KG]^2 = e_2J(KG'P)^2KG = 0$ , and so  $e_2J(KG'P)KG$  is commutative. Thus,  $J(KG)$  is commutative.

REMARK. The condition (4) of Proposition for  $p$  odd is equivalent to the condition of Wallace's result ([6]) that  $G'P$  is a Frobenius group with complement  $P$  and kernel  $G'$ .

Now, in case  $p = 2$ , we shall give the conditions for  $J(KG)$  to be commutative.

**Theorem 2.** Assume that  $p = 2$ ,  $2^a \geq 4$  and  $G' \neq 1$ . Then the following conditions are equivalent:

- (1)  $J(KG)$  is commutative.
- (2)  $G'$  is of odd order and  $|P \cap P^h| \leq 2$  for every  $h \in G'P - P$ .
- (3)  $G'$  is of odd order and  $C_{G'P}(s)/\langle s \rangle$  is either a 2-group or a Frobenius group with complement  $P/\langle s \rangle$  for every involution  $s$  of  $P$ .

Proof. (1) $\Rightarrow$ (2): Suppose that  $J(KG)$  is commutative. Then, by Proposition,  $G'$  is of odd order. Let  $h$  be an arbitrary element of  $G'P - P$ , and  $x$  an arbitrary element of  $P \cap P^h$ . Then  $hxh^{-1}x^{-1} \in P \cap G' = 1$ , and so  $x \in C_{G'P}(h)$ . Thus,  $P \cap P^h \subset C_{G'P}(h)$ . Since we may assume that  $h \in G' - 1$ , we obtain  $|P \cap P^h| \leq 2$  by Proposition.

(2) $\Rightarrow$ (3): Let  $s$  be an arbitrary involution of  $P$  such that  $C_{G'P}(s) \neq P$ . Then  $P \cap P^x = \langle s \rangle$  for  $x \in C_{G'P}(s) - P$ , and so  $C_{G'P}(s)/\langle s \rangle$  is a Frobenius group with complement  $P/\langle s \rangle$ .

(3) $\Rightarrow$ (1): Let  $x$  be an element of  $G' - 1$ , and  $S$  a Sylow 2-subgroup of  $C_{G'P}(x)$ . Suppose that  $S \neq 1$ . Then  $S \subset P^u$  for some  $u \in G'P$ , and  $x \in C_{G'P}(S) \subset C_{G'P}(s)$  for every involution  $s$  of  $S$ . Hence,  $C_{G'P}(s)$  is not a 2-group, and so  $C_{G'P}(s)/\langle s \rangle$  is a Frobenius group with complement  $P^u/\langle s \rangle$ . Thus, we have  $S \subset P^u \cap P^{u^x} = \langle s \rangle$ , and hence  $|C_{G'P}(x)|$  is not divisible by 4, which implies (1) by Proposition.

**Corollary.** Assume that  $p=2$ ,  $2^a \geq 4$  and  $G' \neq 1$ . If  $J(KG)$  is commutative, then a Sylow 2-subgroup of  $G$  is a cyclic group or an abelian group of type  $(2, 2^{a-1})$ .

Proof. Suppose that  $J(KG)$  is commutative. Then, by Theorem 2,  $|P \cap P^h| \leq 2$  for every  $h \in G'P - P$ . If  $P \cap P^h = 1$  for all  $h \in G'P - P$ , then  $G'P$  is a Frobenius group with complement  $P$  and kernel  $G'$ . Hence  $P$  is cyclic. On the other hand, if  $P \cap P^h = \langle s \rangle$  for some  $h \in G'P - P$  and some involution  $s$  of  $P$ , then  $hsh^{-1}s^{-1} \in P \cap G' = 1$ , and so  $h \in C_{G'P}(s)$  and  $h \notin P$ . Therefore  $C_{G'P}(s)$  properly contains  $P$ . Hence,  $C_{G'P}(s)/\langle s \rangle$  is a Frobenius group with complement  $P/\langle s \rangle$  by the condition (3) of Theorem 2. Hence  $P/\langle s \rangle$  is cyclic, and so  $P$  is a cyclic group or an abelian group of type  $(2, 2^{a-1})$ .

REMARK. In case  $G$  is a non-abelian group and  $p^a \geq 3$ , S. Koshitani [1] proved that if  $J(KG)$  is commutative, then  $N_G(P)$  is abelian. This is included in the following proposition: Let  $G$  be a non-abelian group, and  $p^a \geq 3$ . If  $J(KG)$  is commutative then  $G$  is a semi-direct product of  $G'$  by (abelian)  $N_G(P)$ .

Proof. It is easy to see  $G = G'N_G(P)$ . Suppose that  $J(KG)$  is commutative. Let  $x$  be a  $p'$ -element of  $N_{G'P}(P)$ . Since  $G'P$  is a  $p$ -nilpotent group,  $N_{G'P}(P)$  is the direct product of  $P$  and a normal  $p'$ -subgroup, and so  $C_{G'P}(x)$  contains  $P$ . Hence, by Proposition (4), we have  $x=1$ , which implies that  $G' \cap N_G(P) = 1$ .

**References**

- [1] S. Koshitani: *Remarks on the commutativity of the radicals of group algebras*, Glasgow Math. J. **20** (1979), 63–68.
- [2] K. Morita: *On group rings over a modular field which possess radicals expressible as principal ideals*, Sci. Rep. Tokyo Bunrika Daigaku **A 4** (1951), 177–194.
- [3] M. Osima: *On primary decomposable group rings*, Proc. Phys.-Math. Soc. Japan **24** (1942), 1–9.
- [4] D.A.R. Wallace: *Group algebras with central radicals*, Proc. Glasgow Math. Assoc. **5** (1962), 103–108.
- [5] ———: *Group algebras with radicals of square zero*, *ibid.* **5** (1962), 158–159.
- [6] ———: *On the commutativity of the radical of a group algebra*, *ibid.* **7** (1965), 1–8.

Department of Mathematics  
Shinshu University  
Matsumoto 390, Japan