

THE REFLECTIVITY OF SOME CATEGORIES OF T_0 SPACES IN DOMAIN THEORY

CHONG SHEN, XIAOYONG XI, AND DONGSHENG ZHAO

ABSTRACT. Keimel and Lawson proposed a set of conditions for proving the reflectivity of a category of topological spaces in the category of all T_0 spaces. Recently, these conditions were used to prove the reflectivity of the category of all well-filtered spaces. In this paper, we prove that, in certain sense, these conditions are not only sufficient but also necessary for a category of T_0 spaces to be reflective. By applying this general result, we can easily deduce that several categories proposed in domain theory are not reflective, thereby answering a few open problems.

1. Introduction

Given a full subcategory \mathbf{D} of a category \mathbf{C} , one natural and frequently asked question is whether \mathbf{D} is reflective in \mathbf{C} . The objects in \mathbf{D} can be viewed as “special objects”, the reflectivity of \mathbf{D} ensures that every general object in \mathbf{C} can be “completed” to be a special object, or “densely embedded into” a special object.

Keimel and Lawson [12] proved that a full subcategory \mathbf{K} of \mathbf{Top}_0 of all T_0 spaces is reflective if it satisfies the following four conditions:

(K1) \mathbf{K} contains all sober spaces.

(K2) If $X \in \mathbf{K}$ and Y is homeomorphic to X , then $Y \in \mathbf{K}$.

(K3) If $\{X_i : i \in I\} \subseteq \mathbf{K}$ is a family of subspaces of a sober space, then the subspace $\bigcap_{i \in I} X_i \in \mathbf{K}$.

(K4) If $f : X \rightarrow Y$ is a continuous mapping from a sober space X to a sober space Y , then for any subspace Y_1 of Y , $Y_1 \in \mathbf{K}$ implies that $f^{-1}(Y_1) \in \mathbf{K}$.

It has been proved that the categories of d -spaces, well-filtered spaces and sober spaces all satisfy the aforementioned four conditions, as shown in [12, 22, 23]. Therefore, they are all reflective subcategories of \mathbf{Top}_0 .

For a full subcategory \mathbf{K} of \mathbf{Top}_0 , we say that \mathbf{K}

(1) is *productive*, if the product $\prod_{i \in I} X_i \in \mathbf{K}$ whenever $\{X_i : i \in I\} \subseteq \mathbf{K}$, and

(2) is *b-closed-hereditary*, if $Y \in \mathbf{K}$ whenever Y is a b -closed subspace of some $X \in \mathbf{K}$.

The four conditions (K1)–(K4) can, however, usually only be used to confirm the reflectivity of subcategories of \mathbf{Top}_0 . In this paper we shall prove that, in certain sense, they are also necessary conditions, and can therefore be used to disprove the reflectivity of some subcategories of \mathbf{Top}_0 . In

This work was supported by the National Natural Science Foundation of China (1210010153, 12071188), Jiangsu Provincial Department of Education (21KJB110008), and International Postdoctoral Exchange Fellowship Program (No. PC2022013).

2020 *Mathematics Subject Classification*. 06B30, 06B35, 54A05, 18F60.

Key words and phrases. b -topology, consonant space, co-sober space, k -bounded sober spaces, reflective subcategory, sober space.

1 particular, we shall use this result to solve several open problems that were posed in [24]. Our main
2 results are as follows.

3 **Theorem A.** For a full subcategory \mathbf{K} of \mathbf{Top}_0 with $\mathbf{K} \not\subseteq \mathbf{Top}_1$, if \mathbf{K} satisfies (K2), then the following
4 four statements are equivalent:

- 5 (1) \mathbf{K} is reflective in \mathbf{Top}_0 ;
6 (2) \mathbf{K} satisfies conditions (K1)–(K4);
7 (3) \mathbf{K} is productive and b -closed-hereditary;
8 (4) \mathbf{K} is productive and has equalizers.
9

10 **Theorem B.** The categories of co-sober spaces, strongly k -bounded sober spaces, strongly d -spaces,
11 and consonant T_0 spaces, are not reflective in \mathbf{Top}_0 .
12

13 2. Preliminaries

14
15 Let P be a poset. For any $A \subseteq P$, let $\downarrow A = \{x \in P : x \leq a \text{ for some } a \in A\}$ and $\uparrow A = \{x \in P : x \geq$
16 $a \text{ for some } a \in A\}$. For $x \in P$, we write $\downarrow x$ for $\downarrow\{x\}$ and $\uparrow x$ for $\uparrow\{x\}$, respectively. A subset A of P is
17 called a *lower set* (resp. *upper set*) if $A = \downarrow A$ (resp. $A = \uparrow A$).

18 For a T_0 space X , the specialization order \leq on X is defined as $x \leq y$ iff $x \in \text{cl}(\{y\})$, where cl is
19 the closure operator on X . In the following, when we consider a T_0 space X as a poset, it is always
20 equipped with the specialization order.

21 For a T_0 space X , we use $\mathcal{O}(X)$ to denote the topology of X . For any subset A of X , the *saturation*
22 of A , denoted by $\text{Sat}(A)$, is defined to be

$$23 \text{Sat}(A) = \bigcap \{U \in \mathcal{O}(X) : A \subseteq U\}.$$

24
25 A subset A of a T_0 space X is *saturated* if $A = \text{Sat}(A)$.

26 **Remark 2.1** ([6, 7]). Let X be a T_0 space.

- 27 (1) For any subset A of X , $\text{Sat}(A) = \uparrow A$.
28 (2) For any $x \in X$, $\downarrow x = \text{cl}(\{x\})$, and $x \in \text{Sat}(A)$ if and only if $\downarrow x \cap A \neq \emptyset$.
29 (3) For any open subset U of X , $U = \uparrow U$, and for any closed subset F of X , $F = \downarrow F$.
30

31 A nonempty subset A of a T_0 space is called *irreducible* if for any closed sets F_1, F_2 , $A \subseteq F_1 \cup F_2$
32 implies $A \subseteq F_1$ or $A \subseteq F_2$. A T_0 space X is called *sober* if for any irreducible closed set F of X there is
33 a (unique) point $x \in X$ such that $F = \text{cl}(\{x\})$.

34 A very effective tool for studying sober spaces is the b -topology introduced by L. Skula [17] (see
35 also [3]).
36

37 **Definition 2.2** ([3, 17]). Let X be a T_0 space. The b -topology (also called *Skula topology* [17] or *strong*
38 *topology* [6, Exercise V-5.31]) associated with X is the topology having

$$39 \{U \cap \downarrow x : x \in U \in \mathcal{O}(X)\}$$

40
41 as a base. The resulting space will be denoted by bX . A subset B of X is *b -dense* in X , if it is dense in
42 X with respect to the b -topology.

1 The following properties on b -topology will be used later. For further information, one can refer to
 2 [10, 12] and Exercise V-5.31 in [6].

3 **Remark 2.3.** Let X be a T_0 space.

4 (1) The b -topology on X is finer than the original topology on X . This follows trivially from the fact
 5 that for any open set U in X , we have $U = \bigcup_{x \in U} U \cap \downarrow x$.

6 (2) Let X be a T_0 space. For each $x \in X$, we have that $\downarrow x = X \cap \downarrow x$, so $\downarrow x$ is b -open, and it is also
 7 b -closed since $X \setminus \downarrow x$ is b -open. Thus, the b -topology of X is always Hausdorff.

8 (3) Every saturated set A in a T_0 space X is b -closed. In fact, we have that
 9

$$10 \quad X \setminus A = \downarrow(X \setminus A) = \bigcup_{x \in X \setminus A} \downarrow x,$$

11 which is b -open by (2). Therefore, $A = \uparrow A$ is b -closed.

12 (4) For each b -closed set E of X , $E = \bigcap_{i \in I} U_i \cup (X \setminus V_i)$, where $U_i, V_i \in \mathcal{O}(X)$ for any $i \in I$. In fact, since
 13 $X \setminus E$ is b -open, for each $x \notin E$, there exists an open neighborhood V_x of x such that $V_x \cap \downarrow x \subseteq X \setminus E$,
 14 which implies that $X \setminus E = \bigcup_{x \notin E} V_x \cap \downarrow x$; thus $E = \bigcap_{x \notin E} (X \setminus \downarrow x) \cap (X \setminus V_x)$, completing the proof.

15 **Definition 2.4.** (1) A space X is a *retract* of space Y , if there exist two continuous maps $s : X \rightarrow Y$
 16 (the *section*) and $r : Y \rightarrow X$ (the *retraction*) such that $r \circ s = \text{id}_X$, the identity mapping on X [7].

17 (2) We call X a *b-retract* of Y if X is a retraction of Y such that $s(X)$ is b -dense in Y .

18 **Remark 2.5** ([7]). Every section $s : X \rightarrow Y$ is an embedding and every retraction $r : Y \rightarrow X$ is a
 19 quotient mapping.

20 **Proposition 2.6** ([17, 2.6], [19, Proposition 2.11]). *If X and Y are T_0 spaces and X is a b -retract of Y ,*
 21 *then X is homeomorphic to Y .*

22 In what follows, we shall denote by **Top**₀ (resp. **Top**₁, **Sob**) the category of all T_0 spaces (resp. T_1
 23 spaces, sober spaces) with continuous mappings as morphisms. All subcategories of **Top**₀ are assumed
 24 to be full and closed under the formation of homeomorphic objects (i.e., satisfy (K2)).

25 **Definition 2.7** ([15]). A full subcategory **K** of **Top**₀ is *reflective* if, for each $X \in \mathbf{Top}_0$, there exists
 26 $X^k \in \mathbf{K}$ (the **K**-completion for X) and a continuous mapping $\mu_X : X \rightarrow X^k$ (the **K**-reflection for X)
 27 satisfying the universal property: for any continuous mapping $f : X \rightarrow Z$ to a space $Z \in \mathbf{K}$, there
 28 exists a unique continuous mapping $g : X^k \rightarrow Z$ such that $g \circ \mu_X = f$:

$$29 \quad \begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow f & \downarrow g \\ & & Z \end{array}$$

30 Equivalently, **K** is reflective if the inclusion functor $I : \mathbf{K} \rightarrow \mathbf{Top}_0$ has a left adjoint (see IV-3 in
 31 [15]). The category **Sob** is a full reflective subcategory of **Top**₀. The **Sob**-completion for X is usually
 32 called the *sobrification* of X .

33 The following lemma can be easily verified by using the definition of **K**-reflection.

34 **Lemma 2.8.** *Let $\mu_1 : X \rightarrow Y_1$ be a **K**-reflection for X . Then, the following conditions are equivalent:*

1 (1) $\mu_2 : X \longrightarrow Y_2$ is a \mathbf{K} -reflection;

2 (2) there exists a (unique) homeomorphism $h : Y_1 \longrightarrow Y_2$ such that $h \circ \mu_1 = \mu_2$.

3 **Definition 2.9.** A mapping $e : X \longrightarrow Y$ between topological spaces is called a *b-dense embedding*, if it
4 is a topological embedding such that $e(X)$ is *b-dense* in Y .

5
6 **Theorem 2.10** ([12, Proposition 3.4, Corollary 3.5]). Let X be a sober space and $Y \subseteq X$.

7 (1) The subspace Y is sober if and only if Y is *b-closed*.

8 (2) The inclusion mapping $e : Y \longrightarrow Y^s$, $x \mapsto x$, is a sober reflection for Y , where Y^s is the *b-closure* of
9 Y in X .

10 **Theorem 2.11** ([12, Proposition 3.2]). Let X be a T_0 space, Y a sober space and $f : X \longrightarrow Y$ a
11 continuous mapping. Then, f is a sober reflection for X if and only if it is a *b-dense embedding*.

12
13 **Theorem 2.12** ([19, Theorem 3.2]). Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$.
14 Then, each \mathbf{K} -reflection is a *b-dense embedding*.

15 16 3. Main results

17 In this section, we present the main results, starting with a simple yet useful topological space in
18 domain theory.

19
20 **Definition 3.1** ([6, 7]). The *Sierpiński space* is the Scott space $\Sigma 2$, where the underlying set 2 is the
21 two-element chain $2 = \{0, 1\}$ with the order defined by $0 \leq 1$. Note that the open sets in this space are
22 \emptyset , $\{0, 1\}$, and $\{1\}$.

23 **Remark 3.2** ([6, 7]). (1) For any set M , $(\Sigma 2)^M = \Sigma(2^M, \subseteq)$.

24 (2) Let X be a T_0 space and $M = \mathcal{O}(X)$. Then, the mapping $e : X \longrightarrow (\Sigma 2)^M$, $x \mapsto (\chi_U(x))_{U \in M}$, is an
25 embedding. Hence, by Theorem 2.10, X is a sober space iff $e(X)$ is a *b-closed* subset of $(\Sigma 2)^M$.

26
27 **Lemma 3.3.** Let X be a T_0 space. Then, the following statements are equivalent:

28 (1) X is non- T_1 ;

29 (2) $\Sigma 2$ is a retract of X ;

30 (3) $\Sigma 2$ is homeomorphic to a *b-closed* subspace of X ;

31 (4) $\Sigma 2$ is homeomorphic to a subspace of X .

32
33 **Proof.** (1) \Rightarrow (2): Suppose X is non- T_1 . Then, there exist $x_0, x_1 \in X$ such that $x_0 < x_1$. We define two
34 mappings $s : \Sigma 2 \longrightarrow X$ by $s(0) = x_0$ and $s(1) = x_1$, and $r : X \longrightarrow \Sigma 2$ by

$$35 \quad r(x) = \begin{cases} 0, & x \leq x_0; \\ 1, & \text{else,} \end{cases}$$

36
37 for any $x \in X$. It is trivial to verify that both r and s are continuous mappings such that $r \circ s = \text{id}_{\Sigma 2}$,
38 where $\text{id}_{\Sigma 2}$ is the identity mapping on $\Sigma 2$. Therefore, $\Sigma 2$ is a retract of X .

39
40 (2) \Rightarrow (3): If $\Sigma 2$ is a retract of X , then by Remark 2.5, $\Sigma 2$ is homeomorphic to a subspace $\{x_1, x_2\}$
41 of X . In addition, by Remark 2.3(2), we know that bX is Hausdorff; thus $\{x_1, x_2\}$ is *b-closed*.

42 (3) \Rightarrow (4): It is clear.

1 (4) \Rightarrow (1): Note that the T_1 -separation property is hereditary. Then, since $\Sigma 2$ is a non- T_1 subspace of
 2 X up to homeomorphism, it follows that X is also a non- T_1 space. \square

3
 4 As an immediate result of Lemma 3.3, the following corollary is clear.

5 **Corollary 3.4.** *Let \mathbf{K} be a full subcategory of \mathbf{Top}_0 . Then, the following statements are equivalent:*

6 (1) $\mathbf{K} \not\subseteq \mathbf{Top}_1$;

7 (2) *The space $\Sigma 2$ can be topologically embedded into some space Y that belongs to \mathbf{K} .*

8
 9 The following lemma extends Result 2.5 in [17].

10 **Lemma 3.5.** *Let $X, Y, Z \in \mathbf{Top}_0$, $k : X \rightarrow Y$ be a continuous mapping such that $k(X)$ is b -dense in Y ,
 11 and $f : X \rightarrow Z$ a continuous mapping.*

12 (1) *There exists at most one continuous mapping $g : Y \rightarrow Z$ such that $f = g \circ k$.*

13 (2) *If $g : Y \rightarrow Z$ is a continuous mapping such that $f = g \circ k$, then $g(Y) \subseteq \text{cl}_b(f(X))$, where $\text{cl}_b(f(X))$
 14 is the b -closure of $f(X)$ in Z .*

15
 16 **Proof.** (1) Suppose that there exist two continuous mappings $g_1, g_2 : Y \rightarrow Z$ such that $g_1 \circ k = g_2 \circ k =$
 17 f :

$$\begin{array}{ccc} X & \xrightarrow{k} & Y \\ & \searrow f & \downarrow g_1, g_2 \\ & & Z \end{array}$$

18
 19
 20
 21
 22
 23 Let $y \in Y$. Suppose $V \in \mathcal{O}(Z)$ such that $g_1(y) \in V$. Then $y \in g_1^{-1}(V) \in \mathcal{O}(Y)$. Since $k(X)$ is b -dense
 24 in Y and $g_1^{-1}(V) \cap \downarrow y$ is b -open, $k(X) \cap g_1^{-1}(V) \cap \downarrow y \neq \emptyset$. In addition, since $g_1 \circ k = g_2 \circ k = f$, we
 25 deduce that $k(X) \cap g_1^{-1}(V) = k(X) \cap g_2^{-1}(V) \subseteq g_2^{-1}(V)$. It follows that $g_2^{-1}(V) \cap \downarrow y \neq \emptyset$, which implies
 26 that $y \in g_2^{-1}(V)$, i.e., $g_2(y) \in V$. These show that each open neighborhood of $g_1(y)$ contains $g_2(y)$;
 27 thus $g_1(y) \in \text{cl}(\{g_2(y)\})$. Dually, it holds that $g_2(y) \in \text{cl}(\{g_1(y)\})$. Since Z is a T_0 space, we have that
 28 $g_1(y) = g_2(y)$. Therefore, $g_1 = g_2$.

29
 30 (2) Let $y \in Y$ and $V \in \mathcal{O}(Z)$ such that $g(y) \in V$. Then $y \in g^{-1}(V) \in \mathcal{O}(Y)$, and since $k(X)$ is b -dense
 31 in Y , $k(X) \cap g^{-1}(V) \cap \downarrow y \neq \emptyset$. Then there exists $x_0 \in X$ such that $k(x_0) \in g^{-1}(V) \cap \downarrow y$, which implies
 32 that $g(y) \geq g(k(x_0)) \in V$ (note that g is monotone since it is continuous); thus $f(x_0) = g(k(x_0)) \in$
 33 $f(X) \cap V \cap \downarrow g(y) \neq \emptyset$. This shows that $g(y) \in \text{cl}_b(f(X))$. Hence, $g(Y) \subseteq \text{cl}_b(f(X))$. \square

34
 35 **Theorem 3.6.** *Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. Then, the following
 36 statements hold.*

37 (1) \mathbf{K} is b -closed-hereditary.

38 (2) *The Sierpiński space $\Sigma 2 \in \mathbf{K}$. Hence, for any set M , the product $(\Sigma 2)^M \in \mathbf{K}$.*

39 (3) $\mathbf{Sob} \subseteq \mathbf{K}$.

40
 41 **Proof.** (1) Let $X \in \mathbf{K}$, A be a b -closed subspace of X , and $\mu_A : A \rightarrow A^k$ be the \mathbf{K} -reflection for A .
 42 Then, $\mu_A(A)$ is a b -dense subset of A^k by Theorem 2.12. Consider the inclusion mapping $e : A \rightarrow X$,

1 $x \mapsto x$. Then there exists a unique continuous mapping $f : A^k \rightarrow X$ such that $f \circ \mu_A = e$:



7 Then by Lemma 3.5, we have $f(A^k) \subseteq \text{cl}_b(e(A)) = A$, which shows that A is a b -dense retract of A^k .

8 By Proposition 2.6, A is homeomorphic to A^k , and since $A^k \in \mathbf{K}$, it follows that $A \in \mathbf{K}$.

9 (2) Since $\mathbf{K} \not\subseteq \mathbf{Top}_1$, there exists a T_0 and non- T_1 space $X \in \mathbf{K}$. By Lemma 3.3, $\Sigma 2$ is a b -closed
10 subspace of X up to homeomorphism, and from result (1) it follows that $\Sigma 2 \in \mathbf{K}$. Since \mathbf{K} is reflective,
11 \mathbf{K} is productive (see V-6 in [15]), hence $(\Sigma 2)^M \in \mathbf{K}$.

12 (3) Let $X \in \mathbf{Sob}$. By Remark 3.2, there is an embedding $e : X \rightarrow (\Sigma 2)^M$ such that $e(X)$ is a b -closed
13 subspace of $(\Sigma 2)^M$, where $M = \mathcal{O}(X)$. By (2), $(\Sigma 2)^M \in \mathbf{K}$ and since \mathbf{K} is b -closed-hereditary, we have
14 that $e(X) \in \mathbf{K}$, and since X is homeomorphic to $e(X)$, it follows that $X \in \mathbf{K}$. Hence, $\mathbf{Sob} \subseteq \mathbf{K}$. \square

15 Note that every saturated subset of a T_0 space is b -closed by Remark 2.3(3). Thus, the following
16 corollary follows directly from Theorem 3.6(1).

17 **Corollary 3.7.** *Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. If $X \in \mathbf{K}$ and Y is a
18 saturated subspace of X , then Y belongs to \mathbf{K} .*

19 Since there exist sober but non- T_1 spaces (such as $\Sigma 2$), the following corollary follows directly from
20 Theorem 3.6(3).

21 **Corollary 3.8.** *Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 . Then, $\mathbf{K} \not\subseteq \mathbf{Top}_1$ if and only if $\mathbf{Sob} \subseteq \mathbf{K}$.*

22 Recall that the reflective hull of a subcategory \mathbf{C} of \mathbf{Top}_0 is the smallest reflective subcategory of
23 \mathbf{Top}_0 containing \mathbf{C} . Let \mathbf{Sier} be the full subcategory of \mathbf{Top}_0 consisting of all T_0 spaces X which are
24 homeomorphic to $\Sigma 2$.

25 **Corollary 3.9** ([16, Theorem 3.4]). *The reflective hull of \mathbf{Sier} in \mathbf{Top}_0 is \mathbf{Sob} .*

26 **Proof.** Suppose \mathbf{K} is a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{Sier} \subseteq \mathbf{K}$. Note that $\Sigma 2$ is a T_0 but
27 non- T_1 space; thus $\mathbf{K} \not\subseteq \mathbf{Top}_1$. By Theorem 3.6(3), $\mathbf{Sob} \subseteq \mathbf{K}$. Since \mathbf{Sob} is reflective, it is the smallest
28 reflective subcategory of \mathbf{Top}_0 having \mathbf{Sier} as a subcategory. Therefore, \mathbf{Sob} is the reflective hull of
29 \mathbf{Sier} in \mathbf{Top}_0 . \square

30 **Lemma 3.10** ([13, Lemma 5, pp.116]). *If $\{f_i : X \rightarrow Y_i\}_{i \in I}$ is a family of continuous mappings between
31 T_0 spaces, then the diagonal $\Delta_{i \in I} f_i : X \rightarrow \prod_{i \in I} Y_i$ is a continuous mapping, where*

32
33

$$\forall x \in X, (\Delta_{i \in I} f_i)(x) = (f_i(x))_{i \in I}.$$

34 A *skeleton* of a category \mathbf{C} is a full subcategory, denoted by \mathbf{skC} , such that each object of \mathbf{C} is
35 isomorphic to exactly one object of \mathbf{skC} .

36 **Remark 3.11.** Some properties on the skeleton are listed below (see [1, Proposition 4.14, pp. 51]):

37 (1) Every category has a skeleton.
38
39
40
41
42

1 (2) Any two skeletons of a category are isomorphic.

2 A category is a *small category* if its class of objects is a set.

3
4 **Lemma 3.12.** For any cardinal number α , let \mathbf{T}_α be the full subcategory of \mathbf{Top}_0 consisting of all T_0
5 spaces whose cardinality is less than or equal to α . Then, every skeleton of \mathbf{T}_α is a small category.

6
7 **Proof.** Let \mathbf{skT}_α be the full subcategory of \mathbf{T}_α consisting of all T_0 spaces of the form (β, \mathcal{T}) , where
8 β is a cardinal number such that $\beta \leq \alpha$, and \mathcal{T} is an arbitrary T_0 topology on β . Then, it is clear that
9 \mathbf{skT}_α is a skeleton of \mathbf{T}_α , and $|\mathbf{skT}_\alpha| \leq |\bigcup_{\beta \leq \alpha} 2^\beta|$, where $|\mathbf{skT}_\alpha|$ is the cardinality of the class of all
10 objects of \mathbf{skT}_α . Thus, the class of objects of \mathbf{skT}_α is a set. Therefore, \mathbf{skT}_α is a small category. \square

11 **Theorem 3.13.** Let \mathbf{K} be a full subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. Then, the following statements
12 are equivalent:

13 (1) \mathbf{K} is reflective;

14 (2) \mathbf{K} is productive and *b-closed-hereditary*.

15
16 **Proof.** (1) \Rightarrow (2): It is well-known that if \mathbf{K} is reflective, then it is productive (see V-6 in [15]), and by
17 Theorem 3.6, it is *b-closed-hereditary*.

18 (2) \Rightarrow (1): Let $X \in \mathbf{Top}_0$. We will complete the proof in a few steps.

19 *Step 1:* We define the full subcategory $\mathbf{C}(X)$ of \mathbf{K} to consist of all objects Y such that there exists a
20 continuous mapping $f : X \rightarrow Y$ with the property that $f(X)$ is *b-dense* in Y . Then, for each $Y \in \mathbf{C}(X)$,
21 the sobrification $f(X)^s$ of $f(X)$ and the sobrification Y^s of Y are homeomorphic (see [12, Proposition
22 3.4]), which implies that

$$23 \quad |Y| \leq |Y^s| = |f(X)^s| = |\text{Irr}(f(X))| \leq 2^{|f(X)|},$$

24
25 where $\text{Irr}(f(X))$ is the set of all irreducible closed sets in the subspace $f(X)$ of Y . Note that $|f(X)| \leq$
26 $|X|$ (because f is a mapping), so $|Y| \leq 2^{|X|}$.

27 Let $\mathbf{skC}(X)$ be a skeleton of $\mathbf{C}(X)$. Since the cardinality of each space in $\mathbf{skC}(X)$ is less than or
28 equal to $2^{|X|}$, by Lemma 3.12, $\mathbf{skC}(X)$ is a small category, so there is a cardinal number α such that
29 $|\mathbf{skC}(X)| \leq \alpha$.

30
31 *Step 2:* Denote by $\Phi(X)$ the family of all pairs (Y, f) , where $Y \in \mathbf{skC}(X)$ and $f : X \rightarrow Y$ is a
32 continuous mapping such that $f(X)$ is *b-dense* in Y . For each $Y \in \mathbf{skC}(X)$, since the cardinality of the
33 set of all continuous mappings from X to Y is less than or equal to $|Y|^{|X|}$, we have that

$$34 \quad |\Phi(X)| \leq \left| \bigcup_{Y \in \mathbf{skC}(X)} |Y|^{|X|} \right|.$$


35
36 Thus, $\Phi(X)$ is a set, and then we may assume that $\Phi(X) = \{(Y_i, f_i) : i \in I\}$, where I is a set. Therefore,
37 for each $i \in I$, $Y_i \in \mathbf{K}$ and $f_i : X \rightarrow Y_i$ is a continuous mapping such that $f_i(X)$ is *b-dense* in Y_i .

38
39 *Step 3:* Let $X^k = \text{cl}_b((\Delta_{i \in I} f_i)(X))$ be the *b-closure* of $(\Delta_{i \in I} f_i)(X)$ in the product space $\prod_{i \in I} Y_i$, where
40 $\Delta_{i \in I} f_i : X \rightarrow \prod_{i \in I} Y_i$ is the diagonal (i.e., $x \mapsto (f_i(x))_{i \in I}$). Since $\{Y_i : i \in I\} \subseteq \mathbf{K}$ and \mathbf{K} is productive,
41 $\prod_{i \in I} Y_i \in \mathbf{K}$, and since \mathbf{K} is *b-closed-hereditary*, $X^k \in \mathbf{K}$. Let $k : X \rightarrow X^k$ be the restriction of the


1 diagonal $\Delta_{i \in I} f_i$, that is, $k(x) = (f_i(x))_{i \in I}$ for each $x \in X$. It is clear that k is continuous, and since
 2 $X^k = \text{cl}_b((\Delta_{i \in I} f_i)(X))$, it follows that $k(X) = (\Delta_{i \in I} f_i)(X)$ is b -dense in X^k .

3 *Step 4:* Now, we prove that the subspace X^k of $\prod_{i \in I} Y_i$ with the mapping k is the \mathbf{K} -reflection for X .
 4 To see this, suppose $Y \in \mathbf{K}$ and $f : X \rightarrow Y$ is a continuous mapping. We consider the following two
 5 cases:

6
 7 (c1) $f(X)$ is b -dense in Y . Then, $Y \in \mathbf{C}(X)$, and there is a homeomorphism h from Y to a unique
 8 space Z in $\mathbf{skC}(X)$. It is trivial to check that $h \circ f : X \rightarrow Z$ is a continuous mapping such that
 9 $h(f(X))$ is b -dense in Z , so $(Z, h \circ f) \in \Phi(X)$. Assume $(Z, h \circ f) = (Y_j, f_j)$ for some $j \in I$. Let
 10 $p_j : X^k \rightarrow Y_j$ be the restriction of the projection from $\prod_{i \in I} Y_i$ to Y_j (i.e., $(x_i)_{i \in I} \mapsto x_j$). Then p_j is
 11 a continuous mapping, and clearly $p_j \circ k = f_j$:




12
 13
 14
 15
 16
 17
 18
 19
 20 Let $\hat{f} = h^{-1} \circ p_j$. Then $\hat{f} : X^k \rightarrow Y$ is a continuous mapping such that $\hat{f} \circ k = (h^{-1} \circ p_j) \circ k =$
 21 $h^{-1} \circ (p_j \circ k) = h^{-1} \circ f_j = h^{-1} \circ (h \circ f) = (h^{-1} \circ h) \circ f = f$:
 22




23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36 Recall that $k : X \rightarrow X^k$ is a continuous mapping such that $k(X)$ is b -dense in X^k . Then, by
 37 Lemma 3.5, \hat{f} is the unique continuous mapping such that $\hat{f} \circ k = f$.

38
 39 (c2) $f(X)$ is not b -dense in Y . Let $\text{cl}_b(f(X))$ be the b -closure of $f(X)$ in Y with the relative topology.
 40 Then the co-restriction $f^* : X \rightarrow \text{cl}_b(f(X))$ of f (i.e., $\forall x \in X, f^*(x) = f(x)$) is a continuous
 41 mapping such that $f^*(X)$ is b -dense in $\text{cl}_b(f(X))$. Since $Y \in \mathbf{K}$ and \mathbf{K} is b -closed-hereditary, it
 42 follows that $\text{cl}_b(f(X)) \in \mathbf{K}$. Then using the argument of (c1), there is a continuous mapping

1 $g : X^k \rightarrow \text{cl}_b(f(X))$ such that $g \circ k = f^*$:



8 Let $e : \text{cl}_b(f(X)) \rightarrow Y$ be the inclusion mapping. Then, $e \circ f^* = f$. Let $\hat{f} = e \circ g$. Then
9 $\hat{f} : X^k \rightarrow Y$ is a continuous mapping such that $\hat{f} \circ k = (e \circ g) \circ k = e \circ (g \circ k) = e \circ f^* = f$:



19 The uniqueness of \hat{f} follows from Lemma 3.5.

20 All these show that $k : X \rightarrow X^k$ is a \mathbf{K} -reflection for X . Therefore, \mathbf{K} is a reflective subcategory of
21 \mathbf{Top}_0 . □

22 **Definition 3.14.** We say that a full subcategory \mathbf{K} of \mathbf{Top}_0 has equalizers if it has equalizers in the
23 sense of category theory. Specifically, for any continuous mappings $f, g : X \rightarrow Y$ in \mathbf{K} , the set
24 $\{x \in X : f(x) = g(x)\}$ equipped with the subspace topology of X belongs to \mathbf{K} .

25 **Lemma 3.15.** Let $X \in \mathbf{Top}_0$ and $E \subseteq X$. Then, the following statements are equivalent:

- 26
27 (1) E is b -closed in X ;
28 (2) there exist continuous mappings $f, g : X \rightarrow (\Sigma 2)^M$ for some set M such that $E = \{x \in X : f(x) =$
29 $g(x)\}$;
30 (3) there exist continuous mappings $f, g : X \rightarrow Y$ for some $Y \in \mathbf{Top}_0$ such that $E = \{x \in X : f(x) =$
31 $g(x)\}$.
32

33 **Proof.** (1) \Rightarrow (2): Since E is b -closed, by Remark 2.3(4), we have that

34
35

$$E = \bigcap_{i \in M} (U_i \cup (X \setminus V_i)),$$

36 where $U_i, V_i \in \mathcal{O}(X)$ for all $i \in M$. Define $f, g : X \rightarrow (\Sigma 2)^M$ by

37
38

$$f(x)(i) = \chi_{U_i}(x) \quad \text{and} \quad g(x)(i) = \chi_{U_i \cup V_i}(x)$$

39 for any $x \in X$ and $i \in M$. It is easy to verify that both f and g are continuous, and for each $x \in X$,
40 $f(x)(i) = g(x)(i)$ iff $x \in U_i \cup (X \setminus V_i)$ for all $i \in M$. It follows that $E = \{x \in X : f(x) = g(x)\}$.
41

42 (2) \Rightarrow (3): It is clear.

1 (3) \Rightarrow (1): Let $x \notin E$. That is, $f(x) \neq g(x)$. Since Y is T_0 , we may assume $f(x) \not\leq g(x)$ without
 2 loss of generality. Then, there exists $V \in \mathcal{O}(Y)$ such that $f(x) \in V$ and $g(x) \notin V$. It follows that
 3 $x \in f^{-1}(V)$ and $x \notin g^{-1}(V)$. We claim that $E \cap f^{-1}(V) \cap \downarrow x = \emptyset$. In fact, if $y \in \downarrow x \cap f^{-1}(V) \cap E$,
 4 then $g(y) = f(y) \in V$ and $g(y) \leq g(x)$, and hence $g(x) \in V$, a contradiction. This shows that E is
 5 b -closed. \square

6 By Lemma 3.15, the following proposition is clear.

7 **Proposition 3.16.** Let \mathbf{K} be a full subcategory of \mathbf{Top}_0 . If $\{(\Sigma 2)^M : M \text{ is a set}\} \subseteq \mathbf{K}$, then the following
 8 statements are equivalent:
 9

- 10 (1) \mathbf{K} has equalizers;
- 11 (2) \mathbf{K} is b -closed-hereditary.

12 As an immediate result of Theorem 3.13 and Proposition 3.16, we have the following theorem.

13 **Theorem 3.17** ([8, 9.33 and 10.2.1]). Let \mathbf{K} be a full subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. Then,
 14 the following statements are equivalent:
 15

- 16 (1) \mathbf{K} is reflective;
- 17 (2) \mathbf{K} is productive and has equalizers.

18 **Theorem 3.18.** Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$, and Z a sober space.
 19 Then, the following statements hold.

- 20 (1) If $\{X_i : i \in I\} \subseteq \mathbf{K}$ is a family of subspaces of Z , then the subspace $\bigcap_{i \in I} X_i$ of Z belongs to \mathbf{K} .
- 21 (2) For each subspace X of Z , the inclusion mapping $e^k : X \rightarrow \text{cl}_k(X)$ is a \mathbf{K} -reflection for X , where
 22 $\text{cl}_k(X) = \bigcap \{A \in \mathbf{K} : X \subseteq A \subseteq Z\}$.

23 **Proof.** (1) We prove this in a few steps.

24 *Step 1:* Let $X = \bigcap_{i \in I} X_i$. Then, by Theorem 2.10(2), the inclusion mapping $e^s : X \rightarrow X^s$ is a sober
 25 reflection for X , where $X^s = \text{cl}_b(X)$ is the b -closure of X in Z . Assume $\mu_X : X \rightarrow X^k$ is a \mathbf{K} -reflection
 26 for X . By Theorem 3.6(3), $X^s \in \mathbf{K}$, and thus there exists a unique continuous mapping $f : X^k \rightarrow X^s$
 27 such that $f \circ \mu_X = e^s$:
 28


$$\begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow e^s & \downarrow f \\ & & X^s. \end{array}$$

33 *Step 2:* We prove that $f(X^k) = X$. Note that $X = e^s(X) = f(\mu_X(X)) \subseteq f(X^k)$. It remains to prove
 34 that $f(X^k) \subseteq X$ for each $i \in I$.

- 35 (c1) Let $e^i : X \rightarrow X_i$ be the inclusion mapping (note that X is a subspace of X_i). Since $X_i \in \mathbf{K}$, there
 36 exists a unique continuous mapping $f_i : X^k \rightarrow X_i$ such that $f_i \circ \mu_X = e^i$:
 37

$$\begin{array}{ccc} X & \xrightarrow{\mu_X} & X^k \\ & \searrow e^i & \downarrow f_i \\ & & X_i. \end{array}$$


1 (c2) Let $(X_i)^s = \text{cl}_b(X_i)$ be the b -closure of X_i in Z , which belongs to \mathbf{K} by Theorem 3.6(3). Let
 2 $e^{is} : X \rightarrow (X_i)^s$ be the inclusion mapping (note that X is a subspace of X_i and X_i is a subspace of
 3 $(X_i)^s$). Then, there exists a unique continuous mapping $g : X^k \rightarrow (X_i)^s$ such that $g \circ \mu_X = e^{is}$:



9 (c3) Let $e_i^{is} : X_i \rightarrow (X_i)^s$ be the inclusion mapping. Then, for each $x \in X$, by (c1) and (c2), we have
 10 that

11
$$(g \circ \mu_X)(x) \stackrel{\text{(c2)}}{=} e^{is}(x) = x = e^i(x) \stackrel{\text{(c1)}}{=} (f_i \circ \mu_X)(x) = ((e_i^{is} \circ f_i) \circ \mu_X)(x).$$


12 Hence, $g \circ \mu_X = (e_i^{is} \circ f_i) \circ \mu_X$. By the uniqueness of g , we deduce that $g = e_i^{is} \circ f_i$, i.e., the
 13 following diagram commutes:
 14



15 (c4) Let $e_s^{is} : X^s \rightarrow (X_i)^s$ be the inclusion mapping (by noting that $X^s = \text{cl}_b(X) \subseteq \text{cl}_b(X_i) = (X_i)^s$).
 16 Then, for each $x \in X$, we have that

17
$$(g \circ \mu_X)(x) \stackrel{\text{(c2)}}{=} e^{is}(x) = x = e^s(x) \stackrel{\text{Step 1}}{=} (f \circ \mu_X)(x) = ((e_s^{is} \circ f) \circ \mu_X)(x).$$

18 Hence, $g \circ \mu_X = (e_s^{is} \circ f) \circ \mu_X$. By the uniqueness of g , we deduce that $g = e_s^{is} \circ f$, i.e., the
 19 following diagram commutes:
 20



21 (c5) For each $y \in X^k$, we have that

22
$$f(y) = e_s^{is}(f(y)) \stackrel{\text{(c4)}}{=} g(y) \stackrel{\text{(c3)}}{=} e_i^{is}(f_i(y)) = f_i(y) \in X_i.$$

1 Thus, $f(X^k) \subseteq X_i$.

2 Therefore, $f(X^k) = \bigcap_{i \in I} X_i = X$.

3 *Step 3:* Now we have proved that the codomain of $f : X^k \rightarrow X^s$ is X , that is, $f(X^k) = X$. Then,
 4 we define the co-restriction $\hat{f} : X^k \rightarrow f(X^k) = X$ of f , which is a continuous mapping such that
 5 $\hat{f} \circ \mu_X = \text{id}_X$, the identity mapping on X . From Theorem 2.12, μ_X is a b -dense embedding, which
 6 implies that X is a b -retract of X^k ; then by Proposition 2.6, X is homeomorphic to $X^k \in \mathbf{K}$. Therefore,
 7 $X = \bigcap_{i \in I} X_i \in \mathbf{K}$.

8
 9 (2) We prove the conclusion in the following steps.


10 *Step 1:* Suppose that $\mu_X : X \rightarrow X^k$ is a \mathbf{K} -reflection for X . Applying result (1), we have that
 11 $\text{cl}_k(X) \in \mathbf{K}$. Thus, there exists a unique continuous mapping $f : X^k \rightarrow \text{cl}_k(X)$ such that $f \circ \mu_X = e^k$:



18 *Step 2:* Suppose that $\eta_{X^k} : X^k \rightarrow Y$ is a sober reflection for X^k . Let $X^s = \text{cl}_b(X)$ be the b -closure of
 19 X in Z . Then, $X^s \in \mathbf{Sob} \subseteq \mathbf{K}$ by Theorem 3.6(3), which implies that $\text{cl}_k(X) \subseteq X^s$. Let $e_k^s : \text{cl}_k(X) \rightarrow X^s$
 20 be the inclusion mapping. Then, there exists a unique continuous mapping $h : Y \rightarrow X^s$ such that
 21 $h \circ \eta_{X^k} = e_k^s \circ f$:



28 *Step 3:* Let $e^s = e_k^s \circ e^k : X \rightarrow X^s$ be the inclusion mapping. Using results of Step 2 and Step 3,
 29 we have that $h \circ \eta_{X^k} \circ \mu_X = (e_k^s \circ f) \circ \mu_X = e_k^s \circ (f \circ \mu_X) = e_k^s \circ e^k = e^s$:




39 By Theorems 2.11 and 2.12, both μ_X and η_{X^k} are b -dense embeddings, so is their composition
 40 $\eta_{X^k} \circ \mu_X : X \rightarrow Y$. Then, by Theorem 2.11, $\eta_{X^k} \circ \mu_X$ is a sober reflection for X , and by Theorem 2.10,
 41 the inclusion mapping $e^s = e_k^s \circ e^k : X \rightarrow X^s$ is also a sober reflection for X . Applying Lemma 2.8,
 42 we deduce that h is a homeomorphism.

1 *Step 4:* We prove that $f(X^k) = h(\eta_{X^k}(X^k)) = \text{cl}_k(X)$. On the one hand, by Step 3, it is clear that
 2 $X = e^s(X) = h(\eta_{X^k}(\mu_X(X))) \subseteq h(\eta_{X^k}(X^k)) \subseteq X^s \subseteq Z$. On the other hand, since η_{X^k} is an embedding
 3 and h is a homeomorphism, $h(\eta_{X^k}(X^k))$ is homeomorphic to $X^k \in \mathbf{K}$, so $h(\eta_{X^k}(X^k)) \in \mathbf{K}$. Recall that
 4 $\text{cl}_k(X) = \bigcap \{K \in \mathbf{K} : X \subseteq K \subseteq Z\}$, so we have that

5
$$\text{cl}_k(X) \subseteq h(\eta_{X^k}(X^k)) \stackrel{\text{Step 2}}{=} e_k^s(f(X^k)) = f(X^k) \subseteq \text{cl}_k(X).$$

6 Therefore, $f(X^k) = h(\eta_{X^k}(X^k)) = \text{cl}_k(X)$.

7 *Step 5:* Let $\hat{f} : X^k \rightarrow \text{cl}_k(X)$ be the co-restriction of $h \circ \eta_{X^k}$, i.e., $\hat{f}(y) = h(\eta_{X^k}(y))$ for any $y \in X^k$.
 8 Then, \hat{f} is a homeomorphism, since $h \circ \eta_{X^k}$ is a topological embedding. In addition, by Step 4, it
 9 satisfies that $\hat{f}(\mu_X(x)) = h(\eta_{X^k}(\mu_X(x))) = f(\mu_X(x))$ for each $x \in X$. Hence, $\hat{f} \circ \mu_X = f \circ \mu_X$. By the
 10 uniqueness of f , we deduce that $f = \hat{f}$ is a homeomorphism:
 11




18 Therefore, by Lemma 2.8, $e^k : X \rightarrow \text{cl}_k(X)$ is also a \mathbf{K} -reflection for X . □

19 **Theorem 3.19.** Let \mathbf{K} be a reflective subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. If $f : X \rightarrow Y$ is a
 20 continuous mapping from a sober space X to a sober space Y , then for any subspace Y_1 of Y , $Y_1 \in \mathbf{K}$
 21 implies that the subspace $f^{-1}(Y_1)$ of X belongs to \mathbf{K} .
 22

23 **Proof.** Let $X_1 = f^{-1}(Y_1)$ and $(X_1)^k = \bigcap \{K \in \mathbf{K} : X_1 \subseteq K \subseteq X\}$ be the subspace of X . By Theorem 3.18,
 24 the inclusion mapping $e_1 : X_1 \rightarrow (X_1)^k$ is a \mathbf{K} -reflection for X_1 . Consider the restriction $f_1 : X_1 \rightarrow Y_1$
 25 ($x \mapsto f(x)$) of f , then there exists a unique continuous mapping $g_1 : (X_1)^k \rightarrow Y_1$ such that $g_1 \circ e_1 = f_1$:
 26



32 Consider the composition $e_{Y_1} \circ f_1 : X_1 \rightarrow Y$, where $e_{Y_1} : Y_1 \rightarrow Y$ is the inclusion mapping. Since
 33 Y is a sober space, by Theorem 3.6(3), $Y \in \mathbf{K}$. Then, there exists a unique continuous mapping
 34 $g_2 : (X_1)^k \rightarrow Y$ such that $g_2 \circ e_1 = e_{Y_1} \circ f_1$:



41 Let $f_2 : (X_1)^k \rightarrow Y$ ($x \mapsto f(x)$) be the restriction of f . On the one hand, for each $x \in X_1$, we have
 42 $(f_2 \circ e_1)(x) = f(x) = (e_{Y_1} \circ f_1)(x) = (g_2 \circ e_1)(x)$, it follows that $f_2 \circ e_1 = g_2 \circ e_1$, which implies $g_2 = f_2$

1 by the uniqueness of g_2 . On the other hand, $g_2 \circ e_1 = e_{Y_1} \circ f_1 = e_{Y_1} \circ (g_1 \circ e_1) = (e_{Y_1} \circ g_1) \circ e_1$, which
 2 implies that $e_{Y_1} \circ g_1 = g_2 = f_2$ by the uniqueness of g_2 , i.e., the following diagram commutes:

3
4
5
6
7
8

$$\begin{array}{ccc}
 X_1 & \xrightarrow{e_1} & (X_1)^k \\
 & \searrow f_1 & \downarrow g_1 \\
 & & Y_1 \xrightarrow{e_{Y_1}} Y
 \end{array}
 \begin{array}{l}
 \text{---} g_2 = f_2 \text{---} \\
 \text{---} e_{Y_1} \text{---}
 \end{array}$$

9 Then for each $x \in (X_1)^k$, we have $f(x) = f_2(x) = g_2(x) = (e_{Y_1} \circ g_1)(x) = g_1(x) \in Y_1$, which implies
 10 $x \in f^{-1}(Y_1) = X_1$. Hence, $(X_1)^k \subseteq X_1$, and so $X_1 = (X_1)^k \in \mathbf{K}$. \square

11 Using Theorems 3.6, 3.18 and 3.19, and Keimel and Lawson's result in [12], we obtain the main
 12 result in this paper.
 13

14 **Theorem 3.20.** *Let \mathbf{K} be a full subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. Then, the following statements*
 15 *are equivalent:*

- 16 (1) \mathbf{K} is reflective;
 17 (2) \mathbf{K} satisfies conditions (K1)–(K4).
 18

19 By Theorems 3.13, 3.17 and 3.20, several equivalent conditions for the reflectivity of \mathbf{K} are
 20 summarized as follows.

21 **Theorem 3.21.** *Let \mathbf{K} be a full subcategory of \mathbf{Top}_0 such that $\mathbf{K} \not\subseteq \mathbf{Top}_1$. Then, the following statements*
 22 *are equivalent:*

- 23 (1) \mathbf{K} is reflective in \mathbf{Top}_0 ;
 24 (2) \mathbf{K} satisfies conditions (K1)–(K4);
 25 (3) \mathbf{K} is productive and *b-closed-hereditary*;
 26 (4) \mathbf{K} is productive and has equalizers.
 27

28 **Remark 3.22.** In the paper [5], Ershov proved that \mathbf{K} is reflective in \mathbf{Top}_0 if and only if \mathbf{K} satisfies
 29 conditions (K1)–(K4) for every wide category \mathbf{K} , where a *wide category* \mathbf{K} is a full subcategory of
 30 \mathbf{Top}_0 such that every T_0 space X can be topologically embedded into some space Y belonging to \mathbf{K} .
 31 By Corollary 3.4, it is clear that every wide category \mathbf{K} satisfies $\mathbf{K} \not\subseteq \mathbf{Top}_1$. However, the converse is
 32 not true. For example, the full subcategory **Sier** of \mathbf{Top}_0 , consisting of all topological spaces that are
 33 homeomorphic to Σ_2 , satisfies **Sier** $\not\subseteq \mathbf{Top}_1$ but is not a wide category. Consequently, Ershov's result
 34 can be regarded as a corollary of Theorem 3.21. Furthermore, the condition $\mathbf{K} \not\subseteq \mathbf{Top}_1$ of Theorem 3.21
 35 is a common and easily checkable condition in domain theory. Additionally, the approach presented in
 36 this paper differs significantly from that employed in [5].
 37

38 4. Some applications

39
 40 By using the results in the last section, we investigate the reflectivity of several categories of T_0 spaces,
 41 including co-sober spaces, strong d -spaces, k -bounded sober spaces, and consonant T_0 spaces. It is
 42 worth noting that all these classes of spaces are closed under the formation of homeomorphic objects.

1 **4.1. Co-sober spaces.** In order to study the dual Hofmann-Mislove Theorem, Escardó, Lawson and
 2 Simpson [4] introduced the co-sober spaces [4], which are defined below.

3 **Definition 4.1** ([4]). Let X be a T_0 space, and Q a nonempty compact saturated subset of X .

4 (1) Q is called *k-irreducible* if for any compact saturated subsets Q_1, Q_2 of X , $Q = Q_1 \cup Q_2$ implies
 5 $Q = Q_1$ or $Q = Q_2$.

6 (2) X is called *co-sober* if for each *k-irreducible* set Q , there exists a unique $x \in X$ such that $Q = \uparrow x$.

7
 8 For a poset P , the family of all upper sets of P forms a topology, called the *Alexandroff topology* on
 9 P [7].

10 **Lemma 4.2.** (1) *Every poset equipped with the Alexandroff topology is co-sober.*

11 (2) *A poset equipped with the Alexandroff topology is sober if and only if the poset is a dcpo. Hence,*
 12 *co-sober spaces need not be sober.*

13 **Proof.** Let P be a poset equipped with the Alexandroff topology.

14 (1) Note that every nonempty compact saturated set in P is of the form $\uparrow F$, where F is a finite subset
 15 of P . Thus, every *k-irreducible* compact saturated set is of the form $\uparrow x$, where $x \in P$. Therefore, P is
 16 co-sober.
 17

18 (2) This follows immediately from the fact that the irreducible subsets of P are exactly the directed
 19 sets. □

20 Let **Co-Sob** be the full subcategory of **Top₀** consisting of all co-sober spaces.

21 It is worth noting that the topology of the Sierpiński space Σ_2 coincides with the Alexandroff
 22 topology on the two-point chain $2 = \{0, 1\}$. Thus, by Lemma 4.2, Σ_2 is co-sober, and since it is not
 23 T_1 , we can conclude that **Co-Sob** $\not\subseteq$ **Top₁**. The question of whether every sober space is co-sober was
 24 raised in [4]. A negative answer was given by Wen and Xu in [21], where they proved that Isbell's
 25 complete lattice (see [11]) equipped with the lower topology is sober but not co-sober. Furthermore,
 26 it has been proved in [18] that there exists a dcpo that is sober but not co-sober with respect to the
 27 Scott topology. Therefore, we have that **Sob** $\not\subseteq$ **Co-Sob**. Then, by applying Theorem 3.6(3), we obtain
 28 the following result.
 29

30 **Corollary 4.3.** *The category Co-Sob is not reflective in Top₀.*

31 **4.2. Strong d-spaces.** The class of strong *d-spaces* was introduced by Xu and Zhao [25], which lies
 32 between the classes of T_1 spaces and *d-spaces*.

33 **Definition 4.4** ([25]). A T_0 space X is called a *strong d-space* if for any $x \in X$, any directed subset D
 34 of X , and any open subset U of X , $\bigcap_{d \in D} \uparrow d \cap \uparrow x \subseteq U$ implies $\uparrow d_0 \cap \uparrow x \subseteq U$ for some $d_0 \in D$.

35 Let **SD** be the full subcategory of **Top₀** consisting of all strong *d-spaces*.

36 In [25, Example 3.34], it was shown that there exists a continuous dcpo P whose Scott topology
 37 is not a strong *d-space*. However, it is well-known that the Scott topology on any continuous dcpo
 38 is always sober. In addition, it has been noted in [25, Remark 3.21] that the Scott topology on every
 39 continuous lattice is a strong *d-space*. Therefore, **Sob** $\not\subseteq$ **SD** and **SD** $\not\subseteq$ **Top₁**. By applying Theorem
 40 3.6(3), we deduce the following result.
 41

42 **Corollary 4.5.** *The category SD is not reflective in Top₀.*

1 **4.3. k -bounded sober spaces.** In [26], Zhao and Ho introduced another weaker notion of sobriety,
2 called k -bounded sobriety. This notion is defined as follows.

3 **Definition 4.6** ([26]). A T_0 space X is k -bounded sober if for any irreducible closed subset F of X
4 with $\bigvee F$ existing, there is a unique point $x \in X$ such that $F = \downarrow x$.

5 Let **KSob** be the full subcategory of **Top₀** consisting of all k -bounded sober spaces. It is clear that
6 **Sob** \subseteq **KSob** and **Sob** $\not\subseteq$ **Top₁**. Thus, we conclude that **KSob** $\not\subseteq$ **Top₁**.

8 **Example 4.7.** Let $X = [0, 3]$ equipped with the Scott topology (i.e., the open sets are \emptyset , $[0, 3]$ and all
9 sets of the form $(x, 3]$, where $x \in [0, 3]$). Since $[0, 3]$ is a continuous lattice, we know that X is a sober
10 space, and hence it is also k -bounded sober. For each integer $n \geq 2$, let $X_n = [0, 1) \cup (2 - \frac{1}{n}, 2 + \frac{1}{n})$. We
11 have the following facts.

12 (1) Each subspace X_n of X is k -bounded sober. To show this, let F be an irreducible closed set in X_n
13 and $x \in X_n$ such that $\bigvee_{X_n} F = x$. There are two cases:

14 (c1) $x \in [0, 1)$. Then, $F \subseteq \downarrow x \subseteq [0, 1)$, which follows that $\text{cl}_{X_n}(F) = \text{cl}_{X_n}(\{x\})$.

15 (c2) $x \in (2 - \frac{1}{n}, 2 + \frac{1}{n})$. Then, $F \cap (2 - \frac{1}{n}, 2 + \frac{1}{n}) \neq \emptyset$, which implies that

$$16 \quad x = \bigvee_{X_n} F = \bigvee_{X_n} F \cap (2 - \frac{1}{n}, 2 + \frac{1}{n}) = \bigvee_X F \cap (2 - \frac{1}{n}, 2 + \frac{1}{n}) = \bigvee_X F \cap X_n = \bigvee_X F.$$

17 Since X is sober, we have $\text{cl}_X(F) = \text{cl}_X(\{x\})$, and thus $\text{cl}_{X_n}(F) = \text{cl}_X(F) \cap X_n = \text{cl}_X(\{x\}) \cap X_n =$
18 $\text{cl}_{X_n}(\{x\})$, where the last equality holds because $x \in X_n$.

19 All these show that X_n is k -bounded sober.

20 (2) The intersection $Y = \bigcap_{n \geq 2} X_n = [0, 1) \cup \{2\}$ equipped with the subspace topology of X is not
21 k -bounded sober. In fact, the set $F := [0, 1)$ is irreducible since it is directed, and $\bigvee_Y F = 2$. In
22 addition, since $[0, 1]$ is a closed set in X and $F = [0, 1] \cap Y$, we have that F is a closed set in Y . For
23 each $x \in [0, 1)$, we have that $\text{cl}_Y(\{x\}) = [0, x] \neq F$, and $\text{cl}_Y(\{2\}) = Y \neq F$. Therefore, Y is not a
24 k -bounded sober space.

25 The above example shows that **KSob** does not satisfy (K3). Thus, by Theorem 3.21, we obtain the
26 following corollary.

27 **Corollary 4.8** ([14]). *The category **KSob** is not reflective in **Top₀**.*

28 **4.4. Consonant spaces.** The class of consonant spaces was introduced by Dolecki, Greco and Lechicki
29 in [2], which plays an important role in discussion of the equality of the Isbell topology and the compact-
30 open topology on function spaces [20]. The definition is given as follows.

31 **Definition 4.9** ([2]). A topological space X is called *consonant* if for every Scott open subset \mathcal{U} of
32 $\mathcal{O}(X)$, there exists a family $\{K_i : i \in I\}$ of compact subsets of X such that $\mathcal{U} = \bigcup_{i \in I} \mathcal{N}(K_i)$, where
33 $\mathcal{N}(K_i) := \{U \in \mathcal{O}(X) : K_i \subseteq U\}$ for all $i \in I$.

34 Let **Const** be the full subcategory of **Top₀** consisting of all consonant T_0 spaces. We note the
35 following facts:

36 (1) Every finite topological space X is consonant, since every subset of X is compact. As a consequence,
37 $\Sigma 2$ is a consonant T_0 but non- T_1 space, so we have that **Const** $\not\subseteq$ **Top₁**;

38 (2) Nogura and Shakhmatov [20] have shown that there exists a metric space (hence is sober) that is
39 not consonant, so we have that **Sob** $\not\subseteq$ **Const**.

1 Therefore, by Theorem 3.6(3), we obtain the following corollary.

2 **Corollary 4.10.** *The category \mathbf{Const} is not reflective in \mathbf{Top}_0 .*

5. Conclusion

6 In this paper we proved that if a reflective subcategory of \mathbf{Top}_0 contains a non- T_1 space and satisfies
7 the (K2) condition proposed by Lawson and Keimel, then it also satisfies the remaining conditions
8 (K1), (K3) and (K4). Based on this result, we concluded that several subcategories are not reflective,
9 thus giving negative answers to some open problems. We expect that this result might also serve as a
10 tool for verifying the reflectivity of other subcategories of \mathbf{Top}_0 .

11
12
13 **Acknowledgments.** (1) We appreciate the referee for giving us many valuable suggestions to
14 improve our work, particularly the new proof of Theorem 3.18. (2) This work was supported by
15 the National Natural Science Foundation of China (No. 1210010153, 12071188), Jiangsu Provincial
16 Department of Education (No. 21KJB110008), and International Postdoctoral Exchange Fellowship
17 Program (No. PC2022013).

References

- 20 [1] J. Adámek, H. Herrlich, G. E. Strecker, Abstract and Concrete Categories -The Joy of Cats, Dover Publications, 2009.
21 [2] S. Dolecki, G.H. Greco, A. Lechicki, When do the upper Kuratowski topology (homeomorphically, Scott topology) and
22 the co-compact topology coincide?, Trans. Am. Math. Soc. 347 (1995) 2869–2884.
23 [3] C. H. Dowker, D. Papert, Quotient frames and subspaces, Proc. London Math. Soc., 3 (16) (1966) 275–296.
24 [4] M. Escardó, J. Lawson, A. Simpson, Comparing Cartesian closed categories of (core) compactly generated spaces,
25 Topol. Appl. 143 (2004) 105–145.
26 [5] Yu. L. Ershov, K-Completions of T_0 -Spaces, Algebra and Logic 61(4) (2022) 177–187.
27 [6] G. Gierz, K. Hofmann, K. Keimel, J. Lawson, M. Mislove, D. Scott, Continuous lattices and Domains, Encyclopedia of
28 Mathematics and Its Applications, Vol.93, Cambridge University Press, 2003.
29 [7] J. Goubault-Larrecq, Non-Hausdorff topology and Domain Theory, Cambridge University Press, 2013.
30 [8] H. Herrlich, Topologische Reflexionen und Coreflexionen, Lecture Notes Math. Vol. 78, Berlin, 1968.
31 [9] R-E. Hoffmann, Topological functors admitting generalized Cauchy-completions, in Categorical Topology, Proc. of the
32 conf. held at Mannheim 1975 ed. E. Binz and H. Herrlich, pp. 286–344, Lect. Notes in Math., 540, Berlin-Heidelberg-
33 New York: Springer 1976.
34 [10] R-E. Hoffmann, On the sobrification remainder ${}^sX - X$, Pacific J. Math. 83(1) (1979) 145–156.
35 [11] J.R. Isbell, Completion of a construction of Johnstone, Proc. Amer. Math. Soc. 85 (1982) 333–334.
36 [12] K. Keimel, J.D. Lawson, D -completions and the d -topology, Ann. Pure Appl. Logic 159 (2009) 292–306.
37 [13] J.K. Kelly, General Toplogy, Springer-Verlag, 1955.
38 [14] J. Lu, K. Wang, G. Wu, B. Zhao, Nonexistence of k -bounded sobrification, preprint, 2020, arXiv: 2011.11606v1.
39 [15] S. Mac Lane, Categories for the Working Mathematician, Springer, 1997.
40 [16] L.D. Nel, R.G. Wilson, Epireflections in the category of T_0 -spaces, Fund. Math. 75 (1972) 69–74.
41 [17] L. Skula, On a reflective subcategory of the category of all topological spaces, Trans. Amer. Math. Soc. 142 (1969)
42 37–41.
[18] C. Shen, G. Wu, X. Xi, D. Zhao, Sober Scott spaces are not always co-sober, Topol. Appl. 282 (2020) 107316.
[19] C. Shen, X. Xi, D. Zhao, The non-reflectivity of open well-filtered spaces via b -topology, Houston J. Math. (2023) accepted.

- 1 [20] T. Nogura, D. Shakhmatov, When does the Fell topology on a hyperspace of closed sets coincide with the meet of the
 2 upper Kuratowski and the lower Vietoris topologies?, *Topol. Appl.* 70 (2–3) (1996) 213–243.
 3 [21] X.P. Wen and X.Q. Xu, Sober is not always co-sober, *Topol. Appl.* 250 (2018) 48–52.
 4 [22] G. Wu, X. Xi, X. Xu and D. Zhao, Existence of well-filterification, *Topol. Appl.* 267 (2019) 107044.
 5 [23] O. Wyler, Dedekind complete posets and Scott topologies, in: B. Banaschewski, R. E. Hoffman (Eds.), *Continuous
 6 Lattices, Proc. Bremen, 1979, Lecture Notes in Mathematics, vol. 871, 1981, pp. 384–389.*
 7 [24] X. Xu, D. Zhao, Some open problems on well-filtered spaces and sober spaces, *Topol. Appl.* (2020) 107540.
 8 [25] X. Xu, D. Zhao, On topological Rudin’s lemma, well-filtered spaces and sober spaces, *Topol. Appl.* 272 (2020) 107080.
 9 [26] D. Zhao, W. Ho, On topologies defined by irreducible sets, *J. Log. Algebr. Methods Program.* 84 (1) (2015) 185–195.

10 CHONG SHEN, KEY LABORATORY OF MATHEMATICS AND INFORMATION NETWORKS, SCHOOL OF SCIENCE,
 11 BEIJING UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, BEIJING, CHINA 100876
 12 *Email address:* shenchong0520@163.com

13 XIAOYONG XI, SCHOOL OF MATHEMATICS AND STATISTICS, YANCHENG TEACHERS UNIVERSITY, JIANGSU,
 14 YANCHENG, CHINA 224002
 15 *Email address:* xixy@yctu.edu.cn

16 DONGSHENG ZHAO, MATHEMATICS AND MATHEMATICS EDUCATION, NATIONAL INSTITUTE OF EDUCATION,
 17 NANYANG TECHNOLOGICAL UNIVERSITY, 1 NANYANG WALK, SINGAPORE 637616
 18 *Email address:* dongsheng.zhao@nie.edu.sg

19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42